



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 21, 2023 – 12:40 AM EDT

PDB ID : 5J0P
Title : Binary complex crystal structure of DNA polymerase Beta with A:C mismatch at the primer terminus
Authors : Batra, V.K.; Wilson, S.H.
Deposited on : 2016-03-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	
2	T	16	
3	P	10	
4	D	5	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	330	2640	1667	461	503	9	0	0	0

- Molecule 2 is a DNA chain called Template Strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	16	321	153	63	90	15	0	0	0

- Molecule 3 is a DNA chain called Primer Strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	P	10	203	97	38	59	9	0	0	0

- Molecule 4 is a DNA chain called Downstream Primer Strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	5	106	49	20	32	5	0	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	181	Total	O	0	0
			181	181		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	T	46	Total O 46 46	0	0
6	P	21	Total O 21 21	0	0
6	D	13	Total O 13 13	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase beta



- Molecule 2: Template Strand



- Molecule 3: Primer Strand



- Molecule 4: Downstream Primer Strand



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.14Å 79.62Å 54.57Å 90.00° 105.55° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 20.88 – 2.20	Depositor EDS
% Data completeness (in resolution range)	81.2 (50.00-2.20) 81.4 (20.88-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.19Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.249 , 0.320 0.253 , 0.320	Depositor DCC
R_{free} test set	2184 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtrriage
Anisotropy	0.369	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.056 for l,-k,h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3533	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2690	0.57	0/3617
2	T	0.42	0/360	0.84	0/552
3	P	0.49	0/227	0.76	0/349
4	D	0.91	1/118 (0.8%)	0.87	0/179
All	All	0.41	1/3395 (0.0%)	0.64	0/4697

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	DG	OP3-P	-7.24	1.52	1.61

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	2	DT	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2640	0	2650	104	0
2	T	321	0	179	5	0
3	P	203	0	114	8	0
4	D	106	0	57	0	0
5	A	2	0	0	0	0
6	A	181	0	0	7	0
6	D	13	0	0	0	0
6	P	21	0	0	2	0
6	T	46	0	0	1	0
All	All	3533	0	3000	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ARG:HH12	1:A:275:SER:HB3	1.32	0.94
1:A:209:LYS:HA	1:A:212:HIS:HB3	1.51	0.90
1:A:209:LYS:HG2	1:A:213:GLN:HG3	1.56	0.88
1:A:201:THR:H	1:A:204:SER:HB3	1.38	0.87
2:T:6:DA:H1'	2:T:7:DA:H5'	1.65	0.77
3:P:10:DC:H2''	6:P:111:HOH:O	1.82	0.77
1:A:183:ARG:HH12	1:A:275:SER:CB	2.00	0.72
1:A:174:ILE:HD12	1:A:175:ALA:N	2.05	0.72
1:A:234:LYS:HD3	6:T:136:HOH:O	1.89	0.72
1:A:275:SER:HB2	6:A:521:HOH:O	1.90	0.71
1:A:183:ARG:NH1	1:A:275:SER:HB3	2.05	0.70
1:A:296:TYR:OH	2:T:8:DC:H5''	1.92	0.69
3:P:9:DG:H2''	3:P:10:DC:H5''	1.75	0.68
1:A:174:ILE:HG23	1:A:196:THR:HG23	1.76	0.66
1:A:183:ARG:HH11	1:A:275:SER:N	1.94	0.66
1:A:274:GLY:HA2	1:A:279:ASN:ND2	2.10	0.66
1:A:158:MET:O	1:A:162:VAL:HG23	1.95	0.65
1:A:104:SER:OG	1:A:135:HIS:HE1	1.81	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ARG:NH1	1:A:275:SER:N	2.46	0.64
1:A:48:LYS:C	1:A:48:LYS:HD2	2.18	0.64
1:A:212:HIS:HD2	1:A:230:LYS:NZ	2.00	0.59
3:P:1:DG:H2'	3:P:2:DC:C6	2.37	0.59
1:A:174:ILE:HG23	1:A:196:THR:CG2	2.32	0.59
1:A:278:PHE:HA	1:A:335:GLU:O	2.04	0.57
1:A:212:HIS:HB2	6:A:596:HOH:O	2.03	0.57
1:A:311:LEU:HD13	1:A:322:TYR:CE1	2.40	0.57
1:A:7:PRO:HB3	1:A:11:LEU:HD12	1.88	0.56
1:A:292:THR:CG2	1:A:299:ARG:HB2	2.36	0.56
1:A:244:LYS:HB2	1:A:247:GLU:OE1	2.06	0.55
1:A:13:GLY:HA2	1:A:16:THR:OG1	2.06	0.55
1:A:245:ASN:HD22	1:A:245:ASN:N	2.05	0.55
1:A:150:ILE:HG12	1:A:253:ARG:HD2	1.88	0.55
1:A:9:GLU:HG3	1:A:13:GLY:HA3	1.88	0.55
1:A:183:ARG:HH12	1:A:275:SER:CA	2.20	0.54
1:A:212:HIS:HD2	1:A:230:LYS:HZ2	1.57	0.53
2:T:6:DA:C1'	2:T:7:DA:H5'	2.37	0.53
1:A:183:ARG:NH1	1:A:275:SER:CA	2.72	0.53
1:A:135:HIS:HD2	6:A:584:HOH:O	1.92	0.52
1:A:110:ALA:O	1:A:113:LYS:HG2	2.10	0.52
1:A:328:ARG:HD3	1:A:332:ASP:O	2.10	0.52
1:A:205:THR:C	1:A:207:GLN:H	2.12	0.52
1:A:158:MET:HG3	1:A:241:LEU:HD11	1.93	0.51
1:A:31:GLN:HG2	1:A:108:PRO:HB2	1.92	0.51
1:A:102:ARG:HD2	6:A:554:HOH:O	2.12	0.50
1:A:195:LEU:HD21	1:A:214:VAL:HG21	1.92	0.50
3:P:9:DG:H3'	6:P:102:HOH:O	2.12	0.50
1:A:274:GLY:HA2	1:A:279:ASN:HD21	1.76	0.50
1:A:165:GLU:CD	1:A:221:VAL:HG11	2.32	0.49
1:A:218:LEU:HB3	1:A:224:ILE:HG13	1.93	0.49
1:A:262:LYS:HD2	1:A:262:LYS:O	2.12	0.49
1:A:174:ILE:O	1:A:195:LEU:HD12	2.13	0.49
1:A:45:VAL:O	1:A:48:LYS:HE3	2.13	0.49
1:A:49:TYR:CE1	1:A:51:HIS:HB2	2.47	0.49
1:A:75:GLU:O	1:A:79:THR:HG23	2.13	0.48
1:A:123:GLU:CD	1:A:123:GLU:H	2.17	0.48
1:A:84:LYS:O	1:A:88:ILE:HG13	2.13	0.48
1:A:261:PRO:HD2	1:A:264:GLN:NE2	2.27	0.48
1:A:200:PHE:CE2	1:A:261:PRO:HG3	2.48	0.48
1:A:130:ASP:HA	1:A:137:ARG:NH2	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:CYS:HB2	6:A:573:HOH:O	2.11	0.48
1:A:262:LYS:HZ1	1:A:314:ASP:HA	1.78	0.48
1:A:229:SER:OG	1:A:236:MET:HB2	2.13	0.48
1:A:31:GLN:HG2	1:A:108:PRO:CB	2.43	0.47
1:A:218:LEU:HB2	1:A:224:ILE:HD12	1.97	0.47
3:P:9:DG:C2'	3:P:10:DC:C5'	2.93	0.47
1:A:182:ARG:NH2	1:A:316:GLU:OE2	2.42	0.47
1:A:207:GLN:HB3	6:A:576:HOH:O	2.13	0.47
1:A:285:HIS:HD2	1:A:323:ILE:HD12	1.80	0.47
1:A:285:HIS:O	1:A:288:GLU:HG2	2.15	0.47
1:A:327:TYR:C	1:A:327:TYR:CD1	2.89	0.47
1:A:323:ILE:HG13	1:A:325:TRP:HB2	1.96	0.46
1:A:138:ILE:HD13	1:A:226:ASP:HB3	1.97	0.46
1:A:206:LYS:O	1:A:207:GLN:C	2.54	0.46
1:A:328:ARG:NH1	1:A:335:GLU:OXT	2.48	0.46
1:A:231:GLY:HA3	2:T:10:DC:OP1	2.16	0.46
1:A:126:ARG:NE	6:A:504:HOH:O	2.39	0.46
1:A:209:LYS:O	1:A:213:GLN:HB2	2.16	0.46
1:A:174:ILE:CG2	1:A:196:THR:HG23	2.46	0.46
1:A:103:VAL:HB	1:A:106:ILE:HD12	1.98	0.45
1:A:266:TYR:CD2	1:A:315:SER:HA	2.51	0.45
1:A:75:GLU:OE2	1:A:83:ARG:HG3	2.17	0.45
1:A:183:ARG:NH1	1:A:275:SER:CB	2.73	0.45
1:A:165:GLU:HG3	1:A:221:VAL:HG21	2.00	0.44
1:A:152:ARG:HG3	1:A:152:ARG:NH1	2.31	0.44
1:A:197:HIS:ND1	1:A:198:PRO:HD2	2.33	0.43
3:P:9:DG:H2''	3:P:10:DC:C5'	2.44	0.43
1:A:217:GLN:HE21	1:A:221:VAL:HG13	1.83	0.43
1:A:298:ILE:HG23	1:A:298:ILE:O	2.19	0.43
1:A:33:ILE:O	1:A:36:TYR:HB3	2.18	0.43
1:A:158:MET:HA	1:A:241:LEU:HD21	1.99	0.43
1:A:173:TYR:OH	1:A:210:LEU:HA	2.19	0.43
1:A:327:TYR:C	1:A:327:TYR:HD1	2.23	0.43
1:A:262:LYS:NZ	1:A:314:ASP:HA	2.34	0.42
2:T:6:DA:H2''	2:T:7:DA:OP2	2.19	0.42
1:A:152:ARG:HG3	1:A:152:ARG:HH11	1.84	0.42
1:A:159:GLN:HG3	1:A:177:VAL:HG21	2.01	0.42
1:A:9:GLU:O	1:A:10:THR:C	2.58	0.42
1:A:133:ASN:O	1:A:137:ARG:HG3	2.20	0.42
3:P:2:DC:C6	3:P:3:DT:H72	2.54	0.42
1:A:251:PRO:HG2	1:A:253:ARG:CZ	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:SER:O	1:A:279:ASN:ND2	2.53	0.41
1:A:320:PHE:CD2	1:A:327:TYR:HA	2.55	0.41
1:A:48:LYS:C	1:A:48:LYS:CD	2.86	0.41
3:P:2:DC:H2'	3:P:3:DT:C7	2.51	0.41
1:A:245:ASN:HD22	1:A:245:ASN:H	1.67	0.41
1:A:281:ASN:ND2	1:A:335:GLU:O	2.53	0.41
1:A:292:THR:O	1:A:298:ILE:HA	2.20	0.41
1:A:162:VAL:O	1:A:166:VAL:HG23	2.21	0.40
1:A:195:LEU:CD2	1:A:214:VAL:HG21	2.52	0.40
1:A:205:THR:O	1:A:207:GLN:N	2.54	0.40
1:A:174:ILE:HD12	1:A:175:ALA:H	1.82	0.40
1:A:195:LEU:HG	1:A:196:THR:N	2.36	0.40
1:A:279:ASN:O	1:A:282:MET:HB3	2.22	0.40
1:A:15:ILE:O	1:A:18:MET:HB3	2.21	0.40
1:A:113:LYS:CG	1:A:114:PHE:N	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	328/335 (98%)	289 (88%)	34 (10%)	5 (2%)	10 8

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	206	LYS
1	A	265	TYR
1	A	130	ASP
1	A	207	GLN
1	A	208	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	289/295 (98%)	280 (97%)	9 (3%)	40 51

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	40	ARG
1	A	48	LYS
1	A	174	ILE
1	A	216	GLU
1	A	258	ARG
1	A	267	CYS
1	A	325	TRP
1	A	327	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	135	HIS
1	A	159	GLN
1	A	207	GLN
1	A	212	HIS
1	A	217	GLN
1	A	222	HIS
1	A	245	ASN
1	A	252	HIS
1	A	264	GLN
1	A	279	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/335 (98%)	0.76	43 (13%) 3 3	18, 42, 80, 94	0
2	T	16/16 (100%)	0.22	0 100 100	26, 41, 64, 65	0
3	P	10/10 (100%)	0.40	1 (10%) 7 6	23, 32, 50, 64	0
4	D	5/5 (100%)	-0.31	0 100 100	25, 26, 37, 39	0
All	All	361/366 (98%)	0.71	44 (12%) 4 3	18, 42, 79, 94	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	284	ALA	7.6
1	A	287	LEU	7.0
1	A	246	ASP	7.0
1	A	205	THR	6.1
1	A	208	PRO	6.0
1	A	247	GLU	4.5
1	A	244	LYS	4.5
1	A	204	SER	4.4
1	A	325	TRP	4.2
1	A	270	LEU	3.9
1	A	245	ASN	3.8
1	A	206	LYS	3.5
1	A	326	LYS	3.4
1	A	282	MET	3.4
1	A	301	LEU	3.1
3	P	10	DC	3.1
1	A	277	ILE	3.0
1	A	304	THR	3.0
1	A	78	ALA	2.9
1	A	335	GLU	2.8
1	A	203	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	201	THR	2.4
1	A	321	ASP	2.4
1	A	320	PHE	2.4
1	A	257	ILE	2.4
1	A	22	LEU	2.4
1	A	10	THR	2.4
1	A	7	PRO	2.4
1	A	324	GLN	2.4
1	A	259	LEU	2.3
1	A	248	LYS	2.3
1	A	281	ASN	2.3
1	A	293	ILE	2.3
1	A	269	VAL	2.2
1	A	312	PRO	2.2
1	A	306	VAL	2.2
1	A	303	VAL	2.2
1	A	299	ARG	2.1
1	A	239	CYS	2.1
1	A	286	ALA	2.1
1	A	266	TYR	2.1
1	A	211	LEU	2.0
1	A	267	CYS	2.0
1	A	292	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

5	NA	A	402	1/1	0.93	0.17	44,44,44,44	0
5	NA	A	401	1/1	0.96	0.17	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.