



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2023 – 04:00 PM EST

PDB ID : 1J2F
Title : X-ray crystal structure of IRF-3 and its functional implications
Authors : Takahasi, K.; Noda, N.; Horiuchi, M.; Mori, M.; Okabe, Y.; Fukuhara, Y.;
Terasawa, H.; Fujita, T.; Inagaki, F.
Deposited on : 2003-01-04
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	 3% 61% 22% 12%
1	B	258	 3% 69% 17% 10%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon regulatory factor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	226	1757	1126	301	321	9	0	0	0
1	B	231	1811	1158	310	333	10	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	170	GLY	-	cloning artifact	UNP Q14653
A	171	ALA	-	cloning artifact	UNP Q14653
A	172	MET	-	cloning artifact	UNP Q14653
A	173	GLY	-	cloning artifact	UNP Q14653
A	174	SER	-	cloning artifact	UNP Q14653
B	170	GLY	-	cloning artifact	UNP Q14653
B	171	ALA	-	cloning artifact	UNP Q14653
B	172	MET	-	cloning artifact	UNP Q14653
B	173	GLY	-	cloning artifact	UNP Q14653
B	174	SER	-	cloning artifact	UNP Q14653


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	75	Total 75	O 75	0	0
2	B	85	Total 85	O 85	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interferon regulatory factor 3



- Molecule 1: Interferon regulatory factor 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.81Å 134.81Å 69.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.02 – 2.30 61.66 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (62.02-2.30) 94.1 (61.66-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.00Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.217 , 0.242 0.218 , 0.242	Depositor DCC
R_{free} test set	2677 reflections (6.55%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3728	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1808	0.58	0/2468
1	B	0.36	0/1863	0.59	0/2540
All	All	0.35	0/3671	0.59	0/5008

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1757	0	1690	55	0
1	B	1811	0	1752	43	0
2	A	75	0	0	4	0
2	B	85	0	0	4	0
All	All	3728	0	3442	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303:SER:O	1:A:305:HIS:N	2.01	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:326:ILE:HD13	1:B:415:LEU:HD13	1.54	0.89
1:A:338:ARG:HD2	1:A:338:ARG:H	1.38	0.88
1:A:326:ILE:HD13	1:A:415:LEU:HD13	1.59	0.83
1:B:254:ASP:HB3	2:B:42:HOH:O	1.86	0.76
1:B:297:GLU:OE2	1:B:341:ARG:HD3	1.88	0.74
1:A:299:LEU:HD21	1:B:387:LEU:HB2	1.69	0.73
1:A:227:ARG:CZ	1:A:238:LEU:HD13	2.20	0.71
1:A:285:ARG:HH21	1:A:288:HIS:HA	1.57	0.69
1:A:352:TRP:NE1	1:A:354:GLN:HG3	2.09	0.68
1:A:326:ILE:HD12	1:A:419:MET:CE	2.26	0.66
1:A:212:GLY:O	1:B:388:GLU:HG3	1.95	0.65
1:B:331:THR:HG22	1:B:336:SER:HB2	1.78	0.64
1:B:352:TRP:NE1	1:B:354:GLN:HG3	2.13	0.63
1:A:212:GLY:HA2	1:B:387:LEU:HD23	1.80	0.62
1:B:338:ARG:HH11	1:B:338:ARG:HB3	1.65	0.61
1:B:401:LEU:HD22	1:B:403:LEU:HG	1.82	0.60
1:B:277:ALA:HB3	1:B:282:TRP:HE1	1.67	0.59
1:A:387:LEU:O	1:A:404:THR:HA	2.03	0.58
1:B:277:ALA:HB3	1:B:282:TRP:NE1	2.20	0.57
1:A:329:LEU:O	1:A:333:THR:HG23	2.05	0.57
1:A:345:TRP:CZ2	1:A:366:LYS:HD3	2.40	0.57
1:B:326:ILE:CD1	1:B:415:LEU:HD13	2.31	0.56
1:B:281:LEU:HB2	1:B:320:PHE:HB3	1.87	0.56
1:A:218:GLN:NE2	1:A:220:ILE:HD11	2.21	0.55
1:A:381:VAL:HG12	1:A:381:VAL:O	2.06	0.54
1:A:324:PRO:O	1:A:327:VAL:HG22	2.08	0.54
1:B:361:ARG:HB3	1:B:364:MET:CE	2.37	0.54
1:A:198:PRO:HG2	2:A:133:HOH:O	2.07	0.54
1:B:330:ILE:O	1:B:334:GLU:HG3	2.07	0.54
1:A:197:VAL:HG13	1:A:200:GLU:HG3	1.90	0.53
1:A:218:GLN:CD	1:A:220:ILE:HD11	2.29	0.52
1:A:380:ARG:C	1:A:382:GLY:H	2.13	0.52
1:B:322:LEU:HD22	1:B:326:ILE:CD1	2.40	0.52
1:B:232:GLU:HA	2:B:39:HOH:O	2.10	0.51
1:B:293:TRP:CE3	1:B:319:VAL:HG22	2.46	0.51
1:A:276:ARG:HD3	2:A:17:HOH:O	2.11	0.50
1:A:371:CYS:HB3	1:A:393:LEU:O	2.13	0.49
1:A:254:ASP:OD2	1:A:257:VAL:HG23	2.13	0.49
1:A:404:THR:OG1	1:A:407:GLN:HG3	2.13	0.49
1:A:338:ARG:HD2	1:A:338:ARG:N	2.18	0.49
1:A:220:ILE:N	1:A:220:ILE:HD12	2.28	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:206:VAL:O	1:B:217:GLN:HA	2.13	0.49
1:A:196:LEU:O	1:A:276:ARG:NH2	2.46	0.48
1:B:338:ARG:HB3	1:B:338:ARG:NH1	2.28	0.48
1:B:332:PHE:HA	1:B:336:SER:HB3	1.95	0.48
1:B:403:LEU:HB3	1:B:407:GLN:HG3	1.95	0.48
1:A:415:LEU:HD22	1:A:419:MET:HE2	1.96	0.48
1:B:258:MET:O	1:B:262:ARG:HG3	2.14	0.48
1:A:303:SER:C	1:A:305:HIS:H	2.09	0.48
1:B:361:ARG:HB3	1:B:364:MET:HE3	1.95	0.48
1:A:326:ILE:HD12	1:A:419:MET:HE1	1.94	0.48
1:A:373:ARG:O	1:A:376:VAL:HG22	2.13	0.47
1:A:338:ARG:H	1:A:338:ARG:CD	2.15	0.47
1:A:360:LYS:HD3	1:A:360:LYS:H	1.79	0.47
1:B:231:SER:OG	1:B:233:VAL:HG23	2.15	0.47
1:B:326:ILE:HD12	1:B:419:MET:CE	2.45	0.47
1:A:360:LYS:HE3	2:A:141:HOH:O	2.15	0.46
1:B:388:GLU:HG2	1:B:404:THR:HG22	1.98	0.46
1:B:191:PRO:HG3	1:B:401:LEU:HD11	1.98	0.45
1:B:389:ASN:ND2	1:B:390:THR:H	2.15	0.45
1:A:327:VAL:HG23	1:A:328:ASP:N	2.30	0.44
1:B:247:ASP:OD2	1:B:262:ARG:NH1	2.47	0.44
1:A:285:ARG:HD2	1:A:289:CYS:O	2.17	0.44
1:B:371:CYS:HB3	1:B:393:LEU:O	2.17	0.44
1:A:387:LEU:C	1:A:404:THR:HG22	2.38	0.44
1:A:352:TRP:HE1	1:A:354:GLN:HG3	1.83	0.43
1:B:293:TRP:HA	1:B:345:TRP:O	2.18	0.43
1:A:255:ARG:HG3	2:A:1:HOH:O	2.19	0.43
1:A:226:LEU:HD12	1:A:226:LEU:C	2.39	0.43
1:B:274:LEU:CD2	1:B:367:VAL:HG11	2.48	0.43
1:A:238:LEU:HD12	1:A:238:LEU:N	2.32	0.43
1:B:338:ARG:NH1	1:B:338:ARG:CB	2.82	0.43
1:A:281:LEU:HB2	1:A:320:PHE:HB3	2.01	0.43
1:B:315:LYS:HD2	1:B:315:LYS:C	2.39	0.43
1:A:293:TRP:HA	1:A:345:TRP:O	2.19	0.42
1:A:203:GLU:HB3	1:A:370:THR:OG1	2.19	0.42
1:B:274:LEU:HA	1:B:282:TRP:O	2.20	0.42
1:A:213:ARG:HG3	2:B:19:HOH:O	2.18	0.42
1:A:360:LYS:HB2	1:A:362:LEU:HD13	2.01	0.42
1:A:248:PRO:HB2	1:A:258:MET:HG2	2.02	0.42
1:B:315:LYS:HD2	1:B:315:LYS:O	2.19	0.42
1:A:300:LEU:HD23	1:A:301:PRO:HD2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:LYS:HB2	1:A:360:LYS:HE2	1.83	0.41
1:B:363:VAL:O	1:B:364:MET:HE2	2.20	0.41
1:B:262:ARG:HG2	1:B:262:ARG:HH11	1.84	0.41
1:A:224:GLU:HG2	1:A:239:PRO:HB2	2.02	0.41
1:A:322:LEU:HG	1:A:326:ILE:CD1	2.50	0.41
1:B:404:THR:OG1	1:B:407:GLN:HG2	2.20	0.41
1:B:194:ARG:HG2	2:B:30:HOH:O	2.21	0.41
1:A:272:LEU:C	1:A:272:LEU:HD13	2.40	0.41
1:A:260:TYR:O	1:A:263:HIS:HB2	2.20	0.40
1:A:326:ILE:O	1:A:330:ILE:HG13	2.22	0.40
1:B:194:ARG:O	1:B:200:GLU:HG3	2.20	0.40
1:A:258:MET:O	1:A:262:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/258 (86%)	212 (96%)	8 (4%)	2 (1%)	17	20
1	B	227/258 (88%)	216 (95%)	10 (4%)	1 (0%)	34	42
All	All	449/516 (87%)	428 (95%)	18 (4%)	3 (1%)	22	26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	GLY
1	A	381	VAL
1	B	304	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/217 (86%)	172 (92%)	15 (8%)	12	15
1	B	196/217 (90%)	186 (95%)	10 (5%)	24	33
All	All	383/434 (88%)	358 (94%)	25 (6%)	17	23

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	LYS
1	A	194	ARG
1	A	197	VAL
1	A	238	LEU
1	A	281	LEU
1	A	293	TRP
1	A	297	GLU
1	A	299	LEU
1	A	300	LEU
1	A	338	ARG
1	A	360	LYS
1	A	362	LEU
1	A	386	SER
1	A	401	LEU
1	A	415	LEU
1	B	194	ARG
1	B	281	LEU
1	B	315	LYS
1	B	319	VAL
1	B	322	LEU
1	B	375	LEU
1	B	377	GLU
1	B	389	ASN
1	B	401	LEU
1	B	415	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	214	GLN
1	A	284	GLN
1	A	389	ASN
1	B	217	GLN
1	B	284	GLN
1	B	389	ASN
1	B	413	GLN
1	B	422	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/258 (87%)	0.07	8 (3%) 44 51	19, 30, 60, 72	0
1	B	231/258 (89%)	0.04	9 (3%) 39 46	16, 27, 53, 74	0
All	All	457/516 (88%)	0.06	17 (3%) 41 48	16, 29, 58, 74	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	249	GLY	4.6
1	A	255	ARG	3.7
1	B	250	MET	3.6
1	B	252	LEU	3.4
1	A	381	VAL	3.4
1	B	253	THR	3.3
1	A	253	THR	3.1
1	A	384	ALA	3.1
1	A	252	LEU	2.9
1	B	255	ARG	2.9
1	B	233	VAL	2.8
1	A	385	SER	2.6
1	B	303	SER	2.6
1	A	383	GLY	2.4
1	B	232	GLU	2.3
1	A	250	MET	2.2
1	B	254	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.