



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 03:27 PM JST

PDB ID : 5J51
Title : Structure of tetrameric jacalin complexed with Gal alpha-(1,4) Gal
Authors : Abhinav, K.V.; Sharma, K.; Surolia, A.; Vijayan, M.
Deposited on : 2016-04-01
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	95% 5%
1	C	133	3% 92% 8%
1	E	133	% 94% 6%
1	G	133	% 93% 7%
2	B	19	63% 16% 21%
2	D	19	11% 53% 21% 5% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	19	
2	H	19	
3	I	2	
3	J	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	C	204	-	-	X	-
4	EDO	E	202	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Agglutinin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	133	1069	698	166	203	2	0	4	0
1	C	133	1067	696	164	205	2	0	4	0
1	E	133	1062	692	164	204	2	0	3	0
1	G	133	1047	683	161	201	2	0	1	0

- Molecule 2 is a protein called Agglutinin beta-3 chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	15	105	68	18	19	0	0	0
2	D	15	101	65	17	19	0	0	0
2	F	15	103	67	18	18	0	0	0
2	H	15	99	64	17	18	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-galactopyranose-(1-4)-alpha-D-galactopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	I	2	23	12	11	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	J	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0

- Molecule 5 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	3	1		
5	A	1	Total	C	O	0	0
			4	3	1		
5	A	1	Total	C	O	0	0
			4	3	1		
5	B	1	Total	C	O	0	0
			4	3	1		
5	C	1	Total	C	O	0	0
			4	3	1		
5	E	1	Total	C	O	0	0
			4	3	1		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C O	0	0
			7	4 3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	119	Total	O	0	0
			119	119		
7	B	11	Total	O	0	0
			11	11		
7	C	92	Total	O	0	0
			92	92		
7	D	12	Total	O	0	0
			12	12		
7	E	100	Total	O	0	0
			100	100		
7	F	9	Total	O	0	0
			9	9		
7	G	105	Total	O	0	0
			105	105		
7	H	18	Total	O	0	0
			18	18		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Agglutinin alpha chain

Chain A:  95% 5%





- Molecule 1: Agglutinin alpha chain

Chain C:  3% 92% 8%




- Molecule 1: Agglutinin alpha chain

Chain E:  % 94% 6%



- Molecule 1: Agglutinin alpha chain

Chain G:  % 93% 7%




- Molecule 2: Agglutinin beta-3 chain

Chain B:  63% 16% 21%




- Molecule 2: Agglutinin beta-3 chain




- Molecule 2: Agglutinin beta-3 chain



- Molecule 2: Agglutinin beta-3 chain



- Molecule 3: alpha-D-galactopyranose-(1-4)-alpha-D-galactopyranose



- Molecule 3: alpha-D-galactopyranose-(1-4)-alpha-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.56Å 81.09Å 63.25Å 90.00° 108.05° 90.00°	Depositor
Resolution (Å)	35.71 – 1.67 33.62 – 1.67	Depositor EDS
% Data completeness (in resolution range)	96.7 (35.71-1.67) 96.7 (33.62-1.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 1.67Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.164 , 0.213 0.163 , 0.211	Depositor DCC
R_{free} test set	3223 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	15.5	Xtrriage
Anisotropy	0.495	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 56.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5308	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, IPA, PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.10	0/1098	1.03	2/1487 (0.1%)
1	C	0.98	2/1096 (0.2%)	0.94	0/1487
1	E	1.00	0/1091	0.98	1/1478 (0.1%)
1	G	1.05	0/1076	0.98	0/1459
2	B	1.27	2/107 (1.9%)	1.06	0/145
2	D	1.21	1/103 (1.0%)	0.97	0/141
2	F	1.34	0/105	1.15	0/142
2	H	1.30	1/101 (1.0%)	0.84	0/138
All	All	1.06	6/4777 (0.1%)	0.98	3/6477 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	15	TRP	CD2-CE2	6.22	1.48	1.41
1	C	123	TRP	CD2-CE2	5.67	1.48	1.41
2	H	15	TRP	CD2-CE2	5.48	1.48	1.41
2	D	15	TRP	CD2-CE2	5.24	1.47	1.41
1	C	64	TYR	CE1-CZ	5.15	1.45	1.38
2	B	7	SER	CB-OG	-5.01	1.35	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59	ASP	CB-CG-OD1	6.46	124.11	118.30
1	A	27	ASP	CB-CG-OD1	6.20	123.88	118.30
1	E	13	ARG	NE-CZ-NH1	-5.30	117.65	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1037	5	0
1	C	1067	0	1036	19	0
1	E	1062	0	1031	10	0
1	G	1047	0	1015	5	0
2	B	105	0	108	1	0
2	D	101	0	97	9	0
2	F	103	0	103	1	0
2	H	99	0	92	2	0
3	I	23	0	21	2	0
3	J	23	0	21	0	0
4	A	12	0	18	0	0
4	C	40	0	60	5	0
4	E	44	0	66	4	0
4	G	16	0	24	3	0
5	A	12	0	24	4	0
5	B	4	0	8	0	0
5	C	4	0	8	0	0
5	E	4	0	8	2	0
6	A	7	0	10	2	0
7	A	119	0	0	2	0
7	B	11	0	0	0	0
7	C	92	0	0	0	0
7	D	12	0	0	0	0
7	E	100	0	0	1	0
7	F	9	0	0	0	0
7	G	105	0	0	1	0
7	H	18	0	0	0	0
All	All	5308	0	4787	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:106[B]:LEU:HD12	2:D:15:TRP:CH2	2.08	0.89
1:A:13:ARG:HH12	5:A:206:IPA:H32	1.37	0.86
1:C:106[B]:LEU:HD12	2:D:15:TRP:HH2	1.45	0.80
1:C:106[B]:LEU:HD11	2:D:12:VAL:HG21	1.67	0.75
1:E:78:TYR:CE1	3:I:1:GLA:H2	2.22	0.73
1:C:60:PHE:CZ	2:H:6:ILE:HG12	2.28	0.69
1:C:106[B]:LEU:C	1:C:106[B]:LEU:HD23	2.13	0.69
1:E:119:SER:OG	4:E:202:EDO:H22	1.93	0.68
1:C:106[B]:LEU:HD23	1:C:106[B]:LEU:O	1.94	0.67
1:C:5:ASP:H	4:C:204:EDO:C2	2.08	0.66
1:E:64:TYR:CE1	5:E:212:IPA:H33	2.30	0.66
6:A:207:PEG:H41	7:A:393:HOH:O	1.95	0.66
1:C:106[B]:LEU:CD1	2:D:12:VAL:HG21	2.29	0.63
1:C:5:ASP:H	4:C:204:EDO:H21	1.63	0.62
1:A:13:ARG:NH1	5:A:206:IPA:H32	2.14	0.59
2:D:16:GLY:O	2:D:17:ALA:HB2	2.05	0.57
1:E:1:GLY:O	4:E:202:EDO:H21	2.05	0.56
1:E:3:ALA:HB2	4:E:202:EDO:H12	1.88	0.55
1:C:4:PHE:HB2	4:C:204:EDO:H22	1.87	0.55
2:D:3:GLN:N	4:G:204:EDO:HO1	2.08	0.52
1:C:60:PHE:HZ	2:H:6:ILE:HG12	1.76	0.50
6:A:207:PEG:H22	7:A:345:HOH:O	2.11	0.50
1:C:106[B]:LEU:CD1	2:D:15:TRP:CH2	2.91	0.50
1:E:111:GLY:O	5:E:212:IPA:H32	2.12	0.49
1:E:119:SER:OG	4:E:202:EDO:C2	2.60	0.49
1:A:13:ARG:HH12	5:A:206:IPA:C3	2.17	0.48
1:C:5:ASP:O	4:C:204:EDO:H21	2.13	0.48
5:A:204:IPA:H2	5:A:205:IPA:C1	2.44	0.48
1:G:8:ALA:O	4:G:201:EDO:H22	2.13	0.48
2:B:18:LYS:HD3	2:B:18:LYS:HA	1.74	0.47
1:G:49:THR:HB	7:G:319:HOH:O	2.16	0.46
1:G:1:GLY:HA3	1:G:125:ASP:OD2	2.16	0.45
2:D:16:GLY:O	2:D:17:ALA:CB	2.64	0.45
1:G:12:ILE:HD11	1:G:113:ILE:HG22	1.98	0.45
1:C:106[B]:LEU:CD1	2:D:12:VAL:CG2	2.96	0.44
1:C:5:ASP:C	4:C:204:EDO:H21	2.37	0.44
1:A:19:TYR:HE1	1:A:54[A]:VAL:CG2	2.31	0.44
1:C:106[B]:LEU:O	1:C:106[B]:LEU:CD2	2.65	0.44
1:E:78:TYR:CE1	3:I:1:GLA:C2	2.98	0.43
1:E:49[A]:THR:HG22	7:E:318:HOH:O	2.18	0.43
2:F:18:LYS:O	2:F:18:LYS:HG2	2.19	0.43
1:C:10:THR:HG23	4:G:201:EDO:H21	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:PHE:HB2	1:A:44:HIS:CE1	2.54	0.42
1:G:71:TYR:CE1	1:G:99:THR:HB	2.55	0.42
1:C:14:GLU:HB3	1:C:31:VAL:HB	2.03	0.41
1:C:43:ASN:ND2	1:C:45:LYS:HD3	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/133 (102%)	131 (97%)	4 (3%)	0	100	100
1	C	135/133 (102%)	130 (96%)	5 (4%)	0	100	100
1	E	134/133 (101%)	131 (98%)	3 (2%)	0	100	100
1	G	132/133 (99%)	128 (97%)	4 (3%)	0	100	100
2	B	13/19 (68%)	13 (100%)	0	0	100	100
2	D	13/19 (68%)	13 (100%)	0	0	100	100
2	F	13/19 (68%)	13 (100%)	0	0	100	100
2	H	13/19 (68%)	13 (100%)	0	0	100	100
All	All	588/608 (97%)	572 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/113 (102%)	115 (100%)	0	100	100
1	C	117/113 (104%)	117 (100%)	0	100	100
1	E	116/113 (103%)	116 (100%)	0	100	100
1	G	114/113 (101%)	113 (99%)	1 (1%)	78	69
2	B	11/15 (73%)	11 (100%)	0	100	100
2	D	10/15 (67%)	10 (100%)	0	100	100
2	F	10/15 (67%)	9 (90%)	1 (10%)	7	1
2	H	9/15 (60%)	8 (89%)	1 (11%)	6	1
All	All	502/512 (98%)	499 (99%)	3 (1%)	86	79

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	4	SER
1	G	74	ASN
2	H	14	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GLA	I	1	3	12,12,12	1.57	3 (25%)	17,17,17	1.26	2 (11%)
3	GLA	I	2	3	11,11,12	3.99	5 (45%)	15,15,17	1.85	4 (26%)
3	GLA	J	1	3	12,12,12	1.41	2 (16%)	17,17,17	1.03	1 (5%)
3	GLA	J	2	3	11,11,12	1.60	3 (27%)	15,15,17	1.66	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLA	I	1	3	-	2/2/22/22	0/1/1/1
3	GLA	I	2	3	-	1/2/19/22	0/1/1/1
3	GLA	J	1	3	-	0/2/22/22	0/1/1/1
3	GLA	J	2	3	-	1/2/19/22	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	2	GLA	O5-C1	10.21	1.60	1.43
3	I	2	GLA	C4-C5	5.98	1.65	1.53
3	I	2	GLA	C1-C2	3.72	1.60	1.52
3	J	2	GLA	O5-C1	3.16	1.48	1.43
3	J	2	GLA	C2-C3	3.14	1.57	1.52
3	I	1	GLA	C4-C3	2.99	1.59	1.52
3	J	1	GLA	O5-C5	2.92	1.51	1.44
3	I	1	GLA	O1-C1	2.47	1.47	1.39
3	I	2	GLA	O4-C4	2.36	1.48	1.43
3	J	1	GLA	C4-C3	2.25	1.58	1.52
3	I	1	GLA	O5-C1	2.19	1.48	1.42
3	J	2	GLA	C4-C5	2.04	1.57	1.53
3	I	2	GLA	C4-C3	2.03	1.57	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	2	GLA	O5-C5-C6	4.44	114.17	107.20
3	I	2	GLA	O4-C4-C5	3.50	118.00	109.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	2	GLA	O5-C1-C2	3.21	115.72	110.77
3	I	2	GLA	O5-C5-C6	2.60	111.28	107.20
3	I	2	GLA	O2-C2-C3	-2.54	105.04	110.14
3	J	2	GLA	O4-C4-C3	-2.43	104.73	110.35
3	J	1	GLA	C3-C4-C5	-2.17	106.38	110.24
3	I	1	GLA	O1-C1-O5	2.04	116.51	110.38
3	I	1	GLA	O4-C4-C3	2.01	115.00	110.35

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	I	1	GLA	C4-C5-C6-O6
3	I	1	GLA	O5-C5-C6-O6
3	I	2	GLA	C4-C5-C6-O6
3	J	2	GLA	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	GLA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	204	-	3,3,3	0.43	0	2,2,2	0.20	0
4	EDO	A	201	-	3,3,3	0.36	0	2,2,2	0.26	0
4	EDO	A	202	-	3,3,3	0.32	0	2,2,2	1.07	0
4	EDO	A	203	-	3,3,3	0.50	0	2,2,2	0.42	0
4	EDO	C	203	-	3,3,3	0.54	0	2,2,2	0.61	0
4	EDO	E	209	-	3,3,3	0.45	0	2,2,2	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PEG	A	207	-	6,6,6	0.30	0	5,5,5	1.02	0
4	EDO	C	209	-	3,3,3	0.51	0	2,2,2	0.09	0
4	EDO	E	202	-	3,3,3	0.56	0	2,2,2	0.63	0
4	EDO	G	204	-	3,3,3	0.39	0	2,2,2	0.45	0
4	EDO	E	205	-	3,3,3	0.58	0	2,2,2	0.30	0
4	EDO	C	206	-	3,3,3	0.40	0	2,2,2	0.45	0
4	EDO	C	207	-	3,3,3	0.39	0	2,2,2	0.62	0
4	EDO	E	207	-	3,3,3	0.37	0	2,2,2	0.54	0
4	EDO	C	208	-	3,3,3	0.57	0	2,2,2	0.17	0
4	EDO	E	211	-	3,3,3	0.88	0	2,2,2	0.71	0
4	EDO	E	210	-	3,3,3	0.52	0	2,2,2	0.43	0
4	EDO	C	202	-	3,3,3	0.45	0	2,2,2	0.51	0
5	IPA	B	101	-	3,3,3	0.67	0	3,3,3	0.36	0
4	EDO	C	205	-	3,3,3	0.81	0	2,2,2	0.34	0
5	IPA	A	204	-	3,3,3	0.58	0	3,3,3	0.46	0
5	IPA	E	212	-	3,3,3	0.51	0	3,3,3	1.00	0
4	EDO	E	206	-	3,3,3	0.87	0	2,2,2	0.84	0
4	EDO	G	201	-	3,3,3	0.64	0	2,2,2	0.55	0
5	IPA	A	206	-	3,3,3	0.48	0	3,3,3	0.18	0
5	IPA	C	211	-	3,3,3	0.61	0	3,3,3	0.30	0
4	EDO	E	208	-	3,3,3	0.40	0	2,2,2	0.24	0
4	EDO	C	210	-	3,3,3	0.32	0	2,2,2	0.53	0
4	EDO	G	202	-	3,3,3	0.64	0	2,2,2	0.13	0
4	EDO	E	204	-	3,3,3	0.70	0	2,2,2	0.77	0
4	EDO	G	203	-	3,3,3	0.35	0	2,2,2	1.14	0
4	EDO	C	201	-	3,3,3	0.38	0	2,2,2	0.78	0
4	EDO	E	201	-	3,3,3	0.31	0	2,2,2	1.51	0
4	EDO	E	203	-	3,3,3	0.31	0	2,2,2	0.60	0
5	IPA	A	205	-	3,3,3	0.42	0	3,3,3	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	204	-	-	1/1/1/1	-
4	EDO	A	201	-	-	0/1/1/1	-
4	EDO	A	202	-	-	1/1/1/1	-
4	EDO	A	203	-	-	1/1/1/1	-
4	EDO	C	203	-	-	1/1/1/1	-
4	EDO	E	209	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	A	207	-	-	3/4/4/4	-
4	EDO	C	209	-	-	0/1/1/1	-
4	EDO	E	202	-	-	1/1/1/1	-
4	EDO	G	204	-	-	1/1/1/1	-
4	EDO	E	205	-	-	0/1/1/1	-
4	EDO	C	206	-	-	0/1/1/1	-
4	EDO	C	207	-	-	1/1/1/1	-
4	EDO	E	207	-	-	1/1/1/1	-
4	EDO	C	208	-	-	0/1/1/1	-
4	EDO	E	211	-	-	1/1/1/1	-
4	EDO	E	210	-	-	0/1/1/1	-
4	EDO	C	202	-	-	1/1/1/1	-
4	EDO	C	205	-	-	0/1/1/1	-
4	EDO	E	206	-	-	0/1/1/1	-
4	EDO	G	201	-	-	1/1/1/1	-
4	EDO	E	208	-	-	1/1/1/1	-
4	EDO	C	210	-	-	1/1/1/1	-
4	EDO	G	202	-	-	1/1/1/1	-
4	EDO	E	204	-	-	0/1/1/1	-
4	EDO	G	203	-	-	1/1/1/1	-
4	EDO	C	201	-	-	1/1/1/1	-
4	EDO	E	201	-	-	1/1/1/1	-
4	EDO	E	203	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	201	EDO	O1-C1-C2-O2
4	C	203	EDO	O1-C1-C2-O2
4	C	204	EDO	O1-C1-C2-O2
4	E	201	EDO	O1-C1-C2-O2
4	E	211	EDO	O1-C1-C2-O2
4	G	203	EDO	O1-C1-C2-O2
6	A	207	PEG	O1-C1-C2-O2
4	A	202	EDO	O1-C1-C2-O2
4	E	203	EDO	O1-C1-C2-O2
6	A	207	PEG	C4-C3-O2-C2
6	A	207	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	G	201	EDO	O1-C1-C2-O2
4	G	204	EDO	O1-C1-C2-O2
4	E	202	EDO	O1-C1-C2-O2
4	G	202	EDO	O1-C1-C2-O2
4	A	203	EDO	O1-C1-C2-O2
4	C	207	EDO	O1-C1-C2-O2
4	C	210	EDO	O1-C1-C2-O2
4	C	202	EDO	O1-C1-C2-O2
4	E	207	EDO	O1-C1-C2-O2
4	E	208	EDO	O1-C1-C2-O2

There are no ring outliers.

9 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	204	EDO	5	0
6	A	207	PEG	2	0
4	E	202	EDO	4	0
4	G	204	EDO	1	0
5	A	204	IPA	1	0
5	E	212	IPA	2	0
4	G	201	EDO	2	0
5	A	206	IPA	3	0
5	A	205	IPA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	133/133 (100%)	-0.45	0 100 100	9, 13, 20, 25	0
1	C	133/133 (100%)	-0.21	4 (3%) 50 53	10, 14, 22, 36	0
1	E	133/133 (100%)	-0.33	1 (0%) 86 88	9, 13, 22, 37	0
1	G	133/133 (100%)	-0.42	1 (0%) 86 88	9, 14, 23, 33	0
2	B	15/19 (78%)	-0.19	0 100 100	10, 15, 26, 38	0
2	D	15/19 (78%)	0.13	2 (13%) 3 3	10, 15, 37, 39	0
2	F	15/19 (78%)	-0.03	0 100 100	10, 14, 25, 37	0
2	H	15/19 (78%)	-0.16	0 100 100	10, 14, 24, 35	0
All	All	592/608 (97%)	-0.32	8 (1%) 75 79	9, 14, 24, 39	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	17	ALA	3.3
1	C	99	THR	3.1
2	D	3	GLN	2.5
1	C	100	SER	2.4
1	E	74	ASN	2.4
1	G	98	VAL	2.2
1	C	22	GLU	2.1
1	C	21	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLA	I	1	12/12	0.74	0.30	30,43,49,50	0
3	GLA	I	2	11/12	0.83	0.17	15,21,30,35	0
3	GLA	J	1	12/12	0.95	0.17	16,20,24,26	0
3	GLA	J	2	11/12	0.95	0.08	15,16,17,17	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	IPA	C	211	4/4	0.69	0.21	37,38,38,38	0
4	EDO	E	211	4/4	0.77	0.14	29,30,31,34	0
4	EDO	E	205	4/4	0.81	0.13	42,43,45,50	0
4	EDO	E	210	4/4	0.81	0.13	46,46,47,48	0
4	EDO	E	209	4/4	0.82	0.13	37,40,41,44	0
4	EDO	G	201	4/4	0.82	0.15	32,33,33,36	0
4	EDO	E	201	4/4	0.82	0.18	33,34,34,37	0
4	EDO	C	210	4/4	0.83	0.10	43,43,44,45	0
4	EDO	C	206	4/4	0.84	0.34	34,35,37,40	0
5	IPA	B	101	4/4	0.84	0.15	34,35,35,36	0
4	EDO	C	207	4/4	0.84	0.11	49,49,50,52	0
4	EDO	G	203	4/4	0.85	0.14	35,37,37,41	0
4	EDO	C	203	4/4	0.87	0.10	33,35,35,38	0
4	EDO	A	202	4/4	0.88	0.24	44,45,46,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	IPA	E	212	4/4	0.88	0.15	27,28,28,28	0
4	EDO	C	201	4/4	0.89	0.11	40,40,40,41	0
4	EDO	G	202	4/4	0.89	0.16	27,30,30,32	0
4	EDO	E	202	4/4	0.89	0.20	32,33,34,39	0
4	EDO	A	203	4/4	0.90	0.19	39,40,42,45	0
5	IPA	A	204	4/4	0.90	0.12	35,35,36,37	0
5	IPA	A	206	4/4	0.90	0.12	45,45,46,49	0
6	PEG	A	207	7/7	0.90	0.17	25,27,40,42	0
4	EDO	E	207	4/4	0.91	0.18	36,37,38,38	0
5	IPA	A	205	4/4	0.92	0.10	41,42,42,42	0
4	EDO	E	203	4/4	0.92	0.12	42,43,45,48	0
4	EDO	E	204	4/4	0.93	0.14	18,18,18,18	0
4	EDO	C	204	4/4	0.93	0.08	26,27,28,30	0
4	EDO	G	204	4/4	0.94	0.10	25,27,28,32	0
4	EDO	E	206	4/4	0.94	0.09	19,20,20,21	0
4	EDO	C	209	4/4	0.94	0.10	33,33,35,36	0
4	EDO	E	208	4/4	0.94	0.07	26,26,27,29	0
4	EDO	C	202	4/4	0.95	0.08	25,26,26,29	0
4	EDO	A	201	4/4	0.96	0.09	21,22,22,24	0
4	EDO	C	205	4/4	0.96	0.08	18,18,18,18	0
4	EDO	C	208	4/4	0.97	0.07	16,17,17,17	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.