Full wwPDB X-ray Structure Validation Report (i)

Sep 10, 2023 – 11:50 AM EDT

PDB ID	:	4J6P
Title	:	Crystal structure of calcium2+-free wild-type CD23 lectin domain (crystal form F)
Authors	:	Dhaliwal, B.; Yuan, D.; Sutton, B.J.
Deposited on	:	2013-02-11
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	4% 81% 8% • 10%
1	B	143	3% 85% 6% • 9%
1	C	143	6% 82% 12% • 6%
1	D	143	4% 80% 11% • 7%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Low affinity immunoglobulin epsilon Fc receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	128	1050	664	185	188	13	0	4	0
1	B	130	1048	660	185	192	11	0	0	0
1	C	135	1087	684	193	199	11	0	1	0
1	D	133	1071	672	191	197	11	0	1	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	67	67	67	0	0
2	B	83	83	83	0	0
2	C	53	53	53	0	0
2	D	72	72	72	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Low affinity immunoglobulin epsilon Fc receptor

Chain A:

• Molecule 1: Low affinity immunoglobulin epsilon Fc receptor

Chain B:

• Molecule 1: Low affinity immunoglobulin epsilon Fc receptor

Chain C:



• Molecule 1: Low affinity immunoglobulin epsilon Fc receptor

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	62.67Å 89.92Å 101.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.82 – 1.90 45.84 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (27.82-1.90) 99.9 (45.84-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.65 (at 1.89Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.0, BUSTER 2.10.0	Depositor
R, R_{free}	0.186 , 0.214 0.200 , 0.229	Depositor DCC
R_{free} test set	2312 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	27.9	Xtriage
Anisotropy	0.495	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 51.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4531	wwPDB-VP
Average B, all atoms (Å^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	A	0.51	0/1094	0.70	0/1480
1	B	0.52	0/1080	0.67	0/1462
1	C	0.52	0/1124	0.73	1/1524 (0.1%)
1	D	0.55	0/1106	0.77	3/1496 (0.2%)
All	All	0.52	0/4404	0.72	4/5962 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	250	PRO	N-CA-C	5.89	127.41	112.10
1	C	256	GLY	N-CA-C	-5.34	99.74	113.10
1	D	159	VAL	N-CA-C	-5.23	96.88	111.00
1	D	251	THR	C-N-CA	5.05	134.31	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1050	0	987	6	0
1	B	1048	0	964	5	0
1	C	1087	0	1011	11	0
1	D	1071	0	991	18	0
2	A	67	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	83	0	0	0	0
2	C	53	0	0	4	0
2	D	72	0	0	2	0
All	All	4531	0	3953	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:157:GLY:CA	1:D:158:PHE:HB2	1.42	1.43
1:D:157:GLY:HA2	1:D:158:PHE:CB	1.61	1.19
1:D:156:SER:O	1:D:172:ARG:CG	2.02	1.06
1:D:157:GLY:N	1:D:158:PHE:HB2	1.77	0.99
1:D:157:GLY:CA	1:D:158:PHE:CB	2.29	0.98
1:D:157:GLY:HA2	1:D:158:PHE:HB2	0.96	0.96
1:D:156:SER:O	1:D:172:ARG:HG3	1.65	0.95
1:D:156:SER:O	1:D:172:ARG:HG2	1.80	0.79
1:A:160:CYS:HG	1:A:288:CYS:HG	1.32	0.72
1:D:157:GLY:H	1:D:158:PHE:HB2	1.58	0.69
1:C:185[A]:VAL:HG12	2:C:304:HOH:O	1.94	0.67
1:B:160:CYS:HB2	1:B:172:ARG:HD2	1.82	0.61
1:A:185[B]:VAL:HG12	2:A:301:HOH:O	2.02	0.59
1:C:275:ARG:NH1	2:C:340:HOH:O	2.33	0.55
1:C:200:SER:HA	1:C:234:TRP:CE3	2.45	0.52
1:A:200:SER:HA	1:A:234:TRP:CE3	2.45	0.52
1:B:185:VAL:HG11	1:C:257:GLU:HG3	1.93	0.50
1:C:228:LEU:HA	1:C:255:GLN:HB3	1.94	0.50
1:B:200:SER:HA	1:B:234:TRP:CE3	2.48	0.49
1:D:275:ARG:NH1	2:D:305:HOH:O	2.39	0.47
1:C:235:VAL:O	1:D:231:GLU:OE1	2.31	0.47
1:C:188:ARG:NH1	1:D:256:GLY:HA3	2.29	0.47
1:C:160:CYS:HB2	1:C:172:ARG:HD3	1.97	0.47
1:B:224:ARG:HD2	2:C:305:HOH:O	2.15	0.46
1:C:188:ARG:HH12	1:D:256:GLY:HA3	1.84	0.43
1:A:174:CYS:SG	1:A:286:ALA:HA	2.59	0.43
1:D:228:LEU:HA	1:D:255:GLN:HB3	2.00	0.43
1:A:160:CYS:HB2	1:A:172:ARG:HG2	2.00	0.42
1:D:200:SER:HA	1:D:234:TRP:CE3	2.54	0.42
1:D:223:LEU:HB2	1:D:234:TRP:CZ3	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185[B]:VAL:HG11	1:C:189:TYR:CZ	2.55	0.42
1:D:156:SER:O	1:D:172:ARG:CD	2.65	0.42
1:B:223:LEU:HB2	1:B:234:TRP:CZ3	2.56	0.41
1:C:229:LYS:HG3	2:C:348:HOH:O	2.22	0.40
1:D:185[B]:VAL:HG12	2:D:302:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/143 (90%)	122 (95%)	6 (5%)	0	100	100
1	B	126/143 (88%)	122 (97%)	4 (3%)	0	100	100
1	C	134/143 (94%)	125 (93%)	9 (7%)	0	100	100
1	D	132/143 (92%)	122 (92%)	8 (6%)	2 (2%)	10	3
All	All	520/572 (91%)	491 (94%)	27 (5%)	2 (0%)	34	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	158	PHE
1	D	251	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/120 (94%)	107 (95%)	6 (5%)	22	13
1	B	111/120 (92%)	108 (97%)	3 (3%)	44	38
1	C	116/120 (97%)	111 (96%)	5 (4%)	29	19
1	D	114/120 (95%)	110 (96%)	4 (4%)	36	27
All	All	454/480 (95%)	436 (96%)	18 (4%)	33	22

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	208	ASP
1	A	224[A]	ARG
1	A	224[B]	ARG
1	A	231	GLU
1	A	241	ASP
1	A	288	CYS
1	B	224	ARG
1	B	231	GLU
1	B	267	ARG
1	C	159	VAL
1	C	208	ASP
1	C	229	LYS
1	C	241	ASP
1	C	254	SER
1	D	159	VAL
1	D	241	ASP
1	D	252	SER
1	D	258	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	213	HIS
1	A	239	HIS
1	D	197	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	128/143 (89%)	0.34	6 (4%)	31	34	19, 38, 64, 105	0
1	B	130/143 (90%)	0.05	5 (3%)	40	43	21, 34, 58, 86	0
1	C	135/143 (94%)	0.37	9 (6%)	17	20	22, 40, 67, 85	0
1	D	133/143 (93%)	0.34	6 (4%)	33	36	19, 36, 58, 69	0
All	All	526/572 (91%)	0.27	26 (4%)	29	33	19, 37, 64, 105	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	156	SER	10.6
1	A	291	PRO	9.5
1	B	158	PHE	9.0
1	D	158	PHE	8.6
1	A	289	THR	8.5
1	D	157	GLY	6.7
1	A	290	PRO	6.4
1	D	251	THR	5.0
1	B	290	PRO	5.0
1	C	255	GLN	4.8
1	B	216	HIS	4.6
1	C	158	PHE	4.2
1	C	252	SER	3.5
1	C	216	HIS	3.5
1	A	216	HIS	3.3
1	C	229	LYS	3.3
1	D	239	HIS	2.8
1	C	292	ALA	2.5
1	B	255	GLN	2.5
1	A	256	GLY	2.4
1	C	217	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	251	THR	2.3
1	C	228	LEU	2.3
1	A	288	CYS	2.2
1	D	252	SER	2.1
1	C	214	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.