

Nov 20, 2022 – 06:43 AM EST

PDB ID	:	3JBA
EMDB ID	:	EMD-6424
Title	:	The U4 antibody epitope on human papillomavirus 16 identified by cryo-EM
Authors	:	Guan, J.; Bywaters, S.M.; Brendle, S.A.; Lee, H.; Ashley, R.E.; Christensen,
		N.D.; Hafenstein, S.
Deposited on	:	2015-08-11
Resolution	:	12.00 Å(reported)
Based on initial model	:	3J6R

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	0.0.1. dev 43
:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	1.9.9
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.31.3
	: : : : :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 12.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	$\mathop{{\rm EM}}\limits_{(\#{ m Entries})}$		
Clashscore	158937	4297		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	L	115	91%	9%	
2	Н	115	90%	9%	•
3	А	478	92%	7%	•
3	В	478	90%	8%	•
3	С	478	89%	10%	·
3	D	478	90%	8%	•
3	Е	478	91%	8%	
3	F	478	91%	8%	•



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 20721 atoms, of which 7468 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called H16.U4 antibody light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	L	115	Total 1772	C 567	Н 879	N 144	0 178	$\frac{S}{4}$	0	0

• Molecule 2 is a protein called H16.U4 antibody heavy chain.

Mol	Chain	Residues	Atoms				AltConf	Trace		
2	Н	115	Total 1735	C 560	Н 841	N 151	0 178	${ m S}{ m 5}$	0	0

• Molecule 3 is a protein called Major capsid protein L1.

Mol	Chain	Residues		Α	toms			AltConf	Trace
2	Р	178	Total	С	Η	Ν	Ο	0	0
0	D	470	2869	956	958	478	477	0	0
2	Δ	178	Total	С	Η	Ν	Ο	0	0
0	A	470	2869	956	958	478	477	0	0
2	F	478	Total	С	Η	Ν	Ο	0	0
0	5 E		2869	956	958	478	477		0
2	Л	178	Total	С	Η	Ν	Ο	0	0
0	D	470	2869	956	958	478	477	0	0
2	C	178	Total	С	Η	Ν	Ο	0	0
	470	2869	956	958	478	477	0	0	
3	F	178	Total	С	Η	Ν	Ο	0	0
J	T,	410	2869	956	958	478	477		U

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	177	GLN	ASN	CONFLICT	UNP C9E771
В	181	GLN	ASN	CONFLICT	UNP C9E771
В	472	LEU	ALA	CONFLICT	UNP C9E771
А	177	GLN	ASN	CONFLICT	UNP C9E771
А	181	GLN	ASN	CONFLICT	UNP C9E771



Chain	Residue	Modelled	Actual	Comment	Reference
А	472	LEU	ALA	CONFLICT	UNP C9E771
Е	177	GLN	ASN	CONFLICT	UNP C9E771
E	181	GLN	ASN	CONFLICT	UNP C9E771
E	472	LEU	ALA	CONFLICT	UNP C9E771
D	177	GLN	ASN	CONFLICT	UNP C9E771
D	181	GLN	ASN	CONFLICT	UNP C9E771
D	472	LEU	ALA	CONFLICT	UNP C9E771
С	177	GLN	ASN	CONFLICT	UNP C9E771
С	181	GLN	ASN	CONFLICT	UNP C9E771
С	472	LEU	ALA	CONFLICT	UNP C9E771
F	177	GLN	ASN	CONFLICT	UNP C9E771
F	181	GLN	ASN	CONFLICT	UNP C9E771
F	472	LEU	ALA	CONFLICT	UNP C9E771



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: H16.U4 antibody light chain





Chain E:	91%	8%	
A9 110 111 113 113 114 114 145	P48 F83 F83 F110 0117 0197 1165 P1202 F83 F83 F84 F84 F420 F420 F420 F425 F420 F425 F420 F425 F420 F425 F420	H431 T432 P433 P434 A435	P436 K437 E438 D439
K442 461 1474 1474 8475 8475 8477 8477	K479 7480 1482 L492 K486		
• Molecule 3:	Major capsid protein L1		
Chain D:	90%	8% •	
A9 L13 P17 R41 G67	P78 P78 S89 S89 S89 S89 S89 S89 S89 P107 C115 C115 C115 C115 C115 C115 C115 C11	1422 1422 0424 0427 C428 C428	T432 P433 P434
A435 P436 R437 E438 E453 A457	F462 L474 L474 K476 K477 F480 K479 C483 K449 K485 K485 K485 K485		
• Molecule 3:	Major capsid protein L1		
Chain C:	89%	10% •	
A9 110 111 112 113 113 114 115 115 115 115 115	G38 R41 R41 R41 R43 R95 R90 C16 C16 C16 C16 C16 C16 C16 C16 C16 C16	1311 N341 T354 Y355	N359 D371
N395 G405 L406 Q407 P408 P408 P410	1416 1417 1417 1418 1418 1421 1422 1425 1425 1425 1425 1425 1425 1425 1425 1425 1425 1425 1425 1435 1435 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1436 1447 <td></td> <td></td>		
• Molecule 3:	Major capsid protein L1		
Chain F:	91%	8% •	
A9 710 711 113 113 113 113 113 819 819	R41 F51 E106 E106 V180 V180 F241 F247 F241 F241 F241 R315 R315 R315 R315 R315 R315 R315 R31	F420 A425 I426 A427 C428	0429 K430 H431
435 474 8475 8477 8477 8479 8479	R 486 R 486 R 486 R 486		



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	5806	Depositor
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	each particle	Depositor
Microscope	JEOL 2100	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{\AA}^2)$	15	Depositor
Minimum defocus (nm)	1600	Depositor
Maximum defocus (nm)	5430	Depositor
Magnification	50000	Depositor
Image detector	GATAN ULTRASCAN 4000 (4k x 4k)	Depositor
Maximum map value	6.050	Depositor
Minimum map value	-2.131	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	1.0	Depositor
Map size (Å)	894.72, 894.72, 894.72	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.33, 2.33, 2.33	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	ond lengths	Bond angles		
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	L	0.69	1/913~(0.1%)	0.68	0/1236	
2	Н	1.07	1/916~(0.1%)	1.04	5/1236~(0.4%)	
3	А	1.50	3/1910~(0.2%)	1.73	10/2386~(0.4%)	
3	В	1.52	1/1910~(0.1%)	1.76	10/2386~(0.4%)	
3	С	1.55	11/1910~(0.6%)	1.72	11/2386~(0.5%)	
3	D	1.58	7/1910~(0.4%)	1.75	13/2386~(0.5%)	
3	Ε	1.58	5/1910~(0.3%)	1.72	17/2386~(0.7%)	
3	F	1.51	4/1910~(0.2%)	1.73	17/2386~(0.7%)	
All	All	1.47	33/13289~(0.2%)	1.64	83/16788~(0.5%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	А	0	12
3	В	0	10
3	С	0	13
3	D	0	15
3	Е	0	9
3	F	0	11
All	All	0	70

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	Н	94	ARG	C-N	23.30	1.75	1.33
3	Е	206	GLY	N-CA	-6.92	1.35	1.46
3	С	290	ASN	N-CA	-6.25	1.33	1.46
3	А	164	PRO	N-CA	CA -6.25 1.36		1.47
3	Е	165	ILE	C-N	6.13	1.44	1.33
3	Е	482	LEU	C-N	6.10	1.44	1.33



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	
3	С	119	GLY	CA-C	-6.03	1.42	1.51	
3	Е	197	ASP	N-CA	-5.92	1.34	1.46	
3	А	119	GLY	N-CA	-5.85	1.37	1.46	
3	С	85	PHE	CA-C	-5.82	1.37	1.52	
3	С	38	GLY	CA-C	-5.65	1.42	1.51	
3	А	84	GLY	N-CA	-5.64	1.37	1.46	
3	С	268	GLY	N-CA	-5.61	1.37	1.46	
3	Е	83	PHE	C-N	5.58	1.43	1.33	
3	D	110	GLY	N-CA	-5.54	1.37	1.46	
3	D	67	GLY	N-CA	-5.42	1.38	1.46	
3	F	51	PRO	CA-C	-5.33	1.42	1.52	
1	L	30(B)	SER	C-N	-5.32	1.21	1.34	
3	D	159	ILE	C-N	5.29	1.42	1.33	
3	D	435	ALA	N-CA	-5.29	1.35	1.46	
3	С	116	GLY	CA-C	-5.29	1.43	1.51	
3	С	280	SER	C-N	5.25	1.42	1.33	
3	D	214	GLN	N-CA	-5.17	1.36	1.46	
3	D	408	PRO	C-N	5.14	1.44	1.34	
3	F	359	ASN	N-CA	-5.14	1.36	1.46	
3	С	12	TYR	N-CA	-5.11	1.36	1.46	
3	С	53	LYS	C-N	5.11	1.45	1.34	
3	С	149	MET	CA-C	-5.10	1.39	1.52	
3	D	182	PRO	N-CA	-5.09	1.38	1.47	
3	С	38	GLY	N-CA	-5.09	1.38	1.46	
3	В	197	ASP	C-N	5.06	1.42	1.33	
3	F	277	ILE	C-N	5.06	1.45	1.34	
3	F	13	LEU	N-CA	5.04	1.56	1.46	

All (83) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Н	94	ARG	CA-C-N	-22.36	71.48	116.20
2	Н	102	TYR	O-C-N	-8.84	108.56	122.70
3	В	420	PHE	C-N-CA	8.65	143.34	121.70
3	В	15	PRO	C-N-CA	8.15	142.07	121.70
3	F	427	ALA	O-C-N	-8.11	109.73	122.70
3	F	485	ARG	C-N-CA	8.08	141.90	121.70
2	Н	94	ARG	C-N-CA	-7.90	105.71	122.30
3	С	354	THR	O-C-N	7.33	134.43	122.70
3	А	339	SER	C-N-CA	7.11	139.48	121.70
3	F	431	HIS	C-N-CA	7.06	139.34	121.70
3	Е	479	LYS	C-N-CA	6.94	139.05	121.70



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	$Ideal(^{o})$
3	Е	482	LEU	O-C-N	-6.92	111.44	123.20
3	D	256	PHE	O-C-N	-6.76	111.89	122.70
3	В	431	HIS	C-N-CA	6.74	138.55	121.70
3	F	431	HIS	O-C-N	-6.72	111.95	122.70
3	Е	434	PRO	O-C-N	-6.43	112.42	122.70
3	А	45	VAL	N-CA-C	-6.29	94.03	111.00
3	F	426	ILE	CA-C-N	6.25	130.94	117.20
3	D	92	ASN	N-CA-C	-6.22	94.19	111.00
3	С	172	GLY	C-N-CA	6.22	137.25	121.70
3	F	483	GLY	C-N-CA	6.21	137.23	121.70
3	D	474	LEU	C-N-CA	6.21	137.22	121.70
3	D	438	GLU	N-CA-C	-6.19	94.28	111.00
3	D	428	CYS	C-N-CA	6.17	137.12	121.70
3	Е	341	ASN	N-CA-C	-6.08	94.59	111.00
3	В	415	GLU	C-N-CA	6.03	136.79	121.70
3	D	265	GLY	N-CA-C	-5.96	98.21	113.10
3	С	410	PRO	C-N-CA	5.90	134.69	122.30
3	А	406	LEU	C-N-CA	5.90	136.45	121.70
2	Н	102	TYR	CA-C-N	5.89	130.16	117.20
3	А	424	GLN	N-CA-C	-5.84	95.23	111.00
3	С	10	THR	C-N-CA	5.81	136.23	121.70
3	В	425	ALA	C-N-CA	5.70	135.94	121.70
2	Н	102	TYR	C-N-CA	5.66	135.86	121.70
3	В	395	ASN	C-N-CA	5.65	135.83	121.70
3	D	462	PHE	N-CA-C	-5.64	95.76	111.00
3	F	474	LEU	N-CA-C	-5.62	95.82	111.00
3	А	12	TYR	C-N-CA	5.62	135.74	121.70
3	F	9	ALA	CA-C-N	5.61	129.55	117.20
3	В	10	THR	C-N-CA	5.60	135.71	121.70
3	F	429	GLN	C-N-CA	5.46	135.35	121.70
3	D	422	THR	CA-C-N	5.45	129.20	117.20
3	Е	475	LYS	N-CA-C	5.43	125.67	111.00
3	D	479	LYS	O-C-N	-5.42	114.03	122.70
3	С	359	ASN	O-C-N	5.41	131.35	122.70
3	А	431	HIS	O-C-N	-5.38	114.09	122.70
3	С	341	ASN	N-CA-C	-5.36	96.53	111.00
3	Е	474	LEU	C-N-CA	5.36	135.09	121.70
3	В	432	THR	N-CA-C	-5.35	96.55	111.00
3	E	442	LYS	O-C-N	$5.3\overline{1}$	131.20	122.70
3	А	301	THR	N-CA-C	-5.30	96.68	111.00
3	D	414	LEU	C-N-CA	5.30	134.94	121.70
3	F	359	ASN	O-C-N	-5.29	114.24	122.70



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$	
3	С	422	THR	C-N-CA	5.28	134.90	121.70	
3	Е	434	PRO	C-N-CA	5.25	134.81	121.70	
3	Е	482	LEU	CA-C-N	5.24	126.69	116.20	
3	D	107	VAL	O-C-N	5.24	132.11	123.20	
3	D	139	ALA	N-CA-C	-5.23	96.89	111.00	
3	С	296	SER	C-N-CA	5.22	133.25	122.30	
3	Е	461	GLN	C-N-CA	5.21	134.72	121.70	
3	D	481	THR	C-N-CA	5.16	134.59	121.70	
3	А	363	TYR	N-CA-C	-5.15	97.09	111.00	
3	С	311	TYR	N-CA-C	-5.15	97.10	111.00	
3	F	426	ILE	O-C-N	-5.14	114.47	122.70	
3	Е	420	PHE	C-N-CA	5.14	134.54	121.70	
3	F	427	ALA	C-N-CA	5.12	134.51	121.70	
3	F	106	GLU	N-CA-C	-5.12	97.18	111.00	
3	С	371	ASP	N-CA-C	-5.09	97.26	111.00	
3	В	406	LEU	N-CA-C	-5.08	97.27	111.00	
3	F	315	ARG	N-CA-C	-5.08	97.27	111.00	
3	Е	45	VAL	N-CA-C	-5.06	97.33	111.00	
3	В	381	ILE	N-CA-C	-5.04	97.38	111.00	
3	F	342	MET	N-CA-C	-5.04	97.38	111.00	
3	Е	479	LYS	CA-C-N	5.04	128.28	117.20	
3	А	331	VAL	N-CA-C	-5.03	97.41	111.00	
3	Е	323	ILE	N-CA-C	-5.03	97.42	111.00	
3	Е	431	HIS	N-CA-C	-5.03	97.43	111.00	
3	С	10	THR	O-C-N	-5.02	114.67	122.70	
3	А	427	ALA	O-C-N	-5.02	114.67	122.70	
3	F	359	ASN	C-N-CA	5.02	134.25	121.70	
3	E	428	CYS	CA-C-N	5.01	128.23	117.20	
3	F	429	GLN	N-CA-C	-5.01	97.47	111.00	
3	Е	419	ARG	O-C-N	-5.00	114.70	122.70	

There are no chirality outliers.

All (70) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	А	105	VAL	Peptide
3	А	13	LEU	Peptide
3	А	16	VAL	Peptide
3	А	405	GLY	Peptide
3	А	407	GLN	Peptide
3	А	408	PRO	Peptide
3	А	409	PRO	Peptide



Mol	Chain	Res	Type	Group
3	А	415	GLU	Peptide
3	А	433	PRO	Peptide
3	А	436	PRO	Peptide
3	А	437	LYS	Peptide
3	А	477	LYS	Peptide
3	В	13	LEU	Peptide
3	В	14	PRO	Peptide
3	В	181	GLN	Peptide
3	В	408	PRO	Peptide
3	В	409	PRO	Peptide
3	В	414	LEU	Peptide
3	В	424	GLN	Peptide
3	В	433	PRO	Peptide
3	В	443	LYS	Peptide
3	В	481	THR	Peptide
3	С	12	TYR	Peptide
3	С	13	LEU	Peptide
3	С	14	PRO	Peptide
3	С	16	VAL	Peptide
3	С	181	GLN	Peptide
3	С	405	GLY	Peptide
3	С	407	GLN	Peptide
3	С	416	ASP	Peptide
3	С	418	TYR	Peptide
3	С	421	VAL	Peptide
3	С	435	ALA	Peptide
3	С	476	ALA	Peptide
3	С	477	LYS	Peptide
3	D	13	LEU	Peptide
3	D	17	PRO	Peptide
3	D	407	GLN	Peptide
3	D	408	PRO	Peptide
3	D	409	PRO	Peptide
3	D	422	THR	Peptide
3	D	424	GLN	Peptide
3	D	432	THR	Peptide
3	D	433	PRO	Peptide
3	D	435	ALA	Peptide
3	D	438	GLU	Peptide
3	D	473	GLY	Peptide
3	D	477	LYS	Peptide
3	D	481	THR	Peptide

Continued from previous page...



Mol	Chain	Res	Type	Group
3	D	9	ALA	Peptide
3	Е	10	THR	Peptide
3	Ε	13	LEU	Peptide
3	Е	14	PRO	Peptide
3	Е	406	LEU	Peptide
3	Ε	407	GLN	Peptide
3	Е	408	PRO	Peptide
3	Е	433	PRO	Peptide
3	Е	477	LYS	Peptide
3	Е	481	THR	Peptide
3	F	10	THR	Peptide
3	F	12	TYR	Peptide
3	F	13	LEU	Peptide
3	F	16	VAL	Peptide
3	F	19	SER	Peptide
3	F	408	PRO	Peptide
3	F	409	PRO	Peptide
3	F	414	LEU	Peptide
3	F	435	ALA	Peptide
3	F	476	ALA	Peptide
3	F	479	LYS	Peptide

Continued from previous page...

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	893	879	878	8	0
2	Н	894	841	836	20	0
3	А	1911	958	512	2	0
3	В	1911	958	512	1	0
3	С	1911	958	512	0	0
3	D	1911	958	512	0	0
3	Е	1911	958	512	0	0
3	F	1911	958	512	1	0
All	All	13253	7468	4786	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



A + am 1	A + 2	Interatomic	Clash		
Atom-1	Atom-2	distance (\AA)	overlap (Å)		
2:H:94:ARG:C	2:H:95:GLY:N	1.75	1.39		
2:H:94:ARG:CA	2:H:95:GLY:N	1.92	1.30		
2:H:94:ARG:HA	2:H:95:GLY:N	1.70	1.04		
2:H:103:TRP:O	2:H:103:TRP:CE3	2.17	0.96		
2:H:103:TRP:O	2:H:103:TRP:HE3	1.57	0.86		
2:H:103:TRP:CE3	2:H:103:TRP:N	2.47	0.83		
2:H:103:TRP:HE3	2:H:103:TRP:N	1.83	0.74		
2:H:94:ARG:C	2:H:95:GLY:CA	2.56	0.73		
2:H:94:ARG:O	2:H:94:ARG:HG3	1.90	0.69		
1:L:50:TRP:O	1:L:50:TRP:CD2	2.47	0.68		
1:L:44:PRO:O	2:H:103:TRP:NE1	2.28	0.66		
1:L:50:TRP:O	1:L:50:TRP:CE3	2.50	0.64		
2:H:94:ARG:CB	2:H:95:GLY:N	2.62	0.61		
2:H:94:ARG:O	2:H:94:ARG:CG	2.50	0.59		
1:L:50:TRP:O	1:L:50:TRP:CG	2.61	0.54		
2:H:35:SER:N	2:H:93:ALA:O	2.42	0.53		
2:H:103:TRP:CE3	2:H:103:TRP:C	2.84	0.49		
3:F:427:ALA:H	3:F:428:CYS:CA	2.27	0.48		
2:H:42:GLU:O	2:H:43:LYS:HB2	2.14	0.47		
2:H:103:TRP:CE3	2:H:103:TRP:CA	2.99	0.45		
3:A:12:TYR:H	3:A:13:LEU:CA	2.31	0.43		
1:L:19:VAL:HG22	1:L:75:ILE:HB	2.00	0.43		
1:L:33:LEU:HD22	1:L:71:PHE:CG	2.53	0.43		
2:H:109:LEU:C	2:H:109:LEU:HD23	2.40	0.42		
2:H:12:VAL:O	2:H:111:VAL:HA	2.20	0.41		
1:L:43:SER:HB3	2:H:103:TRP:NE1	2.36	0.41		
1:L:32:TYR:CD2	1:L:32:TYR:N	2.88	0.41		
3:A:431:HIS:H	3:A:432:THR:CA	2.34	0.41		
2:H:94:ARG:O	2:H:101:ASP:N	2.53	0.40		
3:B:242:TYR:O	3:B:318:GLY:HA3	2.20	0.40		

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erce	entiles
1	L	113/115~(98%)	109~(96%)	4 (4%)	0	10	00	100
2	Н	113/115~(98%)	111 (98%)	2 (2%)	0	1(00	100
3	А	476/478~(100%)	403 (85%)	55~(12%)	18 (4%)		3	24
3	В	476/478~(100%)	395~(83%)	50 (10%)	31 (6%)		1	16
3	С	476/478~(100%)	402 (84%)	52 (11%)	22~(5%)		2	21
3	D	476/478~(100%)	404 (85%)	47 (10%)	25~(5%)		2	19
3	Ε	476/478~(100%)	417 (88%)	43 (9%)	16 (3%)		3	26
3	F	476/478~(100%)	405 (85%)	52 (11%)	19 (4%)		3	23
All	All	3082/3098~(100%)	2646 (86%)	305 (10%)	131 (4%)		5	22

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (131) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
3	В	15	PRO
3	В	87	ASP
3	В	133	SER
3	В	395	ASN
3	В	408	PRO
3	В	409	PRO
3	В	414	LEU
3	В	415	GLU
3	В	439	ASP
3	А	41	ARG
3	А	86	PRO
3	А	340	THR
3	А	427	ALA
3	Ε	11	VAL
3	Ε	289	SER
3	Ε	475	LYS
3	Ε	480	PHE
3	D	152	LYS
3	D	290	ASN
3	D	476	ALA
3	С	89	SER
3	С	202	ASP
3	С	355	TYR
3	С	395	ASN



Mol	Chain	Res	Type
3	С	408	PRO
3	С	417	THR
3	С	425	ALA
3	С	478	PRO
3	F	339	SER
3	В	143	ASN
3	В	152	LYS
3	В	410	PRO
3	В	421	VAL
3	А	14	PRO
3	А	15	PRO
3	А	26	GLU
3	А	424	GLN
3	Е	425	ALA
3	D	41	ARG
3	D	408	PRO
3	D	409	PRO
3	D	457	ALA
3	D	474	LEU
3	D	482	LEU
3	С	14	PRO
3	С	26	GLU
3	С	41	ARG
3	F	315	ARG
3	F	406	LEU
3	F	415	GLU
3	В	13	LEU
3	В	14	PRO
3	В	132	ALA
3	В	202	ASP
3	В	221	PRO
3	В	326	GLY
3	В	457	ALA
3	А	351	SER
3	A	358	THR
3	Е	48	PRO
3	Е	110	GLY
3	Е	202	ASP
3	E	308	ASN
3	Е	423	SER
3	Е	479	LYS
3	D	78	PRO



Mol	Chain	Res	Type
3	D	89	SER
3	D	395	ASN
3	D	427	ALA
3	D	433	PRO
3	D	436	PRO
3	D	453	GLU
3	D	479	LYS
3	С	133	SER
3	С	182	PRO
3	С	459	LEU
3	С	479	LYS
3	F	314	GLN
3	F	395	ASN
3	F	425	ALA
3	F	477	LYS
3	В	435	ALA
3	В	475	LYS
3	В	478	PRO
3	В	479	LYS
3	В	482	LEU
3	А	13	LEU
3	А	185	CYS
3	А	247	PHE
3	А	308	ASN
3	А	437	LYS
3	А	440	PRO
3	А	462	PHE
3	Е	314	GLN
3	Е	439	ASP
3	D	90	PHE
3	D	139	ALA
3	D	259	HIS
3	D	410	PRO
3	D	434	PRO
3	C	16	VAL
3	С	161	CYS
3	C	173	SER
3	С	436	PRO
3	F	11	VAL
3	F	241	PRO
3	F	247	PHE
3	F	259	HIS



Mal	Chain	Doc	Typo
IVIOI	Unain	nes	Type
3	F	479	LYS
3	В	20	LYS
3	В	173	SER
3	В	403	ASN
3	В	416	ASP
3	В	445	THR
3	Е	409	PRO
3	D	484	LYS
3	D	485	ARG
3	С	90	PHE
3	F	41	ARG
3	F	218	SER
3	F	403	ASN
3	F	420	PHE
3	А	137	ALA
3	Е	41	ARG
3	Е	259	HIS
3	D	403	ASN
3	F	417	THR
3	В	434	PRO
3	F	180	VAL
3	С	407	GLN
3	С	164	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	L	100/100~(100%)	99~(99%)	1 (1%)	76	86
2	Н	95/95~(100%)	94~(99%)	1 (1%)	73	84
All	All	195/195~(100%)	193~(99%)	2(1%)	77	86

All (2) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	L	30(F)	LYS
2	Н	103	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	Н	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	Н	94:ARG	С	95:GLY	N	1.75



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-6424. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 192



Y Index: 192



Z Index: 192



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 217

Y Index: 167

Z Index: 217

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 64244 nm^3 ; this corresponds to an approximate mass of 58034 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.083 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-6424 and PDB model 3JBA. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlays

9.1.1 Map-model overlay (i)



9.1.2 Map-model assembly overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 1.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.0).



9.4 Atom inclusion (i)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (1.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score	
All	0.9825	0.0760	
А	1.0000	0.0880	
В	0.9995	0.0830	
С	0.9990	0.0880	
D	0.9922	0.0830	
Е	0.9885	0.0860	
F	1.0000	0.0810	
Н	0.9217	0.0260	
L	0.8219	0.0090	

