

Full wwPDB X-ray Structure Validation Report (i)

Jan 11, 2024 - 12:09 PM JST

PDB ID	:	8JBC
Title	:	Crystal Structure of the Csm6 K137A mutant from Thermus thermophilus
		HB8 in its apo form
Authors	:	Lin, Z.; Du, L.
Deposited on	:	2023-05-08
Resolution	:	2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$	
R_{free}	130704	4647 (2.44-2.40)	
Clashscore	141614	5161(2.44-2.40)	
Ramachandran outliers	138981	5073(2.44-2.40)	
Sidechain outliers	138945	5074(2.44-2.40)	
RSRZ outliers	127900	4543 (2.44-2.40)	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	468	78%	22%	•
1	В	468	8%	22%	••



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	168	Total	С	Ν	Ο	S	0	0	0
	400	3576	2302	629	640	5	0	0	0	
1	р	456	Total	С	Ν	0	S	0	0	0
I D	400	3536	2276	627	628	5	0	0		

• Molecule 1 is a protein called CRISPR system endoribonuclease Csm6.

Chain	Residue	Modelled	Actual	Comment	Reference
А	-3	SER	-	expression tag	UNP Q53W17
А	-2	GLY	-	expression tag	UNP Q53W17
А	-1	ARG	-	expression tag	UNP Q53W17
А	0	PRO	-	expression tag	UNP Q53W17
А	137	ALA	LYS	engineered mutation	UNP Q53W17
В	-3	SER	-	expression tag	UNP Q53W17
В	-2	GLY	-	expression tag	UNP Q53W17
В	-1	ARG	-	expression tag	UNP Q53W17
В	0	PRO	-	expression tag	UNP Q53W17
В	137	ALA	LYS	engineered mutation	UNP Q53W17

There are 10 discrepancies between the modelled and reference sequences:

• Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Ni 1 1	0	0
2	В	1	Total Ni 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	23	TotalO2323	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	12	Total O 12 12	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: CRISPR system endoribonuclease Csm6







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	94.89Å 205.57Å 59.10Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	29.64 - 2.41	Depositor
Resolution (A)	59.10 - 2.41	EDS
% Data completeness	99.9 (29.64-2.41)	Depositor
(in resolution range)	100.0 (59.10-2.41)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.85 (at 2.42 \text{\AA})$	Xtriage
Refinement program	REFMAC 1.20.1_4487	Depositor
R R.	0.222 , 0.269	Depositor
Π, Π_{free}	0.239 , 0.250	DCC
R_{free} test set	2292 reflections (5.03%)	wwPDB-VP
Wilson B-factor $(Å^2)$	62.3	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 55.1	EDS
L-test for $twinning^2$	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7149	wwPDB-VP
Average B, all atoms $(Å^2)$	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles		
1VIOI	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.46	0/3653	0.67	1/4963~(0.0%)	
1	В	0.48	1/3608~(0.0%)	0.70	2/4887~(0.0%)	
All	All	0.47	1/7261~(0.0%)	0.68	3/9850~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	31	PRO	CG-CD	-7.30	1.26	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	88	LEU	CA-CB-CG	5.69	128.39	115.30
1	А	282	LEU	CA-CB-CG	5.44	127.82	115.30
1	В	7	LEU	CB-CG-CD2	-5.27	102.04	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3576	0	3616	64	1
1	В	3536	0	3620	75	1



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	А	1	0	0	0	0
2	В	1	0	0	0	0
3	А	23	0	0	1	0
3	В	12	0	0	2	0
All	All	7149	0	7236	132	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (132) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:273:ARG:NH1	1:A:277:GLU:OE2	2.05	0.90	
1:B:136:THR:H	1:B:139:MET:HE2	1.38	0.88	
1:A:207:GLU:OE1	1:A:210:GLN:NE2	2.09	0.85	
1:B:39:GLU:HB2	1:B:40:LYS:HD2	1.60	0.83	
1:A:115:GLU:OE1	1:A:118:ARG:NH1	2.14	0.78	
1:A:-2:GLY:N	1:A:0:PRO:HD3	2.00	0.77	
1:B:106:LYS:HG3	1:B:136:THR:HG21	1.68	0.76	
1:A:74:GLU:OE1	1:A:74:GLU:N	2.19	0.76	
1:A:83:GLU:N	1:A:83:GLU:OE2	2.19	0.73	
1:A:0:PRO:HD2	1:A:3:ASP:HB2	1.72	0.71	
1:A:278:ALA:HB2	1:A:454:PRO:HD2	1.73	0.70	
1:A:428:SER:O	1:A:431:ALA:N	2.26	0.69	
1:A:-2:GLY:H	1:A:0:PRO:HD3	1.58	0.67	
1:A:315:ALA:HB1	1:A:319:ALA:HB3	1.76	0.67	
1:B:34:LEU:HD22	1:B:38:ARG:HH11	1.60	0.67	
1:A:218:MET:O	1:A:222:THR:HG22	1.94	0.67	
1:B:207:GLU:OE1	1:B:210:GLN:NE2	2.28	0.65	
1:A:109:VAL:HG21	1:B:184:LEU:HD11	1.78	0.65	
1:A:306:ARG:HH21	1:A:457:PRO:HA	1.61	0.64	
1:A:222:THR:HG23	1:A:224:ASN:H	1.61	0.64	
1:A:168:ASP:HB3	1:A:171:LEU:H	1.62	0.63	
1:B:42:ARG:HG2	1:B:44:TYR:CE1	2.34	0.63	
1:B:30:TRP:HB3	1:B:31:PRO:HD2	1.80	0.62	
1:B:41:PRO:HB2	1:B:70:GLY:HA2	1.81	0.62	
1:B:12:ARG:O	1:B:16:ARG:HG3	2.00	0.62	
1:B:30:TRP:NE1	1:B:94:ASP:OD2	2.32	0.61	
1:A:191:LEU:O	1:A:194:VAL:HG22	2.00	0.61	
1:B:193:GLU:OE1	1:B:227:TYR:OH	2.18	0.61	
1:A:12:ARG:NH1	1:A:168:ASP:OD2	2.33	0.61	



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:115:GLU:O	1:B:119:LEU:HG	2.00	0.61
1:B:24:LEU:O	1:B:28:MET:HG2	2.01	0.60
1:B:85:ALA:O	1:B:88:LEU:HD12	2.02	0.59
1:A:417:SER:HB2	1:A:425:ASP:OD2	2.02	0.59
1:A:452:LEU:HD12	1:A:453:GLY:O	2.04	0.58
1:B:3:ASP:N	1:B:3:ASP:OD1	2.38	0.57
1:A:0:PRO:HG2	1:A:3:ASP:H	1.69	0.57
1:A:117:LYS:NZ	1:A:121:GLU:OE2	2.38	0.56
1:B:20:ASN:CG	1:B:23:ALA:HB3	2.26	0.55
1:A:357:GLU:O	1:A:361:LYS:HG3	2.07	0.55
1:B:34:LEU:HD21	1:B:94:ASP:O	2.06	0.55
1:A:322:LEU:HD21	1:B:322:LEU:HD11	1.88	0.55
1:B:396:ALA:HA	1:B:399:ARG:HH21	1.72	0.55
1:A:306:ARG:NH2	1:A:457:PRO:HA	2.22	0.54
1:A:352:SER:OG	1:A:355:GLU:HG3	2.08	0.54
1:B:195:ASP:OD2	1:B:221:ARG:NH2	2.35	0.54
1:A:40:LYS:O	1:A:40:LYS:HG2	2.09	0.53
1:A:325:LEU:HD21	1:B:420:LEU:HB3	1.91	0.53
1:A:446:LEU:O	1:A:450:THR:HG23	2.09	0.53
1:A:109:VAL:HG21	1:B:184:LEU:CD1	2.38	0.52
1:A:56:THR:HG23	1:A:143:LEU:HD22	1.91	0.52
1:A:103:GLU:O	1:A:104:ILE:HD12	2.07	0.52
1:B:104:ILE:HD12	1:B:104:ILE:O	2.10	0.52
1:B:302:TYR:HB2	1:B:446:LEU:HD22	1.92	0.52
1:A:366:LEU:HD21	1:A:405:LEU:HD22	1.91	0.52
1:A:400:LEU:HD12	1:A:405:LEU:HB2	1.93	0.51
1:B:446:LEU:O	1:B:450:THR:HG22	2.11	0.51
1:B:202:LEU:HD13	1:B:210:GLN:HB3	1.92	0.51
1:A:338:ARG:NH1	1:A:445:ASP:OD2	2.44	0.50
1:B:42:ARG:O	1:B:42:ARG:HD2	2.11	0.50
1:B:264:TRP:O	1:B:270:ASN:ND2	2.44	0.50
1:B:371:PRO:O	1:B:374:VAL:HG22	2.12	0.50
1:A:8:TRP:CE3	1:A:176:ALA:HB2	2.47	0.50
1:A:331:LEU:HD21	1:A:438:LEU:HD23	1.93	0.49
1:A:363:LEU:O	1:A:367:LEU:HG	2.12	0.49
1:A:394:ASP:HB3	1:A:397:LEU:HB2	1.93	0.49
1:A:85:ALA:HA	1:A:88:LEU:HD23	1.95	0.49
1:B:297:LEU:O	1:B:301:VAL:HG23	2.14	0.48
1:B:299:GLU:H	1:B:299:GLU:CD	2.17	0.48
1:B:173:ARG:HG2	1:B:173:ARG:HH11	1.79	0.47
1:B:44:TYR:CZ	1:B:180:LYS:HE3	2.50	0.47



	lo ao pagom	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:244:ASP:OD2	3:B:1101:HOH:O	2.20	0.47
1:B:55:HIS:ND1	1:B:131:ASP:OD2	2.40	0.46
1:B:253:ARG:NH2	1:B:284:GLU:OE1	2.48	0.46
1:B:3:ASP:O	1:B:7:LEU:HB2	2.15	0.46
1:B:23:ALA:HA	1:B:26:GLN:HE21	1.81	0.46
1:B:133:THR:O	3:B:1102:HOH:O	2.21	0.46
1:B:306:ARG:HG3	1:B:443:LEU:HD21	1.97	0.46
1:B:401:SER:OG	1:B:404:GLU:HG2	2.16	0.46
1:B:58:GLY:HA3	1:B:135:GLY:HA2	1.96	0.46
1:B:245:PHE:CE1	1:B:286:VAL:HG11	2.50	0.46
1:B:427:PRO:O	1:B:429:PRO:HD3	2.14	0.46
1:A:10:ARG:HD2	1:A:28:MET:HE3	1.98	0.46
1:B:173:ARG:HG3	1:B:174:PRO:O	2.15	0.46
1:B:380:LEU:HD22	1:B:384:ASP:HB3	1.97	0.46
1:A:366:LEU:HD21	1:A:405:LEU:CD2	2.45	0.46
1:A:184:LEU:HD11	1:B:109:VAL:HG21	1.98	0.45
1:A:254:LYS:O	1:A:258:GLN:HG2	2.17	0.45
1:A:149:PHE:CZ	1:B:194:VAL:HG21	2.51	0.45
1:B:23:ALA:HA	1:B:26:GLN:CG	2.47	0.45
1:A:219:VAL:HG21	1:A:228:ALA:HB2	1.99	0.45
1:B:267:HIS:CE1	1:B:269:LEU:HD12	2.52	0.45
1:B:344:ARG:NH2	1:B:350:GLY:O	2.49	0.45
1:B:7:LEU:HA	1:B:10:ARG:HD3	1.99	0.45
1:B:85:ALA:HA	1:B:88:LEU:HD12	1.99	0.44
1:A:90:ARG:HD2	3:A:1111:HOH:O	2.17	0.44
1:A:408:LEU:HB3	1:A:412:LEU:HD23	2.00	0.44
1:B:392:LYS:HA	1:B:392:LYS:HD3	1.64	0.44
1:B:33:LEU:HD13	1:B:176:ALA:O	2.17	0.44
1:A:81:THR:HA	1:A:104:ILE:O	2.18	0.43
1:B:57:LEU:N	1:B:79:LEU:O	2.50	0.43
1:A:50:PHE:HA	1:A:127:PRO:HB2	2.00	0.43
1:B:411:ALA:HB2	1:B:431:ALA:HB1	2.01	0.43
1:A:82:PRO:HG3	1:A:103:GLU:OE1	2.18	0.43
1:A:104:ILE:HG22	1:A:105:GLY:O	2.19	0.43
1:A:184:LEU:HD11	1:B:109:VAL:HG11	2.01	0.43
1:A:12:ARG:O	1:A:16:ARG:HD3	2.18	0.43
1:B:428:SER:O	1:B:431:ALA:N	2.52	0.43
1:A:366:LEU:HD12	1:A:367:LEU:HD23	2.01	0.42
1:A:110:GLU:O	1:A:114:ARG:HG3	2.18	0.42
1:A:200:LYS:HG3	$1:\overline{A:462:LEU:HD12}$	2.00	0.42
1:B:338:ARG:NH1	1:B:445:ASP:OD1	2.52	0.42



	A 4 am 2	Interatomic	Clash	
Atom-1	Atom-2	distance (\AA)	overlap (Å)	
1:B:342:LEU:O	1:B:344:ARG:HG3	2.19	0.42	
1:A:33:LEU:HD13	1:A:176:ALA:O	2.19	0.42	
1:B:331:LEU:HD22	1:B:435:ILE:HG23	2.00	0.42	
1:A:168:ASP:HB3	1:A:171:LEU:N	2.32	0.42	
1:B:215:PHE:O	1:B:218:MET:HB2	2.20	0.42	
1:A:189:GLU:O	1:A:221:ARG:NH2	2.53	0.42	
1:A:321:VAL:HA	1:A:427:PRO:HG3	2.00	0.41	
1:B:341:LEU:HD23	1:B:341:LEU:HA	1.88	0.41	
1:A:46:PHE:HA	1:A:47:PRO:HD3	1.91	0.41	
1:B:136:THR:N	1:B:139:MET:HE2	2.19	0.41	
1:B:179:GLU:N	1:B:179:GLU:CD	2.74	0.41	
1:B:338:ARG:HA	1:B:338:ARG:HD2	1.90	0.41	
1:A:364:ALA:HB1	1:A:371:PRO:HB3	2.03	0.41	
1:B:35:ALA:HA	1:B:38:ARG:NH1	2.36	0.41	
1:A:25:TYR:CD2	1:A:173:ARG:HD3	2.55	0.41	
1:B:39:GLU:CB	1:B:40:LYS:HD2	2.43	0.41	
1:B:229:LEU:HD12	1:B:267:HIS:CD2	2.56	0.41	
1:B:186:ASN:OD1	1:B:189:GLU:HG2	2.21	0.40	
1:B:20:ASN:O	1:B:24:LEU:HD12	2.21	0.40	
1:B:20:ASN:O	1:B:23:ALA:N	2.55	0.40	
1:B:56:THR:HG23	1:B:143:LEU:HD22	2.02	0.40	

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-3:SER:OG	1:B:188:HIS:NE2[1_556]	1.97	0.23

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	466/468~(100%)	448 (96%)	18 (4%)	0	100	100
1	В	448/468~(96%)	430 (96%)	18 (4%)	0	100	100
All	All	914/936~(98%)	878~(96%)	36~(4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	345/358~(96%)	332~(96%)	13~(4%)	33 50
1	В	347/358~(97%)	331~(95%)	16 (5%)	27 42
All	All	692/716~(97%)	663 (96%)	29 (4%)	30 46

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	12	ARG
1	А	16	ARG
1	А	59	THR
1	А	165	GLU
1	А	168	ASP
1	А	188	HIS
1	А	365	GLU
1	А	376	LEU
1	А	384	ASP
1	А	390	ARG
1	А	415	ARG
1	А	417	SER
1	А	460	VAL
1	В	3	ASP
1	В	5	ASP
1	В	26	GLN
1	В	40	LYS
1	В	42	ARG



Mol	Chain	Res	Type
1	В	84	SER
1	В	107	SER
1	В	134	SER
1	В	258	GLN
1	В	273	ARG
1	В	345	ARG
1	В	352	SER
1	В	401	SER
1	В	415	ARG
1	В	450	THR
1	В	452	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	210	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	468/468~(100%)	0.32	19 (4%) 37	35	41, 68, 116, 137	0
1	В	456/468~(97%)	0.48	38 (8%) 11	10	46, 72, 113, 133	0
All	All	924/936~(98%)	0.40	57 (6%) 20	19	41, 70, 114, 137	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	А	167	TYR	10.4	
1	В	359	LEU	5.0	
1	В	29	VAL	4.5	
1	А	454	PRO	4.4	
1	В	376	LEU	4.4	
1	А	367	LEU	4.3	
1	В	33	LEU	4.3	
1	В	262	ASN	4.3	
1	А	391	LEU	4.3	
1	В	351	LEU	4.2	
1	В	46	PHE	4.0	
1	В	28	MET	3.9	
1	В	4	LEU	3.9	
1	А	372	GLU	3.8	
1	В	15	VAL	3.6	
1	В	66	LEU	3.6	
1	А	17	ALA	3.5	
1	В	388	PHE	3.5	
1	В	107	SER	3.4	
1	В	355	GLU	3.3	
1	В	354	GLU	3.2	
1	А	353	PRO	3.2	
1	A	-3	SER	3.2	
1	В	373	GLU	3.2	



Mol	Chain	Res	Type	RSRZ
1	В	1	MET	3.1
1	А	455	LEU	3.0
1	А	388	PHE	3.0
1	В	176	ALA	3.0
1	А	44	TYR	2.9
1	В	17	ALA	2.8
1	В	31	PRO	2.8
1	А	378	ALA	2.8
1	А	457	PRO	2.8
1	А	389	LEU	2.7
1	В	370	LEU	2.7
1	А	366	LEU	2.6
1	В	177	GLY	2.6
1	В	12	ARG	2.6
1	В	358	ALA	2.5
1	А	15	VAL	2.5
1	В	170	GLU	2.5
1	В	378	ALA	2.5
1	В	391	LEU	2.4
1	В	69	LEU	2.4
1	А	359	LEU	2.3
1	А	400	LEU	2.3
1	В	385	LEU	2.3
1	В	32	ALA	2.2
1	В	339	LEU	2.2
1	В	379	LYS	2.1
1	В	361	LYS	2.1
1	В	36	LEU	2.1
1	A	361	LYS	2.0
1	В	30	TRP	2.0
1	В	41	PRO	2.0
1	В	63	ALA	2.0
1	В	11	TYR	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	NI	А	1001	1/1	0.76	0.06	122,122,122,122	0
2	NI	В	1001	1/1	0.83	0.06	110,110,110,110	0

6.5 Other polymers (i)

There are no such residues in this entry.

