

Full wwPDB X-ray Structure Validation Report (i)

Jan 30, 2024 - 10:13 PM EST

PDB ID	:	1JFD
Title	:	STRUCTURE OF INORGANIC PYROPHOSPHATASE
Authors	:	Oganesyan, V.; Avaeva, S.M.; Huber, R.; Harutyunyan, E.H.
Deposited on		
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$		
Clashscore	141614	5594 (2.20-2.20)		
Ramachandran outliers	138981	5503 (2.20-2.20)		
Sidechain outliers	138945	5504 (2.20-2.20)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	175	81%	15%	•
1	В	175	85%	14%	•




2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- ZeroOcc AltConf Mol Chain Residues Trace Atoms Total С Ν 0 S 0 0 0 1 А 1751381889 2242635С \mathbf{S} Total Ν Ο 1 В 0 0 0 1751381889 2242635
- Molecule 1 is a protein called INORGANIC PYROPHOSPHATASE.

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	103	Total O 103 103	0	0
3	В	104	Total O 104 104	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: INORGANIC PYROPHOSPHATASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	110.40Å 110.40Å 154.80Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 - 2.20	Depositor
% Data completeness	(Not available) (15.00-2.20)	Depositor
(in resolution range)	(100 available) (15.00 2.20)	Depositor
R_{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.192 , 0.266	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2979	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Chain Bond lengths			Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.47	1/1416~(0.1%)	0.68	0/1923	
1	В	0.43	0/1416	0.69	0/1923	
All	All	0.45	1/2832~(0.0%)	0.68	0/3846	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	87	CYS	CB-SG	-5.38	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1381	0	1369	26	0
1	В	1381	0	1369	14	0
2	А	5	0	0	1	0
2	В	5	0	0	0	0
3	А	103	0	0	2	0
3	В	104	0	0	2	0
All	All	2979	0	2738	40	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:ASN:HB2	3:B:278:HOH:O	1.69	0.91
1:B:92:VAL:HG13	1:B:156:GLU:HB2	1.65	0.77
1:A:126:LEU:HB2	1:A:131:LYS:HD2	1.70	0.74
1:B:11:LEU:HD23	1:B:15:ILE:HG22	1.68	0.72
1:A:146:LYS:HD2	1:A:146:LYS:H	1.56	0.71
1:A:91:GLY:HA3	1:A:123:VAL:HG13	1.77	0.67
1:A:151:LYS:HA	1:A:151:LYS:HE2	1.76	0.67
1:A:112:LYS:HD3	1:A:113:LEU:HD12	1.77	0.66
1:B:105:LEU:HD13	1:B:161:ALA:HB1	1.80	0.63
1:B:63:SER:HB2	1:B:65:ASP:OD1	2.00	0.62
1:A:115:LYS:HZ2	1:A:115:LYS:HA	1.66	0.61
1:A:92:VAL:HG13	1:A:156:GLU:HB2	1.84	0.59
1:A:112:LYS:HD2	3:A:237:HOH:O	2.06	0.56
1:A:146:LYS:H	1:A:146:LYS:CD	2.17	0.54
1:A:115:LYS:HA	1:A:115:LYS:NZ	2.22	0.53
1:A:145:GLU:HB3	1:A:148:LYS:HG3	1.91	0.53
1:A:98:GLU:HA	1:A:149:TRP:CZ2	2.46	0.51
1:B:105:LEU:CD1	1:B:161:ALA:HB1	2.39	0.50
1:A:111:SER:HB2	1:A:115:LYS:HZ1	1.75	0.50
1:A:8:GLY:HA3	1:A:15:ILE:HA	1.94	0.49
1:A:27:PRO:HB2	1:A:45:MET:HB2	1.95	0.49
1:A:88:ARG:HG3	3:A:212:HOH:O	2.11	0.49
1:A:120:ILE:HG23	1:A:125:ASP:HB2	1.95	0.48
1:A:93:LEU:HD11	1:A:152:VAL:HG13	1.96	0.48
1:A:151:LYS:HA	1:A:151:LYS:CE	2.43	0.47
1:B:111:SER:HB3	1:B:115:LYS:NZ	2.30	0.47
1:B:115:LYS:NZ	1:B:118:ASP:OD2	2.48	0.47
1:B:68:PRO:HG2	3:B:213:HOH:O	2.16	0.46
1:A:111:SER:CB	1:A:115:LYS:HZ1	2.30	0.44
1:B:123:VAL:HG21	1:B:155:TRP:HB3	1.99	0.44
1:A:123:VAL:HG22	1:A:157:ASN:HA	2.00	0.44
1:A:115:LYS:HG3	1:A:118:ASP:OD2	2.18	0.43
1:A:87:CYS:HB3	1:A:107:ALA:HB1	2.02	0.42
1:A:175:LYS:NZ	1:A:175:LYS:HB2	2.33	0.42
1:B:18:VAL:O	1:B:56:GLY:HA3	2.20	0.42
1:B:11:LEU:CD2	1:B:15:ILE:HG22	2.43	0.42
1:B:5:VAL:HA	1:B:6:PRO:HD3	1.97	0.41

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:GLU:HG3	1:B:152:VAL:HG23	2.02	0.41
1:A:142:LYS:NZ	2:A:176:SO4:O3	2.52	0.41
1:B:2:LEU:HD23	1:B:32:ILE:HD12	2.03	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
1	А	173/175~(99%)	168 (97%)	5(3%)	0	100	100
1	В	173/175~(99%)	168 (97%)	5(3%)	0	100	100
All	All	346/350~(99%)	336~(97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	150/150~(100%)	142~(95%)	8 (5%)	22 27		
1	В	150/150~(100%)	142 (95%)	8 (5%)	22 27		
All	All	300/300~(100%)	284~(95%)	16 (5%)	22 27		

All (16) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	115	LYS
1	А	123	VAL
1	А	141	TYR
1	А	146	LYS
1	А	148	LYS
1	А	151	LYS
1	А	174	ASN
1	А	175	LYS
1	В	2	LEU
1	В	9	LYS
1	В	29	LYS
1	В	105	LEU
1	В	115	LYS
1	В	128	GLU
1	В	141	TYR
1	В	153	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	174	ASN
1	В	24	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Dag	Link	Bond lengths			Bond angles		
MOI	туре	Chain	Res	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	SO4	В	177	-	4,4,4	0.70	0	$6,\!6,\!6$	0.43	0
2	SO4	А	176	-	4,4,4	1.03	0	$6,\!6,\!6$	0.54	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	176	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

