Full wwPDB X-ray Structure Validation Report (i)

Sep 19, 2023 – 09:13 PM EDT

PDB ID	:	5JHW
Title	:	Crystal Structure of the GDF11:Follistatin 288 complex
Authors	:	Walker, R.G.; Thompson, T.B.
Deposited on	:	2016-04-21
Resolution	:	2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	109	2% 96% •
1	B	109	3% 94% 6%
2	C	288	6% 89% 9% •
2	D	288	5% 88% 10% •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6323 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Growth/differentiation factor 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	109	885	557	153	161	14	0	2	0
1	B	109	879	554	150	161	14	0	1	0

- Molecule 2 is a protein called Follistatin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	282	2135	1298	383	415	39	0	1	0
2	D	284	2148	1304	385	420	39	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
3	A	1	5	4	1	0	0
			Total	O	P		
3	C	1	5	4	1	0	0
			Total	O	P		
3	D	1	5	4	1	0	0

- Molecule 4 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	13	6	7	0	0
			Total	C	O		
4	B	1	13	6	7	0	0
			Total	C	O		
4	B	1	13	6	7	0	0
			Total	C	O		
4	C	1	13	6	7	0	0
			Total	C	O		
4	C	1	13	6	7	0	0
			Total	C	O		
4	D	1	13	6	7	0	0
			Total	C	O		
4	D	1	13	6	7	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	29	29	29	0	0
			Total	O		
5	B	41	41	41	0	0
			Total	O		
5	C	53	53	53	0	0
			Total	O		
5	D	47	47	47	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Growth/differentiation factor 11

Chain A:





• Molecule 1: Growth/differentiation factor 11

Chain B:





• Molecule 2: Follistatin

Chain C:





• Molecule 2: Follistatin

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	55.96Å 59.08Å 288.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.28 – 2.35 41.28 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.28-2.35) 99.9 (41.28-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.71 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.202 , 0.247 0.204 , 0.247	Depositor DCC
R_{free} test set	2571 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å^2)	45.2	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 37.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6323	wwPDB-VP
Average B, all atoms (Å^2)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.31	0/910	0.49	0/1227
1	B	0.30	0/905	0.44	0/1223
2	C	0.30	0/2171	0.51	0/2920
2	D	0.29	0/2181	0.51	0/2933
All	All	0.30	0/6167	0.50	0/8303

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	885	0	841	4	0
1	B	879	0	825	5	0
2	C	2135	0	2050	18	0
2	D	2148	0	2056	20	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	13	0	5	0	0
4	B	26	0	10	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	26	0	10	2	0
4	D	26	0	10	2	0
5	A	29	0	0	0	0
5	B	41	0	0	1	0
5	C	53	0	0	0	0
5	D	47	0	0	5	0
All	All	6323	0	5807	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:162:THR:HG23	2:C:165:ARG:HH21	1.51	0.76
2:D:193:LYS:O	2:D:197:LEU:HG	1.98	0.63
2:C:207:GLU:OE1	2:C:222:THR:OG1	2.10	0.62
2:C:165:ARG:N	2:C:165:ARG:HD3	2.15	0.62
2:C:163:CYS:O	2:C:165:ARG:NE	2.33	0.61
4:C:303:FLC:OA2	4:C:303:FLC:OHB	2.16	0.59
1:A:8:GLU:H	1:A:8:GLU:CD	2.06	0.59
2:D:108:LYS:HD2	2:D:109:THR:N	2.17	0.59
2:C:115:ALA:HA	2:C:118:LYS:HE2	1.83	0.59
2:D:92:ASP:OD1	2:D:94:SER:OG	2.20	0.57
2:D:51:ILE:O	5:D:401:HOH:O	2.18	0.56
2:D:267:ALA:O	5:D:402:HOH:O	2.18	0.55
2:C:264:SER:O	2:C:268:MET:N	2.39	0.54
2:D:111:ARG:NH1	2:D:112:ASN:HD21	2.06	0.54
1:A:14:ARG:HH21	2:D:18:LYS:HG2	1.73	0.53
2:D:251:ASP:HB3	2:D:264:SER:HB3	1.90	0.53
2:C:264:SER:H	2:C:267:ALA:HB3	1.73	0.52
1:B:3:GLY:HA3	1:B:105:ARG:HG3	1.91	0.51
1:B:90:LYS:NZ	4:C:302:FLC:HG2	2.26	0.49
2:C:82:LYS:O	2:C:82:LYS:HG3	2.12	0.49
1:B:89:ASP:OD2	5:B:301:HOH:O	2.19	0.48
2:D:136:LYS:HD2	2:D:142:VAL:HG22	1.94	0.48
2:D:219:ILE:O	5:D:403:HOH:O	2.20	0.48
1:A:14:ARG:NH2	4:D:301:FLC:OB1	2.46	0.48
2:D:136:LYS:NZ	2:D:141:ASP:O	2.46	0.48
2:C:164:ASN:C	2:C:165:ARG:HD3	2.35	0.48
2:C:81:LYS:N	2:C:81:LYS:HD3	2.30	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:126:GLU:OE1	4:D:302:FLC:HA2	2.16	0.46
1:B:14:ARG:HH21	2:C:18:LYS:HG2	1.81	0.46
1:B:97:LYS:HE2	1:B:97:LYS:HB3	1.70	0.45
2:D:4:TRP:CG	2:D:13:CYS:HB3	2.52	0.45
2:C:207:GLU:H	2:C:207:GLU:HG3	1.50	0.44
2:D:14:GLN:HG2	5:D:431:HOH:O	2.17	0.44
2:C:113:GLU:OE1	2:C:129:VAL:HG21	2.18	0.44
4:B:202:FLC:OA2	4:B:202:FLC:OHB	2.34	0.44
2:D:113:GLU:OE1	2:D:129:VAL:HG21	2.18	0.44
2:C:78:ARG:HD2	2:C:88:VAL:HG21	2.00	0.43
2:C:265:GLU:O	2:C:269:LYS:HG3	2.19	0.43
2:C:96:ILE:HD13	2:C:113:GLU:HG2	2.01	0.43
2:C:4:TRP:CG	2:C:13:CYS:HB3	2.54	0.42
2:D:179:GLY:O	5:D:404:HOH:O	2.22	0.42
2:D:93:CYS:HA	2:D:96:ILE:HD12	2.02	0.42
2:D:76:LYS:HA	2:D:76:LYS:HD2	1.85	0.42
2:C:193:LYS:HB3	2:C:193:LYS:HE2	1.82	0.41
1:A:81:PRO:HB3	1:A:99:PRO:HA	2.03	0.41
2:D:109:THR:HG23	2:D:134:ARG:HA	2.02	0.41
2:D:122:LYS:HB3	2:D:122:LYS:HE2	1.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/109 (100%)	106 (97%)	3 (3%)	0	100	100
1	B	108/109 (99%)	106 (98%)	2 (2%)	0	100	100
2	C	277/288 (96%)	270 (98%)	6 (2%)	1 (0%)	34	38
2	D	278/288 (96%)	269 (97%)	8 (3%)	1 (0%)	34	38
All	All	772/794 (97%)	751 (97%)	19 (2%)	2 (0%)	41	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	15	VAL
2	D	15	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/96 (102%)	98 (100%)	0	100	100
1	B	97/96 (101%)	97 (100%)	0	100	100
2	C	242/248 (98%)	242 (100%)	0	100	100
2	D	244/248 (98%)	244 (100%)	0	100	100
All	All	681/688 (99%)	681 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	PO4	C	301	-	4,4,4	0.90	0	6,6,6	0.38	0
4	FLC	B	202	-	12,12,12	1.52	1 (8%)	17,17,17	1.08	2 (11%)
4	FLC	D	301	-	12,12,12	1.58	2 (16%)	17,17,17	2.30	2 (11%)
4	FLC	D	302	-	12,12,12	1.53	1 (8%)	17,17,17	1.08	2 (11%)
3	PO4	A	201	-	4,4,4	1.02	0	6,6,6	0.32	0
4	FLC	B	201	-	12,12,12	1.44	1 (8%)	17,17,17	0.91	2 (11%)
4	FLC	A	202	-	12,12,12	1.35	1 (8%)	17,17,17	0.54	0
4	FLC	C	303	-	12,12,12	1.69	2 (16%)	17,17,17	2.45	4 (23%)
3	PO4	D	303	-	4,4,4	0.92	0	6,6,6	0.45	0
4	FLC	C	302	-	12,12,12	1.62	1 (8%)	17,17,17	1.26	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FLC	B	202	-	-	14/16/16/16	-
4	FLC	D	301	-	-	4/16/16/16	-
4	FLC	D	302	-	-	0/16/16/16	-
4	FLC	B	201	-	-	5/16/16/16	-
4	FLC	A	202	-	-	5/16/16/16	-
4	FLC	C	303	-	-	9/16/16/16	-
4	FLC	C	302	-	-	6/16/16/16	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	303	FLC	CB-CBC	3.95	1.57	1.53
4	D	302	FLC	CB-CBC	3.91	1.57	1.53
4	C	302	FLC	CB-CBC	3.88	1.57	1.53
4	A	202	FLC	CB-CBC	3.85	1.57	1.53
4	B	202	FLC	CB-CBC	3.84	1.57	1.53
4	D	301	FLC	CB-CBC	3.83	1.57	1.53
4	B	201	FLC	CB-CBC	3.75	1.57	1.53
4	C	303	FLC	OB1-CBC	2.25	1.29	1.22
4	D	301	FLC	OB1-CBC	2.19	1.29	1.22

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	C	303	FLC	OB1-CBC-CB	-7.04	112.28	122.25
4	D	301	FLC	OB1-CBC-CB	-7.03	112.30	122.25
4	C	303	FLC	OB2-CBC-CB	6.19	123.80	113.05
4	D	301	FLC	OB2-CBC-CB	6.11	123.67	113.05
4	D	302	FLC	OA2-CAC-CA	2.74	123.16	114.35
4	B	202	FLC	OA2-CAC-CA	2.68	122.95	114.35
4	C	302	FLC	OA2-CAC-CA	2.62	122.77	114.35
4	D	302	FLC	OA1-CAC-CA	-2.56	115.47	122.94
4	C	302	FLC	OA1-CAC-CA	-2.47	115.73	122.94
4	B	202	FLC	OA1-CAC-CA	-2.44	115.80	122.94
4	C	303	FLC	OG2-CGC-CG	2.26	121.59	114.35
4	B	201	FLC	OG2-CGC-CG	2.24	121.53	114.35
4	C	302	FLC	OG2-CGC-CG	2.13	121.19	114.35
4	C	303	FLC	OG1-CGC-CG	-2.11	116.79	122.94
4	B	201	FLC	OG1-CGC-CG	-2.10	116.80	122.94
4	C	302	FLC	OG1-CGC-CG	-2.05	116.97	122.94

There are no chirality outliers.

All (43) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	202	FLC	CG-CB-CBC-OB1
4	A	202	FLC	CG-CB-CBC-OB2
4	A	202	FLC	OHB-CB-CBC-OB1
4	A	202	FLC	OHB-CB-CBC-OB2
4	B	202	FLC	CAC-CA-CB-CBC
4	B	202	FLC	CAC-CA-CB-CG
4	B	202	FLC	CAC-CA-CB-OHB
4	B	202	FLC	CG-CB-CBC-OB1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	202	FLC	CG-CB-CBC-OB2
4	B	202	FLC	OHB-CB-CBC-OB1
4	B	202	FLC	OHB-CB-CBC-OB2
4	B	202	FLC	CA-CB-CG-CGC
4	B	202	FLC	CBC-CB-CG-CGC
4	B	202	FLC	OHB-CB-CG-CGC
4	C	302	FLC	CG-CB-CBC-OB1
4	C	302	FLC	CG-CB-CBC-OB2
4	C	302	FLC	OHB-CB-CBC-OB1
4	C	302	FLC	OHB-CB-CBC-OB2
4	C	303	FLC	CAC-CA-CB-CBC
4	C	303	FLC	CAC-CA-CB-CG
4	C	303	FLC	CAC-CA-CB-OHB
4	C	303	FLC	CA-CB-CBC-OB1
4	C	303	FLC	CA-CB-CBC-OB2
4	C	303	FLC	OHB-CB-CBC-OB1
4	C	303	FLC	OHB-CB-CBC-OB2
4	D	301	FLC	CA-CB-CBC-OB1
4	D	301	FLC	CA-CB-CBC-OB2
4	D	301	FLC	OHB-CB-CBC-OB1
4	D	301	FLC	OHB-CB-CBC-OB2
4	B	201	FLC	CAC-CA-CB-CBC
4	B	201	FLC	CAC-CA-CB-OHB
4	C	303	FLC	CB-CA-CAC-OA1
4	C	303	FLC	CB-CA-CAC-OA2
4	B	201	FLC	CAC-CA-CB-CG
4	B	202	FLC	CA-CB-CBC-OB1
4	B	202	FLC	CA-CB-CBC-OB2
4	C	302	FLC	CB-CG-CGC-OG2
4	B	202	FLC	CB-CA-CAC-OA1
4	B	202	FLC	CB-CA-CAC-OA2
4	C	302	FLC	CB-CG-CGC-OG1
4	B	201	FLC	CB-CG-CGC-OG2
4	A	202	FLC	CA-CB-CBC-OB2
4	B	201	FLC	CB-CG-CGC-OG1

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	202	FLC	1	0
4	D	301	FLC	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	302	FLC	1	0
4	C	303	FLC	1	0
4	C	302	FLC	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	109/109 (100%)	0.17	2 (1%)	68	77	30, 44, 81, 105	0
1	B	109/109 (100%)	0.24	3 (2%)	53	64	32, 45, 78, 96	0
2	C	282/288 (97%)	0.22	16 (5%)	23	34	36, 60, 104, 135	0
2	D	284/288 (98%)	0.13	13 (4%)	32	45	39, 61, 98, 122	0
All	All	784/794 (98%)	0.18	34 (4%)	35	47	30, 56, 97, 135	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	252	GLU	5.9
2	C	264	SER	4.5
2	C	287	CYS	4.4
2	C	288	ASN	4.4
1	A	9	HIS	4.3
2	C	266	CYS	4.0
2	D	94	SER	4.0
2	C	242	ASP	3.8
1	B	49[A]	TYR	3.8
2	C	74	GLY	3.5
2	D	74	GLY	3.5
1	B	67	ARG	3.4
2	D	168	PRO	3.3
2	D	73	PRO	3.1
2	C	262	TYR	3.0
2	C	244	LEU	3.0
2	C	73	PRO	2.8
1	A	26	ALA	2.7
2	C	246	PRO	2.6
2	C	81	LYS	2.5
2	D	95	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	165	ARG	2.5
2	C	263	ALA	2.5
2	D	288	ASN	2.5
2	C	251	ASP	2.5
2	D	76	LYS	2.4
2	D	251	ASP	2.4
2	D	264	SER	2.3
2	D	88	VAL	2.3
2	C	253	PRO	2.3
1	B	28	GLY	2.3
2	C	66	CYS	2.1
2	D	66	CYS	2.1
2	D	169	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	FLC	C	302	13/13	0.53	0.29	83,93,99,99	0
4	FLC	D	301	13/13	0.69	0.33	86,95,108,110	0
4	FLC	A	202	13/13	0.71	0.33	83,91,98,99	0
4	FLC	C	303	13/13	0.73	0.22	89,97,105,105	0
4	FLC	B	201	13/13	0.73	0.30	86,89,97,102	0
4	FLC	D	302	13/13	0.82	0.27	89,93,96,96	0
4	FLC	B	202	13/13	0.84	0.18	64,84,95,100	0
3	PO4	D	303	5/5	0.90	0.22	90,91,95,99	0
3	PO4	C	301	5/5	0.97	0.12	56,58,59,63	0
3	PO4	A	201	5/5	0.99	0.15	50,53,55,61	0

6.5 Other polymers (i)

There are no such residues in this entry.