



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 04:14 am BST

PDB ID : 2JIH
Title : Crystal Structure of Human ADAMTS-1 catalytic Domain and Cysteine- Rich Domain (complex-form)
Authors : Gerhardt, S.; Hassall, G.; Hawtin, P.; McCall, E.; Flavell, L.; Minshull, C.; Hargreaves, D.; Ting, A.; Paupit, R.A.; Parker, A.E.; Abbott, W.M.
Deposited on : 2007-06-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	300	
1	B	300	

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 4667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADAMTS-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	284	Total 2181	C 1354	N 379	O 423	S 25	0	0	0
1	B	286	Total 2199	C 1363	N 385	O 426	S 25	0	0	0

- Molecule 2 is (2S,3R)-N 4 -[(1S)-2,2-dimethyl-1-(methylcarbamoyl)propyl]-N 1 ,2-dihydroxy-3-(2-methylpropyl)butanediamide (three-letter code: 097) (formula: C₁₅H₂₉N₃O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 23	C 15	N 3	O 5	0	0
2	B	1	Total 23	C 15	N 3	O 5	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Cd	0	0
			2	2		
4	A	4	Total	Cd	0	0
			4	4		

- Molecule 5 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	5	Total	Ni	0	0
			5	5		
5	A	7	Total	Ni	0	0
			7	7		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	4	Total	Mg	0	0
			4	4		
6	A	2	Total	Mg	0	0
			2	2		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total	Na	0	0
			2	2		
7	A	3	Total	Na	0	0
			3	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	122	Total	O	0	0
			122	122		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	88	Total	O	0	0
			88	88		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADAMTS-1

Chain A: 



- Molecule 1: ADAMTS-1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.58Å 64.40Å 113.52Å 90.00° 90.91° 90.00°	Depositor
Resolution (Å)	56.90 – 2.10 56.75 – 2.09	Depositor EDS
% Data completeness (in resolution range)	82.6 (56.90-2.10) 79.4 (56.75-2.09)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.224 , 0.282 0.281 , 0.297	Depositor DCC
R_{free} test set	1792 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtrriage
Anisotropy	0.494	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.107 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4667	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, ZN, NA, MG, 097, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/2233	0.70	1/3032 (0.0%)
1	B	0.62	0/2251	0.70	1/3055 (0.0%)
All	All	0.63	0/4484	0.70	2/6087 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	483	ASP	CB-CG-OD2	7.15	124.74	118.30
1	A	524	LYS	CD-CE-NZ	5.31	123.91	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	499	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2181	0	2046	26	0
1	B	2199	0	2060	21	1
2	A	23	0	29	5	0
2	B	23	0	28	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	2	0	0	0	0
5	A	7	0	0	0	0
5	B	5	0	0	0	0
6	A	2	0	0	0	0
6	B	4	0	0	0	0
7	A	3	0	0	0	0
7	B	2	0	0	0	0
8	A	122	0	0	1	0
8	B	88	0	0	0	0
All	All	4667	0	4163	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:LEU:CD1	2:A:1001:097:H112	2.07	0.85
1:A:394:GLN:NE2	1:A:526:PHE:CD1	2.59	0.71
1:B:267:ASP:HA	1:B:312:ILE:HD12	1.80	0.64
1:A:434:LEU:HD12	2:A:1001:097:H112	1.79	0.64
1:B:362:CYS:HA	1:B:367:CYS:HA	1.81	0.62
1:A:392:GLY:HA2	1:A:490:PHE:HB3	1.82	0.61
1:A:297:SER:HB3	1:A:441:GLN:HE21	1.67	0.59
1:B:487:GLN:HE22	1:B:529:ALA:H	1.52	0.57
1:A:500:PRO:HD3	1:A:510:TRP:HB2	1.88	0.55
1:B:373:ALA:HB2	1:B:386:SER:HB3	1.92	0.52
1:B:341:PRO:O	1:B:349:HIS:ND1	2.29	0.51
1:A:368:ASP:HB3	2:A:1001:097:H92	1.94	0.50
1:B:441:GLN:HG3	1:B:441:GLN:O	2.14	0.48
1:B:519:LEU:O	1:B:520:VAL:HG23	2.13	0.48
1:A:434:LEU:HD12	2:A:1001:097:C11	2.43	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:411:HIS:HA	1:B:430:MET:O	2.14	0.47
1:B:329:LEU:HD22	1:B:372:MET:HG2	1.97	0.46
1:A:461:GLU:HB3	8:A:2079:HOH:O	2.15	0.46
1:A:286:PHE:CZ	1:A:304:LEU:HB3	2.50	0.46
1:A:394:GLN:NE2	1:A:526:PHE:CE1	2.81	0.46
1:A:524:LYS:CE	1:A:526:PHE:HE1	2.29	0.46
1:B:392:GLY:HA2	1:B:490:PHE:HB3	1.98	0.46
1:A:324:ASN:ND2	1:A:327:LEU:HB2	2.31	0.45
1:A:408:ASN:HB2	1:A:454:PHE:CD2	2.52	0.45
1:A:524:LYS:HE3	1:A:526:PHE:HE1	1.80	0.45
1:A:308:LYS:NZ	1:A:310:LEU:HD21	2.32	0.45
1:A:338:GLN:NE2	1:A:339:HIS:NE2	2.65	0.45
1:B:263:MET:SD	1:B:308:LYS:HD3	2.57	0.45
1:B:498:HIS:CE1	1:B:500:PRO:HA	2.51	0.45
1:A:362:CYS:HA	1:A:367:CYS:HA	1.98	0.45
1:A:486:ARG:HD3	1:B:280:HIS:HB2	2.00	0.44
1:B:477:LEU:HD13	1:B:530:ASP:HB3	2.00	0.43
1:A:499:CYS:HB3	1:A:500:PRO:CD	2.48	0.43
1:B:305:VAL:HG21	1:B:467:PRO:HG3	2.01	0.43
1:A:524:LYS:HE3	1:A:526:PHE:CE1	2.54	0.43
1:A:333:CYS:SG	1:A:383:ARG:HA	2.59	0.43
1:B:314:ASP:HB3	1:B:317:LYS:HG2	2.01	0.43
1:A:434:LEU:HD12	2:A:1001:097:O2	2.18	0.42
1:B:473:LEU:HA	1:B:474:PRO:HD3	1.92	0.41
1:A:299:ARG:HG2	1:A:448:ALA:HB1	2.02	0.41
1:B:499:CYS:O	1:B:500:PRO:C	2.58	0.41
1:A:375:VAL:HA	1:A:405:HIS:O	2.20	0.41
1:B:489:GLN:HA	1:B:493:GLY:O	2.21	0.40
1:B:288:VAL:HB	1:B:396:ALA:HB1	2.04	0.40
1:A:531:GLY:HA2	1:A:539:TRP:CE3	2.57	0.40
1:B:446:CYS:O	1:B:450:MET:HG2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:456:ASP:OD2	1:B:550:VAL:CG2[2_656]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/300 (92%)	268 (97%)	6 (2%)	2 (1%)	22	18
1	B	278/300 (93%)	267 (96%)	10 (4%)	1 (0%)	34	32
All	All	554/600 (92%)	535 (97%)	16 (3%)	3 (0%)	29	26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	499	CYS
1	A	519	LEU
1	B	506	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/259 (95%)	236 (96%)	11 (4%)	27	27
1	B	248/259 (96%)	236 (95%)	12 (5%)	25	24
All	All	495/518 (96%)	472 (95%)	23 (5%)	27	26

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	303	SER
1	A	338	GLN
1	A	364	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	365	GLN
1	A	419	SER
1	A	426	ASP
1	A	434	LEU
1	A	469	ASN
1	A	505	THR
1	A	536	GLU
1	A	539	TRP
1	B	316	GLN
1	B	317	LYS
1	B	343	SER
1	B	352	THR
1	B	382	SER
1	B	426	ASP
1	B	435	SER
1	B	441	GLN
1	B	476	ASP
1	B	481	SER
1	B	520	VAL
1	B	539	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	324	ASN
1	A	338	GLN
1	A	359	GLN
1	A	441	GLN
1	A	457	ASN
1	A	469	ASN
1	B	268	GLN
1	B	408	ASN
1	B	487	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 31 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	097	B	1001	3	22,22,22	0.94	1 (4%)	25,31,31	1.08	1 (4%)
2	097	A	1001	3	22,22,22	1.02	2 (9%)	25,31,31	1.49	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	097	B	1001	3	-	2/34/34/34	-
2	097	A	1001	3	-	2/34/34/34	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	097	C10-N2	2.47	1.36	1.33
2	A	1001	097	C12-N3	2.05	1.36	1.33
2	B	1001	097	C10-N2	2.03	1.35	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	1001	097	C11-N2-C10	3.82	128.82	122.22
2	A	1001	097	C14-C4-N1	3.54	120.22	116.00
2	B	1001	097	C13-C12-N3	2.89	119.83	116.19
2	A	1001	097	O1-C4-C14	-2.66	118.52	121.73
2	A	1001	097	C6-C5-C10	-2.40	110.55	112.81

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1001	097	O4-C12-C13-O5
2	A	1001	097	N3-C12-C13-O5
2	B	1001	097	N3-C12-C13-O5
2	A	1001	097	O4-C12-C13-O5


There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	097	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.