

wwPDB EM Validation Summary Report (i)

Dec 13, 2022 – 04:59 PM JST

PDB ID : 6JMQ EMDB ID : EMD-9849

Title : LAT1-CD98hc complex bound to MEM-108 Fab

Authors: Lee, Y.; Nishizawa, T.; Kusakizako, T.; Oda, K.; Ishitani, R.; Nakane, T.;

Nureki, O.

Deposited on : 2019-03-13

Resolution : 3.31 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 ${\it MapQ}$: ${\it FAILED}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	515	69%	9%	22%				
2	В	631	64%	8%	29%				
3	С	219	88%		10% •				
4	D	219	89%		11%				
5	Е	2	100%						



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 9950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Large neutral amino acids transporter small subunit 1.

\mathbf{Mol}	Chain	Residues	Atoms					AltConf	Trace	
1	A	403	Total 3101	C 2093	N 468	O 525	S 15	0	0	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	ASP	-	expression tag	UNP Q01650
A	509	TYR	-	expression tag	UNP Q01650
A	510	LYS	-	expression tag	UNP Q01650
A	511	ASP	-	expression tag	UNP Q01650
A	512	ASP	-	expression tag	UNP Q01650
A	513	ASP	-	expression tag	UNP Q01650
A	514	ASP	-	expression tag	UNP Q01650
A	515	LYS	-	expression tag	UNP Q01650

• Molecule 2 is a protein called 4F2 cell-surface antigen heavy chain.

Mol	Chain	Residues	Atoms				AltConf	Trace	
2	В	451	Total 3520	C 2248	N 605	O 660	S 7	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference	
В	0	GLY	-	expression tag	UNP P08195	
В	1	SER	-	expression tag	UNP P08195	

• Molecule 3 is a protein called Antibody.

Mol	Chain	Residues	Atoms			AltConf	Trace		
3	С	214	Total 1526	C 960	N 260	O 300	S 6	0	0



• Molecule 4 is a protein called Antibody.


Mol	Chain	Residues	Atoms				AltConf	Trace	
4	D	210	Total	С	N	О	S	0	0
4	D	219	1593	992	268	326	7	U	U

• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			AltConf	Trace	
5	Е	2	Total 28	C 16	N 2	O 10	0	0

 \bullet Molecule 6 is CHOLESTEROL (three-letter code: CLR) (formula: $\mathrm{C_{27}H_{46}O}).$



Mol	Chain	Residues	Atoms	AltConf
6	A	1	Total C O 112 108 4	0
6	A	1	Total C O 112 108 4	0
6	A	1	Total C O 112 108 4	0
6	A	1	Total C O 112 108 4	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
6	В	1	Total C O 28 27 1	0

 \bullet Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$




Mol	Chain	Residues	Ato		AltConf	
7	В	1	Total C 42 24			0
7	В	1	Total C 42 24		_	0
7	В	1	Total C 42 24			0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Large neutral amino acids transporter small subunit 1



• Molecule 4: Antibody

Chain D: 89% 11%



 \bullet Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	250712	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	7.14	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.29	0/3184	0.54	1/4356~(0.0%)	
2	В	0.29	0/3601	0.53	0/4886	
3	С	0.31	0/1403	0.55	0/1914	
4	D	0.29	0/1456	0.50	0/1972	
All	All	0.29	0/9644	0.53	1/13128 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	435	LEU	CA-CB-CG	5.23	127.33	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3101	0	3214	25	0
2	В	3520	0	3500	27	0
3	С	1526	0	1403	14	0
4	D	1593	0	1413	12	0
5	Е	28	0	25	0	0
6	A	112	0	184	2	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	В	28	0	46	1	0
7	В	42	0	39	0	0
All	All	9950	0	9824	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:B:361:ALA:HA	2:B:395:ASN:HD21	1.60	0.66
1:A:106:LEU:HD13	1:A:415:MET:HB3	1.80	0.62
2:B:600:GLN:HB2	2:B:620:GLU:HG3	1.82	0.62
1:A:141:ARG:HE	1:A:346:SER:HB2	1.68	0.59
2:B:430:TRP:HE1	2:B:463:THR:HG23	1.69	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	397/515~(77%)	378 (95%)	18 (4%)	1 (0%)	41	71
2	В	$449/631 \ (71\%)$	427 (95%)	22 (5%)	0	100	100
3	С	179/219~(82%)	170 (95%)	9 (5%)	0	100	100
4	D	184/219~(84%)	181 (98%)	3 (2%)	0	100	100
All	All	1209/1584~(76%)	1156 (96%)	52 (4%)	1 (0%)	54	81

All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	428	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	336/427 (79%)	334 (99%)	2 (1%)	86	92
2	В	378/514 (74%)	378 (100%)	0	100	100
3	С	162/162 (100%)	162 (100%)	0	100	100
4	D	165/166 (99%)	165 (100%)	0	100	100
All	All	1041/1269 (82%)	1039 (100%)	2 (0%)	93	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	TRP
1	A	454	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
3	С	86	GLN
4	D	130	GLN
4	D	144	ASN
2	В	395	ASN
2	В	542	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain		Chain	Res	Tiple	Bo	ond leng	ths	В	ond ang	les
IVIOI	туре	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	Е	1	2,5	14,14,15	0.22	0	17,19,21	0.50	0
5	NAG	Е	2	5	14,14,15	0.25	0	17,19,21	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	Ε	1	2,5	-	1/6/23/26	0/1/1/1
5	NAG	Е	2	5	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:


\mathbf{Mol}	Chain	Res	Type	Atoms
5	Ε	2	NAG	O5-C5-C6-O6
5	Ε	2	NAG	C4-C5-C6-O6
5	Ε	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	$ ag{ths}$	В	ond ang	les
WIOI	Туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	CLR	A	601	-	31,31,31	0.34	0	48,48,48	0.47	0



Mol	Tuno	Chain	Dec	Res Link	Вс	Bond lengths			Bond angles		
MIOI	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
6	CLR	A	602	-	31,31,31	0.29	0	48,48,48	0.48	0	
6	CLR	В	701	-	31,31,31	0.33	0	48,48,48	0.48	0	
6	CLR	A	604	-	31,31,31	0.32	0	48,48,48	0.46	0	
7	NAG	В	705	2	14,14,15	0.36	0	17,19,21	0.43	0	
7	NAG	В	704	2	14,14,15	1.46	2 (14%)	17,19,21	1.36	1 (5%)	
6	CLR	A	603	-	31,31,31	0.33	0	48,48,48	0.53	0	
7	NAG	В	706	2	14,14,15	0.48	0	17,19,21	0.63	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CLR	A	601	-	-	0/10/68/68	0/4/4/4
6	CLR	A	602	-	-	4/10/68/68	0/4/4/4
6	CLR	В	701	ı	-	4/10/68/68	0/4/4/4
6	CLR	A	604	-	-	4/10/68/68	0/4/4/4
7	NAG	В	705	2	-	2/6/23/26	0/1/1/1
7	NAG	В	704	2	-	2/6/23/26	0/1/1/1
6	CLR	A	603	-	-	0/10/68/68	0/4/4/4
7	NAG	В	706	2	-	4/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$ \operatorname{Ideal}(ext{ iny A}) $
7	В	704	NAG	O5-C1	4.61	1.51	1.43
7	В	704	NAG	C1-C2	2.56	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
7	В	704	NAG	C1-O5-C5	5.25	119.31	112.19

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	В	706	NAG	O5-C5-C6-O6
7	В	706	NAG	C4-C5-C6-O6

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	В	705	NAG	C4-C5-C6-O6
7	В	704	NAG	C4-C5-C6-O6
7	В	706	NAG	C8-C7-N2-C2

There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	602	CLR	1	0
6	В	701	CLR	1	0
6	A	603	CLR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

