



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 11:22 PM EDT

PDB ID : 5JON  
Title : Crystal structure of the unliganded form of HCN2 CNBD  
Authors : Klenchin, V.A.; Chanda, B.  
Deposited on : 2016-05-02  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	 2% 94%
1	B	517	 3% 92% 6%
2	C	2	 100%
2	D	2	 50% 50%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8340 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein,Potassium/sodium hyper polarization-activated cyclic nucleotide-gated channel 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	509	Total	C	N	O	S	0	6	0
			3977	2555	652	755	15			
1	B	506	Total	C	N	O	S	0	2	0
			3936	2526	646	749	15			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-369	GLY	-	expression tag	UNP P0AEY0
A	-2	ASN	-	linker	UNP P0AEY0
A	-1	ALA	-	linker	UNP P0AEY0
A	0	ALA	-	linker	UNP P0AEY0
B	-369	GLY	-	expression tag	UNP P0AEY0
B	-2	ASN	-	linker	UNP P0AEY0
B	-1	ALA	-	linker	UNP P0AEY0
B	0	ALA	-	linker	UNP P0AEY0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N O 4 1 3	0	0
3	B	1	Total N O 4 1 3	0	0
3	B	1	Total N O 4 1 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	193	Total O 193 193	0	0
4	B	176	Total O 176 176	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltose-binding periplasmic protein,Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2



- Molecule 1: Maltose-binding periplasmic protein,Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 2



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.52Å 42.01Å 198.38Å 90.00° 90.85° 90.00°	Depositor
Resolution (Å)	24.82 – 2.04 24.82 – 2.07	Depositor EDS
% Data completeness (in resolution range)	95.7 (24.82-2.04) 95.7 (24.82-2.07)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.180 , 0.221 0.187 , 0.223	Depositor DCC
$R_{free}$ test set	3134 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 22.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.186 for h,-k,-l	Xtriage
Reported twinning fraction	0.878 for H, K, L 0.122 for -h,-k,l	Depositor
Outliers	0 of 62510 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8340	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/4089	0.66	1/5546 (0.0%)
1	B	0.48	0/4032	0.65	0/5468
All	All	0.48	0/8121	0.65	1/11014 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-326	LEU	CA-CB-CG	6.05	129.22	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3977	0	3930	10	0
1	B	3936	0	3875	13	0
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	4	0	0	0	0
3	B	8	0	0	0	0
4	A	193	0	0	0	0
4	B	176	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8340	0	7847	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-28:TYR:O	1:A:-24:THR:HG23	2.02	0.59
1:A:-217:GLN:HA	1:A:-21[A]:ILE:HD11	1.85	0.58
1:B:615:LEU:HD21	1:B:625:PHE:CD2	2.42	0.55
1:A:-301:GLY:HA3	1:A:-37:ASN:O	2.07	0.54
1:B:587:THR:HG23	1:B:589:GLY:H	1.72	0.54
1:A:510:LYS:O	1:A:514:SER:OG	2.19	0.54
1:B:507:ASN:OD1	1:B:558:GLN:NE2	2.42	0.53
1:A:-222:LEU:HD23	1:A:-165:MET:CE	2.40	0.50
1:A:-124[A]:THR:HG22	1:A:-123:VAL:H	1.75	0.50
1:A:-185:ASP:HB2	1:A:-4:GLN:HB2	1.94	0.50
1:B:-301:GLY:HA3	1:B:-37:ASN:O	2.15	0.46
1:B:-280:LEU:HD22	1:B:-275:TRP:CZ2	2.50	0.46
1:A:-272:VAL:HG21	1:A:-264:ALA:O	2.17	0.44
1:B:-75:ASN:HD22	1:B:-70:LEU:H	1.64	0.44
1:A:-124[A]:THR:HG22	1:A:-123:VAL:N	2.33	0.43
1:B:584:CYS:HA	1:B:587:THR:HG22	2.01	0.43
1:B:-364:GLY:H	1:B:-97:ASN:HD21	1.66	0.42
1:A:-124[A]:THR:HG22	1:A:-123:VAL:HG12	2.01	0.42
1:B:-128:ASN:ND2	4:B:802:HOH:O	2.51	0.42
1:B:-211:TRP:N	1:B:-210:PRO:CD	2.83	0.42
1:B:-274:ASP:OD1	1:B:-271:ARG:NH1	2.53	0.41
1:B:509:ARG:NH1	1:B:527:THR:OG1	2.53	0.41
1:B:-136:SER:OG	1:B:-71:PRO:HD3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	513/517 (99%)	505 (98%)	8 (2%)	0	100	100
1	B	506/517 (98%)	495 (98%)	11 (2%)	0	100	100
All	All	1019/1034 (98%)	1000 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/423 (98%)	407 (98%)	8 (2%)	57	53
1	B	407/423 (96%)	400 (98%)	7 (2%)	60	57
All	All	822/846 (97%)	807 (98%)	15 (2%)	59	55

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-363	LYS
1	A	-326	LEU
1	A	-323	LYS
1	A	-111	PHE
1	A	-25	ARG
1	A	496	ASN
1	A	558	GLN
1	A	620	MET
1	B	-111	PHE
1	B	-47	GLU
1	B	517	LEU
1	B	553	LYS
1	B	558	GLN
1	B	590	ARG
1	B	623	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-351	ASN
1	A	-269	ASN
1	A	-168	ASN
1	A	496	ASN
1	B	-97	ASN
1	B	-75	ASN
1	B	610	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	C	1	2	12,12,12	0.44	0	17,17,17	1.21	2 (11%)
2	GLC	C	2	2	11,11,12	0.37	0	15,15,17	0.89	1 (6%)
2	GLC	D	1	2	12,12,12	0.43	0	17,17,17	0.86	0
2	GLC	D	2	2	11,11,12	0.25	0	15,15,17	0.87	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	1/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	C1-O5-C5	3.24	119.78	113.66
2	D	2	GLC	C1-O5-C5	2.78	115.95	112.19
2	C	2	GLC	C1-O5-C5	2.70	115.84	112.19
2	C	1	GLC	O5-C1-C2	2.33	114.44	110.28

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	GLC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NO3	A	702	-	1,3,3	0.70	0	0,3,3	-	-
3	NO3	B	702	-	1,3,3	0.57	0	0,3,3	-	-
3	NO3	B	703	-	1,3,3	0.58	0	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	509/517 (98%)	-0.08	12 (2%) 59 63	12, 24, 57, 100	4 (0%)
1	B	506/517 (97%)	-0.05	16 (3%) 47 52	11, 23, 64, 94	5 (0%)
All	All	1015/1034 (98%)	-0.07	28 (2%) 53 58	11, 23, 61, 100	9 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	628	VAL	9.3
1	A	625	PHE	5.4
1	A	517	LEU	4.7
1	B	497	GLY	4.4
1	B	625	PHE	3.9
1	A	-315	GLY	3.5
1	A	633	LEU	3.4
1	B	512	VAL	3.4
1	A	628	VAL	3.2
1	B	569	ASN	2.9
1	A	506	PHE	2.8
1	A	630	ILE	2.8
1	B	624	ALA	2.8
1	B	623	ARG	2.7
1	B	500	ARG	2.7
1	B	495	LEU	2.6
1	B	523	PRO	2.6
1	A	-367	ILE	2.5
1	B	567	LYS	2.5
1	B	520	ASN	2.4
1	A	568	GLY	2.3
1	A	567	LYS	2.2
1	B	502	GLU	2.2
1	B	498	PRO	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	629	ALA	2.2
1	B	506	PHE	2.2
1	A	498	PRO	2.1
1	A	620	MET	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	C	1	12/12	0.97	0.08	14,17,21,21	0
2	GLC	D	1	12/12	0.97	0.09	14,18,21,24	0
2	GLC	C	2	11/12	0.98	0.09	10,11,13,13	0
2	GLC	D	2	11/12	0.99	0.08	11,13,14,15	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NO3	A	702	4/4	0.75	0.36	39,43,44,45	0
3	NO3	B	702	4/4	0.83	0.37	38,40,41,43	0
3	NO3	B	703	4/4	0.90	0.32	36,37,39,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.