Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 04:10 PM EDT

PDB ID	:	4JPH
Title	:	Crystal structure of Protein Related to DAN and Cerberus (PRDC)
Authors	:	Deng, X.; Nolan, K.T.; Kattamuri, C.; Thompson, T.B.
Deposited on	:	2013-03-19
Resolution	:	2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	11% 71% 7% 22%
1	B	148	12% 63% 12% 25%
1	C	148	7% 59% 9% 32%
1	D	148	16% 64% 10% • 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	203	-	-	-	X
2	GOL	A	204	-	-	-	X
2	GOL	B	203	-	-	-	X
3	GSH	A	206	-	-	-	X
3	GSH	C	204	-	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3854 atoms, of which 106 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gremlin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	115	942	589	173	170	10	0	2	0
1	B	111	885	554	161	160	10	0	0	0
1	C	101	795	500	142	143	10	0	0	0
1	D	111	867	545	155	157	10	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	INITIATING METHIONINE	UNP O88273
B	21	MET	-	INITIATING METHIONINE	UNP O88273
C	21	MET	-	INITIATING METHIONINE	UNP O88273
D	21	MET	-	INITIATING METHIONINE	UNP O88273

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	14	3	8	3	0	0
2	A	1	14	3	8	3	0	0
2	A	1	14	3	8	3	0	0
2	A	1	14	3	8	3	0	0
2	A	1	14	3	8	3	0	0
2	B	1	14	3	8	3	0	0
2	B	1	14	3	8	3	0	0
2	B	1	14	3	8	3	0	0
2	C	1	14	3	8	3	0	0
2	C	1	14	3	8	3	0	0
2	C	1	14	3	8	3	0	0
2	D	1	14	3	8	3	0	0

- Molecule 3 is GLUTATHIONE (three-letter code: GSH) (formula: $C_{10}H_{17}N_3O_6S$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O S 20 10 3 6 1	0	0
3	B	1	Total C N O S 14 9 2 2 1	0	0
3	C	1	Total C N O S 8 4 2 1 1	0	0
3	D	1	Total C N S 6 4 1 1	0	0

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	B	1	18	6	5	7	0	0
			Total	C	H	O		
4	D	1	18	6	5	7	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	49	49	49	0	0
			Total	O		
5	B	37	37	37	0	0
			Total	O		
5	C	14	14	14	0	0
			Total	O		
5	D	13	13	13	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Gremlin-2

Chain A:



• Molecule 1: Gremlin-2

Chain B:



• Molecule 1: Gremlin-2

Chain C:



• Molecule 1: Gremlin-2

Chain D:




E114
F122
C123
K124
S130
V131
I132
V133
P138
G139
L140
D141
P142
P143
F144
R145
I146
I149
M158
S159
V160
ASN
LEU
SER
ASP
SER
ASP
LYS
GLN

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	73.31Å 65.56Å 85.08Å 90.00° 105.50° 90.00°	Depositor
Resolution (Å)	17.60 – 2.25 17.60 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.3 (17.60-2.25) 99.3 (17.60-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.86 (at 2.25Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.202 , 0.222 0.211 , 0.233	Depositor DCC
R_{free} test set	1838 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	60.8	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 71.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GSH, CIT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.56	0/961	0.73	0/1294
1	B	0.54	0/902	0.69	0/1214
1	C	0.48	0/812	0.70	0/1097
1	D	0.47	0/884	0.78	0/1191
All	All	0.52	0/3559	0.73	0/4796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	942	0	926	7	0
1	B	885	0	874	9	0
1	C	795	0	766	6	0
1	D	867	0	846	7	0
2	A	30	40	40	1	0
2	B	18	24	24	0	0
2	C	18	24	24	0	0
2	D	6	8	8	0	0
3	A	20	0	14	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	14	0	11	0	0
3	C	8	0	4	0	0
3	D	6	0	4	0	0
4	B	13	5	5	0	0
4	D	13	5	5	0	0
5	A	49	0	0	0	0
5	B	37	0	0	0	0
5	C	14	0	0	0	0
5	D	13	0	0	1	0
All	All	3748	106	3551	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:108:ARG:HB3	1:B:115:ASP:O	1.86	0.76
1:D:124:LYS:HG3	1:D:158:MET:HE2	1.79	0.65
1:B:80:GLN:HE22	1:B:94:ASN:HD22	1.45	0.64
1:D:64:GLU:HA	1:D:67:TYR:HB2	1.77	0.64
1:D:124:LYS:HG3	1:D:158:MET:CE	2.32	0.60
1:A:104:PHE:HE1	1:A:117:PHE:HD2	1.46	0.60
1:A:118[A]:GLN:HE22	3:A:206:GSH:HG13	1.69	0.57
1:B:80:GLN:NE2	1:B:94:ASN:HD22	2.01	0.57
1:A:104:PHE:HE1	1:A:117:PHE:CD2	2.27	0.52
1:D:90:ARG:NH2	5:D:310:HOH:O	2.44	0.49
1:A:98:TYR:CE2	1:B:67:TYR:HB3	2.47	0.49
1:B:132:ILE:HG23	1:B:146:ILE:HG23	1.94	0.48
1:A:146:ILE:HD13	1:C:146:ILE:HD11	1.95	0.48
1:A:145:ARG:HE	2:A:205:GOL:H11	1.78	0.48
1:B:106:ILE:HB	1:B:117:PHE:HB2	1.98	0.46
1:D:132:ILE:HG23	1:D:146:ILE:HG23	1.98	0.46
1:C:110:VAL:HG21	1:D:149:ILE:HD12	1.98	0.45
1:C:149:ILE:HD12	1:D:110:VAL:HG21	1.99	0.45
1:B:111:LYS:HA	1:B:112:LYS:HA	1.79	0.44
1:B:68:LEU:HD13	1:B:102:ASN:HB3	2.00	0.44
1:C:90:ARG:HD3	1:C:134:GLU:O	2.18	0.43
1:A:87:CYS:HA	1:A:137:CYS:HA	1.99	0.43
1:C:127:ARG:HG2	1:C:153:LYS:HD3	2.01	0.43
1:B:63:THR:HA	1:B:67:TYR:HB2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:132:ILE:HG23	1:C:146:ILE:HG23	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/148 (78%)	108 (94%)	7 (6%)	0	100	100
1	B	109/148 (74%)	106 (97%)	3 (3%)	0	100	100
1	C	99/148 (67%)	95 (96%)	2 (2%)	2 (2%)	7	4
1	D	109/148 (74%)	101 (93%)	6 (6%)	2 (2%)	8	4
All	All	432/592 (73%)	410 (95%)	18 (4%)	4 (1%)	17	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	144	PHE
1	C	115	ASP
1	C	112	LYS
1	D	110	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/138 (78%)	106 (98%)	2 (2%)	57	66
1	B	101/138 (73%)	97 (96%)	4 (4%)	31	37
1	C	90/138 (65%)	86 (96%)	4 (4%)	28	32
1	D	97/138 (70%)	89 (92%)	8 (8%)	11	10
All	All	396/552 (72%)	378 (96%)	18 (4%)	27	31

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LYS
1	A	130	SER
1	B	53	LEU
1	B	65	ARG
1	B	114	GLU
1	B	130	SER
1	C	68	LEU
1	C	69	LYS
1	C	78	LEU
1	C	130	SER
1	D	53	LEU
1	D	67	TYR
1	D	78	LEU
1	D	81	THR
1	D	91	THR
1	D	93	LEU
1	D	130	SER
1	D	141	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	80	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GSH	A	206	1	18,19,19	2.28	3 (16%)	21,24,24	1.24	2 (9%)
4	CIT	B	201	-	12,12,12	0.32	0	17,17,17	0.54	0
2	GOL	A	204	-	5,5,5	0.07	0	5,5,5	0.14	0
4	CIT	D	201	-	12,12,12	0.26	0	17,17,17	0.35	0
2	GOL	A	201	-	5,5,5	0.08	0	5,5,5	0.28	0
2	GOL	B	202	-	5,5,5	0.26	0	5,5,5	0.49	0
2	GOL	A	202	-	5,5,5	0.08	0	5,5,5	0.21	0
2	GOL	B	203	-	5,5,5	0.09	0	5,5,5	0.35	0
2	GOL	C	201	-	5,5,5	0.21	0	5,5,5	0.32	0
3	GSH	C	204	1	6,7,19	1.80	2 (33%)	4,7,24	1.37	0
2	GOL	C	203	-	5,5,5	0.09	0	5,5,5	0.17	0
3	GSH	B	205	1	13,13,19	2.53	3 (23%)	14,15,24	1.31	3 (21%)
2	GOL	A	205	-	5,5,5	0.16	0	5,5,5	0.74	0
2	GOL	A	203	-	5,5,5	0.10	0	5,5,5	0.24	0
2	GOL	D	202	-	5,5,5	0.09	0	5,5,5	0.19	0
2	GOL	C	202	-	5,5,5	0.05	0	5,5,5	0.14	0
2	GOL	B	204	-	5,5,5	0.11	0	5,5,5	0.23	0
3	GSH	D	203	1	5,5,19	1.07	0	4,5,24	3.14	2 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GSH	A	206	1	-	1/24/24/24	-
4	CIT	B	201	-	-	0/16/16/16	-
2	GOL	A	204	-	-	0/4/4/4	-
4	CIT	D	201	-	-	0/16/16/16	-
2	GOL	A	201	-	-	2/4/4/4	-
2	GOL	B	202	-	-	0/4/4/4	-
2	GOL	A	202	-	-	0/4/4/4	-
2	GOL	B	203	-	-	0/4/4/4	-
2	GOL	C	201	-	-	2/4/4/4	-
3	GSH	C	204	1	-	1/6/7/24	-
2	GOL	C	203	-	-	0/4/4/4	-
3	GSH	B	205	1	-	6/16/16/24	-
2	GOL	A	205	-	-	4/4/4/4	-
2	GOL	A	203	-	-	0/4/4/4	-
2	GOL	D	202	-	-	0/4/4/4	-
2	GOL	C	202	-	-	0/4/4/4	-
2	GOL	B	204	-	-	0/4/4/4	-
3	GSH	D	203	1	-	3/3/4/24	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	206	GSH	C2-N3	6.40	1.48	1.33
3	B	205	GSH	C2-N3	6.34	1.48	1.33
3	B	205	GSH	CD1-N2	5.90	1.46	1.34
3	A	206	GSH	CD1-N2	5.77	1.46	1.34
3	C	204	GSH	CD1-N2	3.62	1.45	1.33
3	A	206	GSH	CB2-CA2	-2.20	1.50	1.53
3	B	205	GSH	CB2-CA2	-2.09	1.50	1.53
3	C	204	GSH	CA2-N2	-2.08	1.45	1.47

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	203	GSH	CD1-N2-CA2	5.70	121.20	114.39
3	B	205	GSH	CB2-CA2-N2	2.79	115.26	111.28
3	A	206	GSH	O12-C1-O11	-2.34	118.77	124.08
3	A	206	GSH	CA2-C2-N3	2.28	121.43	116.54
3	D	203	GSH	CA2-CB2-SG2	-2.23	111.65	114.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	205	GSH	CA2-N2-CD1	2.04	126.81	121.68
3	B	205	GSH	CA2-C2-N3	2.03	120.89	116.54

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	GOL	C1-C2-C3-O3
2	A	205	GOL	C1-C2-C3-O3
2	A	205	GOL	O2-C2-C3-O3
2	C	201	GOL	C1-C2-C3-O3
3	B	205	GSH	CA1-CB1-CG1-CD1
3	B	205	GSH	N2-CA2-CB2-SG2
3	B	205	GSH	C2-CA2-CB2-SG2
3	C	204	GSH	OE1-CD1-N2-CA2
3	D	203	GSH	CB2-CA2-N2-CD1
3	A	206	GSH	CA1-CB1-CG1-CD1
2	A	205	GOL	O1-C1-C2-C3
2	A	201	GOL	O2-C2-C3-O3
3	B	205	GSH	O2-C2-CA2-CB2
3	D	203	GSH	C2-CA2-N2-CD1
3	B	205	GSH	N3-C2-CA2-CB2
2	C	201	GOL	O2-C2-C3-O3
2	A	205	GOL	O1-C1-C2-O2
3	D	203	GSH	N2-CA2-CB2-SG2
3	B	205	GSH	C2-CA2-N2-CD1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	206	GSH	1	0
2	A	205	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand GSH A 206



Ligand GSH C 204







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	115/148 (77%)	0.70	17 (14%)	2	2	44, 69, 111, 148	0
1	B	111/148 (75%)	0.76	18 (16%)	1	1	46, 62, 119, 142	0
1	C	101/148 (68%)	0.65	11 (10%)	5	5	49, 80, 132, 147	0
1	D	111/148 (75%)	1.13	23 (20%)	1	1	65, 95, 134, 149	0
All	All	438/592 (73%)	0.81	69 (15%)	2	1	44, 77, 130, 149	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	138	PRO	7.0
1	A	46	HIS	6.8
1	B	160	VAL	6.1
1	C	117	PHE	5.7
1	A	127	ARG	4.9
1	B	110	VAL	4.7
1	B	111	LYS	4.7
1	B	50	LYS	4.6
1	C	160	VAL	4.6
1	B	113	GLU	4.6
1	C	96	PHE	4.2
1	D	50	LYS	4.1
1	D	141	ASP	4.1
1	B	106	ILE	4.1
1	C	60	LEU	4.1
1	D	139	GLY	4.1
1	D	142	PRO	4.0
1	D	160	VAL	3.9
1	B	51	GLU	3.8
1	A	47	HIS	3.8
1	B	115	ASP	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	113	GLU	3.6
1	A	112	LYS	3.6
1	B	107	PRO	3.6
1	D	114	GLU	3.4
1	D	140	LEU	3.4
1	A	96	PHE	3.4
1	C	139	GLY	3.3
1	D	109	HIS	3.2
1	A	122	PHE	3.2
1	D	70	SER	3.2
1	B	112	LYS	3.1
1	D	123	CYS	3.1
1	D	79	ARG	3.1
1	D	55	SER	3.0
1	A	48	GLN	2.9
1	B	152	VAL	2.9
1	C	78	LEU	2.9
1	C	144	PHE	2.8
1	A	160	VAL	2.8
1	A	51	GLU	2.8
1	B	144	PHE	2.7
1	A	69	LYS	2.6
1	D	67	TYR	2.6
1	C	109	HIS	2.6
1	C	122	PHE	2.6
1	B	135	LEU	2.5
1	A	97	CYS	2.4
1	B	114	GLU	2.4
1	A	78	LEU	2.3
1	D	93	LEU	2.3
1	B	105	TYR	2.3
1	C	115	ASP	2.3
1	D	133	VAL	2.3
1	D	111	LYS	2.3
1	A	156	ARG	2.3
1	A	49	ILE	2.3
1	D	99	GLY	2.2
1	B	123	CYS	2.2
1	D	68	LEU	2.2
1	B	146	ILE	2.2
1	D	122	PHE	2.2
1	C	159	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	103	SER	2.1
1	A	107	PRO	2.1
1	A	123	CYS	2.0
1	B	65	ARG	2.0
1	D	51	GLU	2.0
1	D	112	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GOL	A	204	6/6	0.65	0.52	137,138,138,138	0
3	GSH	A	206	20/20	0.65	0.44	114,128,133,134	0
3	GSH	C	204	8/20	0.69	0.56	123,125,129,129	0
2	GOL	A	203	6/6	0.71	0.53	114,121,122,122	0
4	CIT	D	201	13/13	0.71	0.36	160,164,166,166	0
3	GSH	D	203	6/20	0.78	0.26	111,113,113,113	0
2	GOL	C	201	6/6	0.79	0.25	82,90,91,92	0
2	GOL	C	203	6/6	0.79	0.35	103,109,111,111	0
2	GOL	B	203	6/6	0.80	0.50	134,136,137,137	0
2	GOL	B	204	6/6	0.82	0.20	83,86,91,91	0
2	GOL	D	202	6/6	0.82	0.33	114,115,118,118	0
2	GOL	C	202	6/6	0.83	0.44	150,151,152,152	0
2	GOL	A	201	6/6	0.84	0.19	119,120,122,122	0
2	GOL	A	202	6/6	0.84	0.22	124,124,124,125	0
3	GSH	B	205	14/20	0.85	0.31	103,119,123,124	0
2	GOL	B	202	6/6	0.85	0.12	79,84,87,87	0
2	GOL	A	205	6/6	0.87	0.24	78,83,91,92	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	CIT	B	201	13/13	0.91	0.13	83,93,99,100	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GSH A 206:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around GSH C 204:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GSH D 203:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GSH B 205:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.