



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 06:15 PM EST

PDB ID : 4JQ5  
Title : Crystal structure of the human Nup49CCS2+3\* coiled-coil segment  
Authors : Stuwe, T.; Bley, C.J.; Mayo, D.J.; Hoelz, A.  
Deposited on : 2013-03-20  
Resolution : 2.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	86	12% (Poor fit) 81% (0-1 outliers) 17% (2-3 outliers)
1	B	86	7% (Poor fit) 85% (0-1 outliers) 14% (2-3 outliers)
1	C	86	14% (Poor fit) 76% (0-1 outliers) 21% (2-3 outliers)
1	D	86	8% (Poor fit) 86% (0-1 outliers) 13% (2-3 outliers)
1	E	86	12% (Poor fit) 79% (0-1 outliers) 21% (2-3 outliers)

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	86	<p>9% 83% 17%</p>
1	G	86	<p>15% 90% 10%</p>
1	H	86	<p>7% 84% 15%</p>
1	I	86	<p>13% 79% 20%</p>
1	J	86	<p>10% 85% 13%</p>
1	K	86	<p>8% 83% 17%</p>
1	L	86	<p>9% 87% 13%</p>

## 2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 17026 atoms, of which 8500 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoporin p58/p45.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	85	1413	452	706	122	130	3	0	0	0
1	B	85	1413	452	706	122	130	3	0	0	0
1	C	85	1413	452	706	122	130	3	0	0	0
1	D	85	1413	452	706	122	130	3	0	0	0
1	E	86	1423	455	710	123	132	3	0	0	0
1	F	86	1423	455	710	123	132	3	0	0	0
1	G	86	1423	455	710	123	132	3	0	0	0
1	H	85	1413	452	706	122	130	3	0	0	0
1	I	86	1423	455	710	123	132	3	0	0	0
1	J	86	1423	455	710	123	132	3	0	0	0
1	K	86	1423	455	710	123	132	3	0	0	0
1	L	86	1423	455	710	123	132	3	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	SER	-	expression tag	UNP Q9BVL2
B	326	SER	-	expression tag	UNP Q9BVL2
C	326	SER	-	expression tag	UNP Q9BVL2
D	326	SER	-	expression tag	UNP Q9BVL2
E	326	SER	-	expression tag	UNP Q9BVL2

*Continued on next page...*


*Continued from previous page...*

<b>Chain</b>	<b>Residue</b>	<b>Modelled</b>	<b>Actual</b>	<b>Comment</b>	<b>Reference</b>
F	326	SER	-	expression tag	UNP Q9BVL2
G	326	SER	-	expression tag	UNP Q9BVL2
H	326	SER	-	expression tag	UNP Q9BVL2
I	326	SER	-	expression tag	UNP Q9BVL2
J	326	SER	-	expression tag	UNP Q9BVL2
K	326	SER	-	expression tag	UNP Q9BVL2
L	326	SER	-	expression tag	UNP Q9BVL2


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleoporin p58/p45




- Molecule 1: Nucleoporin p58/p45




- Molecule 1: Nucleoporin p58/p45





- Molecule 1: Nucleoporin p58/p45




- Molecule 1: Nucleoporin p58/p45





- Molecule 1: Nucleoporin p58/p45




- Molecule 1: Nucleoporin p58/p45





- Molecule 1: Nucleoporin p58/p45




- Molecule 1: Nucleoporin p58/p45





- Molecule 1: Nucleoporin p58/p45



- Molecule 1: Nucleoporin p58/p45



- Molecule 1: Nucleoporin p58/p45



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.82Å 77.10Å 77.20Å 120.04° 90.12° 89.96°	Depositor
Resolution (Å)	19.83 – 2.19 19.83 – 2.19	Depositor EDS
% Data completeness (in resolution range)	97.0 (19.83-2.19) 89.6 (19.83-2.19)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 2.19Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, $R_{free}$	0.285 , 0.293 0.287 , 0.292	Depositor DCC
$R_{free}$ test set	1990 reflections (3.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.3	Xtriage
Anisotropy	0.604	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 23.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.458 for h,-l,k+1 0.458 for h,k+1,-k 0.457 for h,-k-l,k 0.457 for h,l,-k-l 0.487 for h,-k,-l 0.458 for -h,k,-k-l 0.460 for -h,l,k 0.477 for -h,-k-l,l 0.477 for -h,k+1,-l 0.459 for -h,-k,k+1 0.459 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	17026	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/720	0.36	0/972
1	B	0.23	0/720	0.37	0/972
1	C	0.23	0/720	0.35	0/972
1	D	0.24	0/720	0.38	0/972
1	E	0.22	0/726	0.34	0/980
1	F	0.23	0/726	0.36	0/980
1	G	0.23	0/726	0.35	0/980
1	H	0.23	0/720	0.35	0/972
1	I	0.22	0/726	0.34	0/980
1	J	0.23	0/726	0.37	0/980
1	K	0.23	0/726	0.34	0/980
1	L	0.22	0/726	0.35	0/980
All	All	0.23	0/8682	0.36	0/11720

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	707	706	702	8	0
1	B	707	706	702	3	0
1	C	707	706	702	6	0
1	D	707	706	702	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	713	710	707	11	0
1	F	713	710	707	10	0
1	G	713	710	707	5	0
1	H	707	706	702	6	0
1	I	713	710	707	13	0
1	J	713	710	707	9	0
1	K	713	710	707	11	0
1	L	713	710	707	10	0
All	All	8526	8500	8459	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:326:SER:OG	1:L:329:ALA:CB	2.34	0.75
1:L:326:SER:OG	1:L:329:ALA:HB3	1.89	0.72
1:B:362:ASN:O	1:C:333:ARG:NE	2.24	0.70
1:E:332:PHE:O	1:E:335:LEU:N	2.26	0.69
1:K:358:ALA:O	1:K:362:ASN:ND2	2.30	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/86 (96%)	81 (98%)	2 (2%)	0	100	100
1	B	83/86 (96%)	80 (96%)	3 (4%)	0	100	100
1	C	83/86 (96%)	79 (95%)	4 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	83/86 (96%)	80 (96%)	3 (4%)	0	100	100
1	E	84/86 (98%)	75 (89%)	9 (11%)	0	100	100
1	F	84/86 (98%)	84 (100%)	0	0	100	100
1	G	84/86 (98%)	83 (99%)	1 (1%)	0	100	100
1	H	83/86 (96%)	78 (94%)	5 (6%)	0	100	100
1	I	84/86 (98%)	79 (94%)	5 (6%)	0	100	100
1	J	84/86 (98%)	81 (96%)	3 (4%)	0	100	100
1	K	84/86 (98%)	82 (98%)	2 (2%)	0	100	100
1	L	84/86 (98%)	82 (98%)	2 (2%)	0	100	100
All	All	1003/1032 (97%)	964 (96%)	39 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/77 (99%)	71 (93%)	5 (7%)	16	19
1	B	76/77 (99%)	68 (90%)	8 (10%)	7	6
1	C	76/77 (99%)	63 (83%)	13 (17%)	2	1
1	D	76/77 (99%)	69 (91%)	7 (9%)	9	9
1	E	77/77 (100%)	72 (94%)	5 (6%)	17	19
1	F	77/77 (100%)	72 (94%)	5 (6%)	17	19
1	G	77/77 (100%)	74 (96%)	3 (4%)	32	41
1	H	76/77 (99%)	71 (93%)	5 (7%)	16	19
1	I	77/77 (100%)	72 (94%)	5 (6%)	17	19
1	J	77/77 (100%)	74 (96%)	3 (4%)	32	41
1	K	77/77 (100%)	75 (97%)	2 (3%)	46	58
1	L	77/77 (100%)	76 (99%)	1 (1%)	69	81

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	919/924 (100%)	857 (93%)	62 (7%)	16	18

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	369	GLN
1	I	409	MET
1	E	363	ASN
1	I	399	LEU
1	K	380	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	342	GLN
1	E	365	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	85/86 (98%)	1.10	10 (11%) 4 4	28, 50, 86, 93	0
1	B	85/86 (98%)	0.95	6 (7%) 16 14	28, 49, 81, 94	0
1	C	85/86 (98%)	1.17	12 (14%) 2 2	28, 49, 87, 93	0
1	D	85/86 (98%)	0.96	7 (8%) 11 10	28, 49, 79, 86	0
1	E	86/86 (100%)	1.14	10 (11%) 4 4	28, 50, 86, 95	0
1	F	86/86 (100%)	0.98	8 (9%) 8 7	29, 50, 83, 95	0
1	G	86/86 (100%)	1.18	13 (15%) 2 2	27, 51, 82, 97	0
1	H	85/86 (98%)	1.03	6 (7%) 16 14	29, 50, 86, 93	0
1	I	86/86 (100%)	1.04	11 (12%) 3 3	29, 49, 84, 93	0
1	J	86/86 (100%)	0.89	9 (10%) 6 5	27, 49, 80, 89	0
1	K	86/86 (100%)	1.04	7 (8%) 12 10	28, 48, 84, 92	0
1	L	86/86 (100%)	1.01	8 (9%) 8 7	27, 48, 81, 93	0
All	All	1027/1032 (99%)	1.04	107 (10%) 6 5	27, 50, 85, 97	0

The worst 5 of 107 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	366	ILE	7.7
1	E	411	LEU	6.7
1	K	326	SER	6.4
1	C	411	LEU	5.6
1	G	366	ILE	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.