

Full wwPDB X-ray Structure Validation Report (i)

Mar 4, 2024 – 08:21 PM JST

PDB ID : 8JQ6

Title : Crystal structure of Lactobacillus rhamnosus L-rhamnose isomerase in complex

with D-allose

Authors : Yoshida, H.; Yoshihara, A.

Deposited on : 2023-06-13

Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Wiedite	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	434	88%	6%	6%
1	В	434	88%	6%	6%
1	С	434	88%	6%	6%
1	D	434	90%	5%	5%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 14417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called L-rhamnose isomerase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	409	Total	С	N	О	S	0	0	0
1	A	409	3322	2128	566	617	11	0		
1	В	408	Total	С	N	О	S	0	0	0
1			3318	2126	565	616	11	U	U	
1	С	409	Total	С	N	О	S	0	0	0
1			3322	2128	566	617	11	0	U	
1	1 D	414	Total	С	N	О	S	0	0	0
1		414	3365	2156	574	624	11	U	U	U

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	В	2	Total Mn 2 2	0	0
2	С	2	Total Mn 2 2	0	0
2	D	2	Total Mn 2 2	0	0

• Molecule 3 is beta-D-allopyranose (three-letter code: ALL) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 12 6 6	0	0
3	A	1	Total C O 12 6 6	0	0
3	В	1	Total C O 12 6 6	0	0
3	D	1	Total C O 12 6 6	0	0
3	D	1	Total C O 12 6 6	0	0
3	D	1	Total C O 12 6 6	0	0

• Molecule 4 is alpha-D-allopyranose (three-letter code: AFD) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	Total C O 12 6 6	0	0
4	D	1	Total C O 12 6 6	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	240	Total O 240 240	0	0
5	В	265	Total O 265 265	0	0
5	С	237	Total O 237 237	0	0
5	D	244	Total O 244 244	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	89.93Å 140.06Å 147.47Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.43 - 1.71	Depositor
rtesolution (A)	46.38 - 1.71	EDS
% Data completeness	100.0 (46.43-1.71)	Depositor
(in resolution range)	100.0 (46.38-1.71)	EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.23 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.8	Depositor
D D.	0.140 , 0.187	Depositor
R, R_{free}	0.153 , 0.196	DCC
R_{free} test set	10079 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	15.7	Xtriage
Anisotropy	1.173	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 39.6	EDS
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14417	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, AFD, ALL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
IVIOI	Chain	RMSZ $ \# Z > 5$		$\mid \text{RMSZ} \mid \# Z > 1$	
1	A	0.61	0/3413	0.66	0/4623
1	В	0.61	0/3409	0.66	0/4618
1	С	0.62	0/3413	0.66	0/4623
1	D	0.61	0/3457	0.67	0/4683
All	All	0.61	0/13692	0.66	0/18547

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3322	0	3218	13	0
1	В	3318	0	3215	17	0
1	С	3322	0	3218	13	0
1	D	3365	0	3264	10	0
2	A	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
3	A	24	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	12	0	0	0	0
3	D	36	0	0	0	0
4	С	12	0	4	0	0
4	D	12	0	4	0	0
5	A	240	0	0	0	0
5	В	265	0	0	0	0
5	С	237	0	0	3	0
5	D	244	0	0	0	0
All	All	14417	0	12923	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
7100111 1	7100111 2	$\operatorname{distance} (\text{\AA})$	overlap (Å)
1:C:370:ASP:HB2	5:C:804:HOH:O	1.50	1.10
1:D:3:LYS:HB2	1:D:6:GLU:HG3	1.57	0.86
1:C:149:VAL:O	1:C:157:ARG:HG3	1.85	0.75
1:B:149:VAL:HG22	1:B:188:LEU:HD13	1.69	0.73
1:B:260:ILE:O	1:B:287:MET:HE3	1.88	0.73
1:B:149:VAL:CG2	1:B:188:LEU:HD13	2.28	0.63
1:B:1:MET:HA	1:B:1:MET:HE2	1.83	0.61
1:D:67:TYR:CG	1:D:68:PRO:HD2	2.35	0.60
1:A:157:ARG:HH11	1:A:157:ARG:HB3	1.68	0.58
1:C:149:VAL:O	1:C:157:ARG:CG	2.53	0.54
1:C:46:ASP:HA	1:C:330:PHE:CE2	2.45	0.52
1:B:51:LEU:HD11	1:B:107:ARG:HB2	1.93	0.50
1:D:149:VAL:HG13	1:D:209:ALA:HB1	1.94	0.50
1:D:67:TYR:CD2	1:D:68:PRO:HD2	2.47	0.49
1:A:279:PHE:N	1:A:280:PRO:CD	2.75	0.49
1:C:279:PHE:N	1:C:280:PRO:CD	2.76	0.49
1:D:289:VAL:HG23	1:D:301:VAL:HG21	1.95	0.49
1:C:227:PHE:HB3	1:C:247:TYR:CD1	2.48	0.49
1:A:149:VAL:HG13	1:A:209:ALA:HB1	1.95	0.48
1:B:230:LYS:HE3	1:B:296:ASP:OD2	2.13	0.48
1:B:146:ASN:O	1:B:190:ASP:HA	2.14	0.48
1:A:3:LYS:HB2	1:A:6:GLU:HG3	1.96	0.47
1:B:227:PHE:HB3	1:B:247:TYR:CD1	2.48	0.47
1:D:46:ASP:HA	1:D:330:PHE:CE1	2.50	0.47
1:D:227:PHE:HB3	1:D:247:TYR:CD1	2.49	0.47

Continued on next page...



Continued from previous page...

A . 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)
1:B:1:MET:HE3	1:B:407:ASN:ND2	2.30	0.46
1:C:311:THR:O	1:C:315:VAL:HG13	2.16	0.46
1:A:227:PHE:HB3	1:A:247:TYR:CD1	2.50	0.46
1:C:151:SER:O	1:C:157:ARG:HD3	2.16	0.46
1:A:201:THR:N	1:A:202:PRO:CD	2.78	0.46
1:A:46:ASP:HA	1:A:330:PHE:CE2	2.50	0.46
1:A:146:ASN:O	1:A:190:ASP:HA	2.17	0.45
1:A:140:HIS:CG	1:A:141:PRO:HD2	2.52	0.45
1:C:4:PRO:HA	1:C:395:GLN:OE1	2.17	0.45
1:B:303:MET:CE	1:B:308:ILE:HD11	2.47	0.45
1:C:363:LYS:NZ	5:C:604:HOH:O	2.48	0.45
1:D:279:PHE:N	1:D:280:PRO:CD	2.80	0.45
1:D:51:LEU:HD12	1:D:51:LEU:HA	1.84	0.44
1:C:146:ASN:O	1:C:190:ASP:HA	2.17	0.44
1:B:30:LYS:HD2	1:B:30:LYS:HA	1.80	0.44
1:B:279:PHE:N	1:B:280:PRO:CD	2.80	0.44
1:C:149:VAL:HG13	1:C:209:ALA:HB1	1.99	0.43
1:A:413:GLU:HA	1:A:417:GLN:HB2	2.01	0.43
1:A:51:LEU:HD23	1:A:51:LEU:HA	1.78	0.43
1:B:1:MET:CE	1:B:407:ASN:ND2	2.82	0.43
1:B:1:MET:CE	1:B:407:ASN:HD22	2.31	0.42
1:A:149:VAL:O	1:A:157:ARG:HG2	2.20	0.42
1:C:324:HIS:HE1	5:C:606:HOH:O	2.02	0.42
1:B:149:VAL:CG2	1:B:188:LEU:CD1	2.95	0.41
1:D:146:ASN:O	1:D:190:ASP:HA	2.20	0.41
1:A:40:HIS:HA	1:A:95:GLN:HB2	2.02	0.41
1:B:46:ASP:HA	1:B:330:PHE:CE2	2.56	0.40
1:B:51:LEU:HD12	1:B:51:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	405/434~(93%)	396 (98%)	9 (2%)	0	100 100
1	В	404/434~(93%)	394 (98%)	10 (2%)	0	100 100
1	C	405/434~(93%)	393 (97%)	12 (3%)	0	100 100
1	D	410/434~(94%)	401 (98%)	9 (2%)	0	100 100
All	All	$1624/1736 \ (94\%)$	1584 (98%)	40 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	358/378 (95%)	354 (99%)	4 (1%)	73 62
1	В	358/378 (95%)	351 (98%)	7 (2%)	55 37
1	С	358/378 (95%)	352 (98%)	6 (2%)	60 44
1	D	363/378 (96%)	357 (98%)	6 (2%)	60 44
All	All	1437/1512 (95%)	1414 (98%)	23 (2%)	62 47

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	51	LEU
1	A	97	HIS
1	A	104	ASP
1	В	1	MET
1	В	97	HIS
1	В	155	LYS
1	В	255	ASN
1	В	286	PHE
1	В	287	MET
1	В	414	LYS
1	С	1	MET
1	С	91	LYS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	С	97	HIS
1	С	224	ILE
1	С	286	PHE
1	С	364	LYS
1	D	2	VAL
1	D	9	LYS
1	D	97	HIS
1	D	126	GLN
1	D	286	PHE
1	D	423	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	ASN
1	A	324	HIS
1	A	407	ASN
1	В	407	ASN
1	С	324	HIS
1	D	48	HIS
1	D	407	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ALL	D	505	-	12,12,12	0.53	0	17,17,17	0.92	0
3	ALL	D	506	-	12,12,12	0.73	0	17,17,17	0.79	0
3	ALL	В	503	2	12,12,12	0.66	0	17,17,17	0.89	0
3	ALL	A	503	2	12,12,12	0.79	0	17,17,17	0.65	0
4	AFD	D	503	2	12,12,12	0.36	0	17,17,17	0.35	0
4	AFD	С	503	2	12,12,12	0.43	0	17,17,17	0.34	0
3	ALL	D	504	-	12,12,12	0.55	0	17,17,17	0.73	0
3	ALL	A	504	-	12,12,12	0.63	0	17,17,17	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ALL	D	505	-	-	0/2/22/22	0/1/1/1
3	ALL	D	506	-	-	0/2/22/22	0/1/1/1
3	ALL	В	503	2	-	0/2/22/22	0/1/1/1
3	ALL	A	503	2	-	2/2/22/22	0/1/1/1
4	AFD	D	503	2	-	0/2/22/22	0/1/1/1
4	AFD	С	503	2	-	2/2/22/22	0/1/1/1
3	ALL	D	504	-	-	0/2/22/22	0/1/1/1
3	ALL	A	504	-	-	1/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	С	503	AFD	C4-C5-C6-O6
3	A	503	ALL	C4-C5-C6-O6
3	A	503	ALL	O5-C5-C6-O6

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	С	503	AFD	O5-C5-C6-O6
3	A	504	ALL	O5-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	409/434 (94%)	-0.20	6 (1%) 73 78	14, 19, 32, 56	1 (0%)
1	В	408/434 (94%)	-0.17	4 (0%) 82 85	13, 18, 29, 53	0
1	С	409/434 (94%)	-0.08	7 (1%) 70 74	14, 20, 33, 56	0
1	D	414/434 (95%)	-0.12	7 (1%) 70 74	13, 18, 33, 57	0
All	All	1640/1736 (94%)	-0.14	24 (1%) 73 78	13, 19, 32, 57	1 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	GLY	4.3
1	В	53	PRO	4.2
1	D	425	VAL	4.1
1	D	426	HIS	3.8
1	С	53	PRO	3.8
1	A	106	LYS	3.7
1	D	53	PRO	3.6
1	D	106	LYS	3.6
1	С	106	LYS	3.4
1	С	1	MET	3.1
1	D	424	LEU	3.0
1	С	5	GLU	2.7
1	В	106	LYS	2.7
1	A	418	PHE	2.6
1	В	418	PHE	2.5
1	A	72	ARG	2.5
1	В	231	LEU	2.5
1	A	330	PHE	2.4
1	A	104	ASP	2.3
1	С	16	GLN	2.3
1	С	104	ASP	2.2

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	65	GLY	2.2
1	D	66	ASN	2.1
1	D	1	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	ALL	D	506	12/12	0.70	0.30	30,33,35,37	0
3	ALL	A	504	12/12	0.84	0.21	30,34,38,40	0
3	ALL	A	503	12/12	0.84	0.18	27,38,43,48	0
4	AFD	С	503	12/12	0.84	0.18	25,34,35,38	0
4	AFD	D	503	12/12	0.86	0.15	26,31,34,37	0
3	ALL	В	503	12/12	0.88	0.14	27,37,42,49	0
3	ALL	D	504	12/12	0.89	0.16	26,33,34,34	0
3	ALL	D	505	12/12	0.90	0.22	40,44,51,51	0
2	MN	С	501	1/1	0.99	0.04	42,42,42,42	0
2	MN	D	501	1/1	0.99	0.04	37,37,37,37	0
2	MN	A	501	1/1	1.00	0.04	40,40,40,40	0
2	MN	С	502	1/1	1.00	0.04	26,26,26,26	0
2	MN	A	502	1/1	1.00	0.04	28,28,28,28	0
2	MN	D	502	1/1	1.00	0.05	25,25,25,25	0
2	MN	В	501	1/1	1.00	0.04	40,40,40,40	0
2	MN	В	502	1/1	1.00	0.05	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


































Electron density around ALL D 505: $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around MN C 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around MN A 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around MN C 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around MN A 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around MN D 502:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around MN B 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

