

Full wwPDB NMR Structure Validation Report (i)

Jun 3, 2023 – 05:48 AM EDT

PDB ID : 2JVD BMRB ID : 15476

Title : Solution NMR structure of the folded N-terminal fragment of UPF0291 protein

ynzC from Bacillus subtilis. Northeast Structural Genomics target SR384-1-46

Authors: Aramini, J.M.; Sharma, S.; Huang, Y.J.; Zhao, L.; Owens, L.A.; Stokes, K.;

Jiang, M.; Xiao, R.; Baran, M.C.; Swapna, G.V.T.; Acton, T.B.; Montelione,

G.T.; Northeast Structural Genomics Consortium (NESG)

Deposited on : 2007-09-18

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &: & v1.2 \\ BMRB \ Restraints \ Analysis &: & v1.2 \\ \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive	
Metric	$(\# \mathrm{Entries})$	$(\# \mathrm{Entries})$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	54	63%	26%	11%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mode				
1	A:4-A:37 (34)	0.34	4	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 8 single-model clusters were found.

Cluster number	Models
1	4, 5, 6, 14, 16
2	9, 13, 15, 19
3	7, 10, 11
Single-model clusters	1; 2; 3; 8; 12; 17; 18; 20



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 786 atoms, of which 406 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called UPF0291 protein ynzC.

Mol	Chain	Residues		P	Atom	S			Trace
1	Λ	10	Total	С	Н	N	О	S	0
1	A	48	786	237	406	69	72	2	U

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	LEU	-	expression tag	UNP O31818
A	48	GLU	-	expression tag	UNP O31818
A	49	HIS	-	expression tag	UNP O31818
A	50	HIS	-	expression tag	UNP O31818
A	51	HIS	_	expression tag	UNP O31818
A	52	HIS	-	expression tag	UNP O31818
A	53	HIS	-	expression tag	UNP O31818
A	54	HIS	-	expression tag	UNP O31818



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: UPF0291 protein ynzC




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: UPF0291 protein ynzC



4.2.2 Score per residue for model 2

• Molecule 1: UPF0291 protein ynzC






4.2.3 Score per residue for model 3

• Molecule 1: UPF0291 protein ynzC


Chain A: 61% . 26% 11%



4.2.4 Score per residue for model 4 (medoid)

• Molecule 1: UPF0291 protein ynzC


Chain A: 63% 26% 11%



4.2.5 Score per residue for model 5

• Molecule 1: UPF0291 protein ynzC

Chain A: 57% 6% 26% 11%



4.2.6 Score per residue for model 6

• Molecule 1: UPF0291 protein ynzC


Chain A: 63% 26% 11%



4.2.7 Score per residue for model 7

• Molecule 1: UPF0291 protein ynzC

Chain A: 59% • 26% 11%





4.2.8 Score per residue for model 8

• Molecule 1: UPF0291 protein ynzC



4.2.9 Score per residue for model 9

• Molecule 1: UPF0291 protein ynzC



4.2.10 Score per residue for model 10

• Molecule 1: UPF0291 protein ynzC



4.2.11 Score per residue for model 11

• Molecule 1: UPF0291 protein ynzC



4.2.12 Score per residue for model 12

• Molecule 1: UPF0291 protein ynzC






4.2.13 Score per residue for model 13

• Molecule 1: UPF0291 protein ynzC

Chain A: 57% 6% 26% 11%



4.2.14 Score per residue for model 14

• Molecule 1: UPF0291 protein ynzC

Chain A: 63% 26% 11%

4.2.15 Score per residue for model 15

• Molecule 1: UPF0291 protein ynzC

Chain A: 63% 26% 11%

4.2.16 Score per residue for model 16

• Molecule 1: UPF0291 protein ynzC

Chain A: 54% 7% · 26% 11%



4.2.17 Score per residue for model 17

• Molecule 1: UPF0291 protein ynzC

Chain A: 54% 9% 26% 11%





4.2.18 Score per residue for model 18

• Molecule 1: UPF0291 protein ynzC



4.2.19 Score per residue for model 19

• Molecule 1: UPF0291 protein ynzC



4.2.20 Score per residue for model 20

• Molecule 1: UPF0291 protein ynzC





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	629
Number of shifts mapped to atoms	629
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

	Mol	Chain	Non-H	H(model)	H(added)	Clashes
	1	A	269	281	281	1±1
ĺ	All	All	5380	5620	5620	19

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom 1	Atom-2	Cleab (Å)	Distance(Å)	Mod	dels
Atom-1	Atom-2	$\operatorname{Clash}(\mathrm{\AA})$	Distance(A)	Worst	Total
1:A:16:LYS:HB3	1:A:22:ILE:HB	0.55	1.78	8	2
1:A:9:ARG:NH1	1:A:13:LEU:HD21	0.52	2.19	1	1
1:A:30:GLN:O	1:A:34:ARG:HG3	0.47	2.09	16	4
1:A:31:GLN:O	1:A:35:GLN:HG2	0.44	2.12	10	2
1:A:27:LYS:O	1:A:31:GLN:HG2	0.44	2.13	13	1
1:A:31:GLN:HG2	1:A:35:GLN:OE1	0.43	2.13	5	1
1:A:29:GLU:O	1:A:33:LEU:HG	0.43	2.13	17	1
1:A:18:LYS:NZ	1:A:18:LYS:HB3	0.43	2.29	11	1
1:A:9:ARG:O	1:A:13:LEU:HG	0.42	2.15	12	2
1:A:16:LYS:NZ	1:A:21:VAL:O	0.41	2.53	16	1
1:A:6:LYS:O	1:A:10:ILE:HG13	0.41	2.16	7	2
1:A:23:THR:O	1:A:27:LYS:HG3	0.40	2.17	19	1



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Allowed	Outliers	Percentiles		
1	A	34/54 (63%)	34±1 (99±2%)	0±1 (1±2%)	0±0 (0±1%)	54	85	
All	All	680/1080 (63%)	671 (99%)	8 (1%)	1 (0%)	54	85	

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	21	VAL	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	26/45~(58%)	26±0 (99±1%)	0±0 (1±1%)	86 97
All	All	520/900 (58%)	517 (99%)	3 (1%)	86 97

All 3 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	GLU	1
1	A	32	LYS	1
1	A	18	LYS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.



6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 92% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	629
Number of shifts mapped to atoms	629
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	48	-0.31 ± 0.07	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	46	0.16 ± 0.10	None needed (< 0.5 ppm)
¹³ C′	47	-0.63 ± 0.14	Should be applied
^{15}N	46	-0.82 ± 0.24	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 451 atoms were assigned a chemical shift out of a possible 479. 0 out of 3 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	171/171 (100%)	69/69 (100%)	68/68 (100%)	34/34 (100%)
Sidechain	272/299 (91%)	182/191 (95%)	84/92 (91%)	6/16 (38%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	8/9 (89%)	4/4 (100%)	4/5~(80%)	0/0 (%)
Overall	451/479 (94%)	255/264 (97%)	156/165~(95%)	40/50 (80%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 92%, i.e. 629 atoms were assigned a chemical shift out of a possible 682. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	$237/242 \ (98\%)$	96/98 (98%)	95/96~(99%)	46/48 (96%)
Sidechain	374/421 (89%)	251/271 (93%)	117/129 (91%)	6/21 (29%)
Aromatic	18/19 (95%)	9/9 (100%)	9/10 (90%)	0/0 (%)
Overall	629/682 (92%)	356/378 (94%)	221/235 (94%)	52/69 (75%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1099
Intra-residue ($ i-j =0$)	332
Sequential ($ i-j =1$)	209
Medium range ($ i-j >1$ and $ i-j <5$)	318
Long range (i-j ≥5)	192
Inter-chain	0
Hydrogen bond restraints	48
Disulfide bond restraints	0
Total dihedral-angle restraints	68
Number of unmapped restraints	0
Number of restraints per residue	21.6
Number of long range restraints per residue ¹	3.6

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.3	0.14
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	$\mathbf{Max} \ (^{\circ})$
1.0-10.0 (Small)	0.1	1.2
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dantuninta tema	C	% ¹	\mathbf{V}_{1} Violated ³			Consis	tentl	$\overline{ m y~Violated^4}$
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	332	30.2	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	242	22.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	90	8.2	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	209	19.0	2	1.0	0.2	0	0.0	0.0
Backbone-Backbone	41	3.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	145	13.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	23	2.1	2	8.7	0.2	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	318	28.9	1	0.3	0.1	0	0.0	0.0
Backbone-Backbone	70	6.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	129	11.7	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	119	10.8	1	0.8	0.1	0	0.0	0.0
Long range ($ i-j \ge 5$)	192	17.5	3	1.6	0.3	0	0.0	0.0
Backbone-Backbone	3	0.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	57	5.2	1	1.8	0.1	0	0.0	0.0
Sidechain-Sidechain	132	12.0	2	1.5	0.2	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	48	4.4	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1099	100.0	6	0.5	0.5	0	0.0	0.0
Backbone-Backbone	162	14.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	573	52.1	1	0.2	0.1	0	0.0	0.0
Sidechain-Sidechain	364	33.1	5	1.4	0.5	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	tions	3	Maan (Å)	Max (Å)	${ m SD}^6$ (Å)	Modian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (A)	$SD^*(A)$	Median (Å)
1	0	1	0	0	0	1	0.11	0.11	0.0	0.11
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	1	0	1	0.12	0.12	0.0	0.12
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	1	0	0	1	0.12	0.12	0.0	0.12
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	1	0	1	0.14	0.14	0.0	0.14
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0
11	0	1	0	0	0	1	0.13	0.13	0.0	0.13

Continued on next page...




Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	0	1	0	0	0	1	0.11	0.11	0.0	0.11
13	0	0	0	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	1	0	1	0.11	0.11	0.0	0.11
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0	0	0	0.0	0.0	0.0	0.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1045(IR:332, SQ:207, MR:317, LR:189, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	1	1	3	0	5	1	5.0
0	1	0	0	0	1	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	${f Models}^1$	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,312)	1:A:23:THR:HG21	1:A:24:GLU:HG3	2	0.12	0.01	0.12
(1,312)	1:A:23:THR:HG22	1:A:24:GLU:HG3	2	0.12	0.01	0.12
(1,312)	1:A:23:THR:HG23	1:A:24:GLU:HG3	2	0.12	0.01	0.12


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,883)	1:A:6:LYS:HE2	1:A:36:GLU:HG2	9	0.14
(1,883)	1:A:6:LYS:HE2	1:A:36:GLU:HG3	9	0.14
(1,883)	1:A:6:LYS:HE3	1:A:36:GLU:HG2	9	0.14
(1,883)	1:A:6:LYS:HE3	1:A:36:GLU:HG3	9	0.14
(1,312)	1:A:23:THR:HG21	1:A:24:GLU:HG3	11	0.13
(1,312)	1:A:23:THR:HG22	1:A:24:GLU:HG3	11	0.13
(1,312)	1:A:23:THR:HG23	1:A:24:GLU:HG3	11	0.13
(1,866)	1:A:2:ILE:HD11	1:A:6:LYS:HG2	7	0.12
(1,866)	1:A:2:ILE:HD11	1:A:6:LYS:HG3	7	0.12
(1,866)	1:A:2:ILE:HD12	1:A:6:LYS:HG2	7	0.12
(1,866)	1:A:2:ILE:HD12	1:A:6:LYS:HG3	7	0.12
(1,866)	1:A:2:ILE:HD13	1:A:6:LYS:HG2	7	0.12
(1,866)	1:A:2:ILE:HD13	1:A:6:LYS:HG3	7	0.12
(1,806)	1:A:14:ALA:HA	1:A:30:GLN:HE21	4	0.12
(1,390)	1:A:16:LYS:HE3	1:A:21:VAL:HG21	17	0.11
(1,390)	1:A:16:LYS:HE3	1:A:21:VAL:HG22	17	0.11

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,390)	1:A:16:LYS:HE3	1:A:21:VAL:HG23	17	0.11
(1,312)	1:A:23:THR:HG21	1:A:24:GLU:HG3	1	0.11
(1,312)	1:A:23:THR:HG22	1:A:24:GLU:HG3	1	0.11
(1,312)	1:A:23:THR:HG23	1:A:24:GLU:HG3	1	0.11
(1,311)	1:A:23:THR:HG21	1:A:24:GLU:HG2	12	0.11
(1,311)	1:A:23:THR:HG22	1:A:24:GLU:HG2	12	0.11
(1,311)	1:A:23:THR:HG23	1:A:24:GLU:HG2	12	0.11



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	Count	$\%^{1}$	Vio	lated	3	Consis	tent	${ m ly~Violated^4}$
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PSI	34	50.0	1	2.9	1.5	0	0.0	0.0
PHI	34	50.0	1	2.9	1.5	0	0.0	0.0
Total	68	100.0	2	2.9	2.9	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	f violations	Moon (°)	Mov (°)	SD (°)	Modian (°)
Wiodei 1D	PSI	PHI	Total	$Mean (^{\circ})$	$\mathbf{Max} \ (^{\circ})$	\mathbf{SD} (°)	\mid Median (°) \mid
1	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0.0	0.0	0.0	0.0
11	0	0	0	0.0	0.0	0.0	0.0
12	0	0	0	0.0	0.0	0.0	0.0
13	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0.0	0.0	0.0	0.0
17	1	0	1	1.2	1.2	0.0	1.2
18	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0.0	0.0	0.0	0.0
20	0	1	1	1.2	1.2	0.0	1.2



10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fraction of the ensemble			
PSI	PHI	Total	Count ¹	%		
1	1	2	1	5.0		
0	0	0	2	10.0		
0	0	0	3	15.0		
0	0	0	4	20.0		
0	0	0	5	25.0		
0	0	0	6	30.0		
0	0	0	7	35.0		
0	0	0	8	40.0		
0	0	0	9	45.0		
0	0	0	10	50.0		
0	0	0	11	55.0		

Continued on next page...




Continued from previous page...

Nun	nber o	of violated restraints	Fraction of the ensemble				
PSI	PHI	Total	Count ¹	%			
0	0	0	12	60.0			
0	0	0	13	65.0			
0	0	0	14	70.0			
0	0	0	15	75.0			
0	0	0	16	80.0			
0	0	0	17	85.0			
0	0	0	18	90.0			
0	0	0	19	95.0			
0	0	0	20	100.0			

¹ Number of models with violations

10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)



10.4 Most violated dihedral-angle restraints in the ensemble (i)

No violations found



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,68)	1:A:40:GLY:N	1:A:40:GLY:CA	1:A:40:GLY:C	1:A:41:PHE:N	17	1.2
(1,67)	1:A:39:LYS:C	1:A:40:GLY:N	1:A:40:GLY:CA	1:A:40:GLY:C	20	1.2

