



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2024 – 03:36 PM JST

PDB ID : 6JYS
Title : GII.13/21 noroviruses recognize glycans with a terminal beta-galactose via an unconventional glycan binding site
Authors : Duan, Z.; Xin, C.
Deposited on : 2019-04-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	
1	C	308	
1	D	308	
2	E	2	
2	G	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	2	 100%
3	F	3	 67%  33%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 11371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called human norovirus P protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	308	Total 2406	C 1526	N 410	O 463	S 7	0	3	0
1	B	308	Total 2406	C 1526	N 410	O 463	S 7	0	3	0
1	C	308	Total 2403	C 1524	N 410	O 462	S 7	0	2	0
1	D	308	Total 2400	C 1523	N 409	O 461	S 7	0	2	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-alpha-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	2	Total 25	C 14	N 1	O 10	0	0	0
2	G	2	Total 25	C 14	N 1	O 10	0	0	0
2	H	2	Total 25	C 14	N 1	O 10	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	3	39	22	2	15	0	0	0

- Molecule 4 is THREONINE (three-letter code: THR) (formula: C₄H₉NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	B	1	8	4	1	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	418	Total	O	0	0
			418	418		
5	B	419	Total	O	0	0
			419	419		
5	C	386	Total	O	0	0
			386	386		
5	D	411	Total	O	0	0
			411	411		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: human norovirus P protein




- Molecule 1: human norovirus P protein



- Molecule 1: human norovirus P protein



- Molecule 1: human norovirus P protein



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-alpha-D-galactopyranose



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-alpha-D-galactopyranose

Chain G:  100%

A2G1
GAL2

- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-alpha-D-galactopyranose

Chain H:  100%

A2G1
GAL2

- Molecule 3: beta-D-galactopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose

Chain F:  67% 33%

A2G1
GAL2
NAG3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.73Å 99.69Å 221.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.63 – 1.70 48.63 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.1 (48.63-1.70) 96.1 (48.63-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 1.70Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.165 , 0.188 0.165 , 0.188	Depositor DCC
R_{free} test set	7153 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtrriage
Anisotropy	0.110	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11371	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, A2G, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2478	0.47	0/3387
1	B	0.25	0/2478	0.47	0/3387
1	C	0.26	0/2472	0.47	0/3379
1	D	0.25	0/2472	0.46	0/3379
All	All	0.25	0/9900	0.47	0/13532

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	33	ASN	Peptide
1	B	33	ASN	Peptide
1	C	33	ASN	Peptide
1	D	33	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2406	0	2323	12	0
1	B	2406	0	2323	5	0
1	C	2403	0	2318	7	0
1	D	2400	0	2319	10	0
2	E	25	0	21	0	0
2	G	25	0	21	0	0
2	H	25	0	21	0	0
3	F	39	0	33	0	0
4	B	8	0	5	1	0
5	A	418	0	0	5	0
5	B	419	0	0	0	0
5	C	386	0	0	5	0
5	D	411	0	0	1	0
All	All	11371	0	9384	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:SER:HG	4:B:404:THR:N	1.76	0.83
1:C:130:LYS:NZ	5:C:601:HOH:O	1.99	0.75
1:C:156:ASP:OD2	5:C:602:HOH:O	2.14	0.66
1:D:26:GLU:OE1	1:D:215:ARG:NH1	2.29	0.62
1:B:66:ARG:HD2	1:B:86:PRO:HA	1.84	0.60
1:A:261:ASP:OD2	5:A:501:HOH:O	2.17	0.59
1:A:66:ARG:NH2	5:A:505:HOH:O	2.35	0.58
1:A:215:ARG:NH1	5:A:507:HOH:O	2.36	0.57
1:C:256:ARG:NH1	5:C:610:HOH:O	2.37	0.57
1:D:302:ASN:ND2	1:D:304:PHE:H	2.02	0.57
1:D:3:THR:N	5:D:503:HOH:O	2.37	0.57
1:A:68:ARG:NH1	5:A:509:HOH:O	2.38	0.57
1:A:219:VAL:HA	1:D:128:ASN:HB3	1.86	0.56
1:C:127:LYS:NZ	5:C:614:HOH:O	2.40	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ASN:HB3	1:D:219:VAL:HA	1.88	0.54
1:C:3:THR:N	5:C:613:HOH:O	2.40	0.54
1:D:252:VAL:HG21	1:D:303:GLN:HA	1.93	0.51
1:B:256:ARG:NH1	1:B:298:ASP:OD1	2.32	0.50
1:D:170:ILE:HD13	1:D:178:PHE:CG	2.47	0.50
1:A:256:ARG:NH1	1:A:298:ASP:OD1	2.34	0.49
1:C:92:ASP:OD2	1:C:94:THR:OG1	2.21	0.47
1:A:270:LYS:HG3	1:A:307:LEU:HD11	1.97	0.47
1:C:262:THR:O	1:C:264:ARG:N	2.48	0.46
1:A:164:ARG:NH2	5:A:519:HOH:O	2.49	0.46
1:D:215:ARG:HG2	1:D:226:ASP:HB3	2.00	0.44
1:A:296:ARG:HE	1:A:296:ARG:HB3	1.61	0.44
1:D:66:ARG:HD2	1:D:86:PRO:HA	2.00	0.43
1:B:87:ASN:OD1	1:B:88:GLY:N	2.51	0.43
1:D:47:LEU:HG	1:D:273:ARG:HA	2.02	0.42
1:A:170:ILE:HD13	1:A:178:PHE:CG	2.56	0.41
1:A:26:GLU:OE1	1:A:215:ARG:NE	2.54	0.40
1:B:38:VAL:HG13	1:B:181:TRP:CZ3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/308 (100%)	301 (97%)	8 (3%)	0	100	100
1	B	309/308 (100%)	299 (97%)	10 (3%)	0	100	100
1	C	308/308 (100%)	299 (97%)	9 (3%)	0	100	100
1	D	308/308 (100%)	299 (97%)	9 (3%)	0	100	100
All	All	1234/1232 (100%)	1198 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/266 (101%)	267 (99%)	2 (1%)	84	77
1	B	269/266 (101%)	269 (100%)	0	100	100
1	C	268/266 (101%)	268 (100%)	0	100	100
1	D	268/266 (101%)	266 (99%)	2 (1%)	84	77
All	All	1074/1064 (101%)	1070 (100%)	4 (0%)	91	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	LEU
1	A	303	GLN
1	D	68	ARG
1	D	273	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	36	ASN
1	D	302	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A2G	E	1	2	14,14,15	0.49	0	17,19,21	0.95	0
2	GAL	E	2	2	11,11,12	0.54	0	15,15,17	1.03	2 (13%)
3	A2G	F	1	4,3	14,14,15	0.48	0	17,19,21	0.85	1 (5%)
3	GAL	F	2	3	11,11,12	0.58	0	15,15,17	0.91	0
3	NAG	F	3	3	14,14,15	0.28	0	17,19,21	0.38	0
2	A2G	G	1	2	14,14,15	0.53	0	17,19,21	0.73	0
2	GAL	G	2	2	11,11,12	0.44	0	15,15,17	0.94	0
2	A2G	H	1	2	14,14,15	0.54	0	17,19,21	0.71	0
2	GAL	H	2	2	11,11,12	0.56	0	15,15,17	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A2G	E	1	2	-	2/6/23/26	0/1/1/1
2	GAL	E	2	2	-	0/2/19/22	0/1/1/1
3	A2G	F	1	4,3	-	0/6/23/26	0/1/1/1
3	GAL	F	2	3	-	0/2/19/22	0/1/1/1
3	NAG	F	3	3	-	0/6/23/26	0/1/1/1
2	A2G	G	1	2	-	2/6/23/26	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	A2G	H	1	2	-	2/6/23/26	0/1/1/1
2	GAL	H	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	E	2	GAL	C1-O5-C5	2.12	115.06	112.19
2	E	2	GAL	C1-C2-C3	2.02	112.15	109.67
3	F	1	A2G	C1-O5-C5	2.01	114.92	112.19

There are no chirality outliers.


All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	1	A2G	C8-C7-N2-C2
2	H	1	A2G	C8-C7-N2-C2
2	H	1	A2G	O7-C7-N2-C2
2	E	1	A2G	O7-C7-N2-C2
2	G	1	A2G	O7-C7-N2-C2
2	G	1	A2G	C8-C7-N2-C2


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	THR	B	404	3	6,7,7	0.92	1 (16%)	7,9,9	1.27	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	THR	B	404	3	-	0/8/8/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	404	THR	OXT-C	-2.11	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	THR	OXT-C-O	-2.62	118.13	124.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	404	THR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/308 (100%)	0.05	10 (3%) 47 52	7, 14, 30, 60	0
1	B	308/308 (100%)	0.01	7 (2%) 60 65	7, 15, 35, 74	0
1	C	308/308 (100%)	0.05	8 (2%) 56 60	7, 17, 35, 72	0
1	D	308/308 (100%)	0.02	11 (3%) 42 47	7, 15, 32, 74	0
All	All	1232/1232 (100%)	0.03	36 (2%) 51 56	7, 15, 33, 74	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	89	ALA	5.9
1	B	3	THR	4.8
1	D	3	THR	4.5
1	C	264	ARG	4.5
1	A	304	PHE	4.5
1	A	85	TYR	4.4
1	D	91	TYR	3.7
1	D	89	ALA	3.7
1	A	86	PRO	3.6
1	A	3	THR	3.5
1	B	85	TYR	3.5
1	D	85	TYR	3.4
1	A	89	ALA	3.4
1	C	3	THR	3.4
1	D	304	PHE	3.2
1	C	262	THR	3.1
1	A	303	GLN	3.1
1	D	303	GLN	2.8
1	B	91	TYR	2.8
1	B	87	ASN	2.8
1	C	263	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	88	GLY	2.7
1	D	87	ASN	2.5
1	A	190	LEU	2.5
1	B	304	PHE	2.5
1	C	304	PHE	2.4
1	A	34	GLU	2.3
1	C	70	LEU	2.2
1	C	87	ASN	2.2
1	D	86	PRO	2.2
1	A	87	ASN	2.2
1	C	34	GLU	2.2
1	D	10	ILE	2.2
1	D	73	ASN	2.1
1	B	86	PRO	2.1
1	A	84	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A2G	G	1	14/15	0.71	0.20	26,35,46,57	0
2	A2G	H	1	14/15	0.72	0.22	31,38,50,54	0
2	A2G	E	1	14/15	0.82	0.14	19,28,37,41	0
3	NAG	F	3	14/15	0.87	0.15	20,27,33,43	0
2	GAL	H	2	11/12	0.91	0.12	14,20,27,34	0
3	A2G	F	1	14/15	0.93	0.10	11,15,22,22	0
2	GAL	G	2	11/12	0.94	0.10	15,19,26,28	0
3	GAL	F	2	11/12	0.95	0.07	8,12,16,16	0
2	GAL	E	2	11/12	0.95	0.09	11,14,20,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	THR	B	404	8/8	0.84	0.17	18,25,32,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.