# Full wwPDB NMR Structure Validation Report (i)

Mar 5, 2022 – 10:53 AM EST

| | | |
|---|---|---|
| PDB ID | : | 2K1W |
| Title | : | NMR solution structure of M-crystallin in calcium loaded form(holo). |
| Authors | : | Barnwal, R.; Jobby, M.; Devi, K.; Sharma, Y.; Chary, K. |
| Deposited on | : | 2008-03-17 |

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4.02b-467 |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| RCI | : | v_1n_11_5_13_A (Berjanski et al., 2005) |
| PANAV | : | Wang et al. (2010) |
| ShiftChecker | : | 2.27 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.27 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|---|---|---|
| Clashscore | 158937 | 12864 |
| Ramachandran outliers | 154571 | 11451 |
| Sidechain outliers | 154315 | 11428 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 85 | 39% 35% • • 21% |

# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

| Well-defined (core) protein residues | | | |
|---|---|---|---|
| **Well-defined core** | **Residue range (total)** | **Backbone RMSD (Å)** | **Medoid model** |
| 1 | A:5-A:19, A:34-A:85 (67) | 0.28 | 1 |

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

| Cluster number | Models |
|---|---|
| 1 | 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 13, 14, 16 |
| 2 | 15, 17, 18, 19, 20 |
| Single-model clusters | 10; 12 |

# 3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1277 atoms, of which 612 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Beta/gama crystallin family protein.

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | H | N | O | S | |
| 1 | A | 85 | 1275 | 418 | 612 | 110 | 134 | 1 | 0 |

There is a discrepancy between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | 1 | MET | - | expression tag | UNP Q8TMX3 |

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | |
|---|---|---|---|---|
| | | | Total | Ca |
| 2 | A | 2 | 2 | 2 |

# 4 Residue-property plots (i)

## 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Beta/gama crystallin family protein

Chain A: 39% 35% • • 21%

M1 N2 A3 A4 E5 V6 H11 V12 N13 F14 G15 G16 K17 S18 F19 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 I39 K40 V41 K42 W46 R47 Y51 I52 N53 Y54 G55 L61 G62 Y66 V69 E70 G73 I74 P75

D76 N77 S81 Q84 I85

## 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

### 4.2.1 Score per residue for model 1 (medoid)

• Molecule 1: Beta/gama crystallin family protein

Chain A: 35% 38% 5% • 21%

M1 N2 A3 A4 E5 V6 I7 V8 H11 V12 N13 F14 G15 G16 K17 S18 F19 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 V41 K42 S43 W46 R47 Y51 I52 N53 Y54 G55 G56 R57 L61 G62 E65 Y66 V69

E70 G73 I74 P75 D76 N77 S81 F82 R83 Q84 I85

### 4.2.2 Score per residue for model 2

• Molecule 1: Beta/gama crystallin family protein

Chain A: 41% 32% 5% • 21%




### 4.2.3 Score per residue for model 3

- Molecule 1: Beta/gama crystallin family protein

Chain A: 42% 28% 7% • 21%




### 4.2.4 Score per residue for model 4

- Molecule 1: Beta/gama crystallin family protein

Chain A: 45% 27% 6% • 21%




### 4.2.5 Score per residue for model 5

- Molecule 1: Beta/gama crystallin family protein

Chain A: 48% 25% 5% • 21%




### 4.2.6 Score per residue for model 6

- Molecule 1: Beta/gama crystallin family protein

Chain A: 41% 29% 7% • 21%




### 4.2.7 Score per residue for model 7

- Molecule 1: Beta/gama crystallin family protein

Chain A: 40% 33% 5% • 21%




### 4.2.8 Score per residue for model 8

- Molecule 1: Beta/gama crystallin family protein

Chain A: 45% 28% 5% • 21%

### 4.2.9 Score per residue for model 9

- Molecule 1: Beta/gama crystallin family protein

Chain A: 41% 32% 5% • 21%




### 4.2.10 Score per residue for model 10

- Molecule 1: Beta/gama crystallin family protein

Chain A: 40% 32% 6% • 21%

M1 N2 A3 A4 E5 V6 H11 V12 N13 F14 G15 G16 K17 S18 F19 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 I39 K40 V41 W46 R47 E50 Y51 Y54 R57 L61 V69 E70 G73 I74 P75 D76 N77

S80 S81 F82 R83 Q84 I85

### 4.2.11 Score per residue for model 11

- Molecule 1: Beta/gama crystallin family protein

Chain A: 41% 31% 6% • 21%

M1 N2 A3 A4 E5 V6 E10 H11 V12 N13 F14 K17 S18 F19 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 I39 K40 V41 K42 S43 W46 Y51 G55 G56 R57 L61 G62 P63 V69 E70 G73 I74 P75 D76

N77 S81 Q84 I85

### 4.2.12 Score per residue for model 12

- Molecule 1: Beta/gama crystallin family protein

Chain A: 32% 39% 7% • 21%

M1 N2 A3 A4 E5 V6 I7 V8 H11 V12 N13 F14 G15 G16 K17 S18 F19 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 I39 K40 V41 W46 R47 Y51 I52 N53 Y54 G55 G56 R57 L61 G62 E65 Y66 V69 E70

S71 A72 G73 I74 P75 S78 I79 S80 S81 F82 R83 Q84 I85

### 4.2.13 Score per residue for model 13

- Molecule 1: Beta/gama crystallin family protein

Chain A: 40% 34% • • 21%

M1 N2 A3 A4 E5 V6 I7 V8 H11 V12 G15 G16 K17 D20 A21 T22 S23 D24 Q25 P26 G27 A28 G29 D30 N31 L32 N33 D34 K35 I36 S37 S38 I39 K40 V41 W46 R47 Y51 I52 N53 Y54 G55 L61 G62 E65 Y66 V69 E70 G73 I74 P75

D76 N77 S80 S81 F82 R83 Q84 I85

### 4.2.14 Score per residue for model 14

- Molecule 1: Beta/gama crystallin family protein

Chain A: 41% 28% 8% • 21%




### 4.2.15 Score per residue for model 15

- Molecule 1: Beta/gama crystallin family protein

Chain A: 44% 32% • • 21%




### 4.2.16 Score per residue for model 16

- Molecule 1: Beta/gama crystallin family protein

Chain A: 38% 34% 6% • 21%




### 4.2.17 Score per residue for model 17

- Molecule 1: Beta/gama crystallin family protein

Chain A: 36% 33% 8% • 21%



### 4.2.18 Score per residue for model 18

- Molecule 1: Beta/gama crystallin family protein

Chain A: 33% 39% 6% • 21%




### 4.2.19 Score per residue for model 19

- Molecule 1: Beta/gama crystallin family protein

Chain A: 39% 29% 9% • 21%




### 4.2.20 Score per residue for model 20

- Molecule 1: Beta/gama crystallin family protein

Chain A: 34% 38% 6% • 21%



# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics, simulated annealing.*

Of the 400 calculated structures, 20 were deposited, based on the following criterion: *target function.*

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---|---|---|
| CYANA | structure solution | 3.0 |
| CYANA | refinement | 3.0 |

No chemical shift data was provided.

# 6 Model quality (i)

## 6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|---|---|---|---|---|---|
| | | RMSZ | #Z>5 | RMSZ | #Z>5 |
| 1 | A | 0.43±0.01 | 1±0/558 ( 0.2± 0.0%) | 0.49±0.01 | 1±0/753 ( 0.1± 0.0%) |
| All | All | 0.43 | 20/11160 ( 0.2%) | 0.49 | 20/15060 ( 0.1%) |

All unique bond outliers are listed below.

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) | Models | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Worst | Total |
| 1 | A | 75 | PRO | CG-CD | 9.36 | 1.81 | 1.50 | 16 | 20 |

All unique angle outliers are listed below.

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) | Models | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Worst | Total |
| 1 | A | 75 | PRO | N-CD-CG | -10.64 | 87.23 | 103.20 | 2 | 20 |

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|---|---|---|---|---|---|
| 1 | A | 542 | 509 | 508 | 13±2 |
| All | All | 10880 | 10180 | 10170 | 256 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|---|---|---|---|---|---|
| | | | | Worst | Total |
| 1:A:75:PRO:CD | 1:A:75:PRO:CG | 1.55 | 1.79 | 3 | 20 |
| 1:A:34:ASP:O | 1:A:74:ILE:HG23 | 0.65 | 1.91 | 11 | 17 |
| 1:A:66:TYR:OH | 1:A:72:ALA:HB2 | 0.65 | 1.92 | 2 | 2 |
| 1:A:34:ASP:OD2 | 1:A:75:PRO:HD2 | 0.62 | 1.93 | 11 | 6 |
| 1:A:73:GLY:C | 1:A:75:PRO:HD3 | 0.61 | 2.16 | 2 | 14 |
| 1:A:36:ILE:HD12 | 1:A:69:VAL:HB | 0.61 | 1.72 | 3 | 11 |
| 1:A:52:ILE:HD12 | 1:A:54:TYR:CE1 | 0.60 | 2.31 | 8 | 4 |
| 1:A:6:VAL:HG22 | 1:A:41:VAL:HG11 | 0.60 | 1.74 | 8 | 19 |
| 1:A:38:SER:C | 1:A:39:ILE:HD12 | 0.60 | 2.17 | 12 | 9 |
| 1:A:73:GLY:O | 1:A:75:PRO:HD3 | 0.59 | 1.98 | 11 | 13 |
| 1:A:6:VAL:HG11 | 1:A:84:GLN:OE1 | 0.59 | 1.96 | 19 | 8 |
| 1:A:6:VAL:HG22 | 1:A:41:VAL:CG1 | 0.57 | 2.29 | 4 | 2 |
| 1:A:36:ILE:HB | 1:A:69:VAL:HG23 | 0.57 | 1.76 | 9 | 13 |
| 1:A:73:GLY:O | 1:A:74:ILE:HD13 | 0.56 | 2.00 | 5 | 2 |
| 1:A:34:ASP:HA | 1:A:74:ILE:HG22 | 0.55 | 1.78 | 10 | 1 |
| 1:A:6:VAL:HG21 | 1:A:84:GLN:OE1 | 0.55 | 2.01 | 6 | 4 |
| 1:A:12:VAL:HG23 | 1:A:36:ILE:HD13 | 0.55 | 1.78 | 13 | 1 |
| 1:A:61:LEU:HD12 | 1:A:62:GLY:O | 0.54 | 2.02 | 11 | 16 |
| 1:A:6:VAL:HG21 | 1:A:84:GLN:NE2 | 0.54 | 2.18 | 12 | 2 |
| 1:A:52:ILE:HD12 | 1:A:54:TYR:CD1 | 0.54 | 2.38 | 1 | 5 |
| 1:A:8:VAL:HG21 | 1:A:82:PHE:CZ | 0.53 | 2.39 | 17 | 6 |
| 1:A:12:VAL:HG21 | 1:A:14:PHE:CZ | 0.52 | 2.39 | 2 | 3 |
| 1:A:84:GLN:O | 1:A:85:ILE:HD13 | 0.52 | 2.03 | 18 | 3 |
| 1:A:12:VAL:HG23 | 1:A:36:ILE:HG21 | 0.52 | 1.80 | 17 | 2 |
| 1:A:6:VAL:HG13 | 1:A:41:VAL:HG21 | 0.51 | 1.80 | 4 | 1 |
| 1:A:6:VAL:HG11 | 1:A:84:GLN:CD | 0.51 | 2.27 | 7 | 4 |
| 1:A:36:ILE:HD12 | 1:A:69:VAL:HG12 | 0.50 | 1.83 | 13 | 1 |
| 1:A:6:VAL:HG21 | 1:A:84:GLN:HE21 | 0.49 | 1.67 | 12 | 4 |
| 1:A:12:VAL:HG23 | 1:A:36:ILE:CG2 | 0.48 | 2.38 | 11 | 3 |
| 1:A:12:VAL:HG13 | 1:A:14:PHE:H | 0.48 | 1.68 | 18 | 5 |
| 1:A:46:TRP:CZ3 | 1:A:61:LEU:HD11 | 0.48 | 2.43 | 6 | 4 |
| 1:A:12:VAL:HG13 | 1:A:14:PHE:N | 0.48 | 2.24 | 14 | 8 |
| 1:A:6:VAL:HG11 | 1:A:84:GLN:NE2 | 0.47 | 2.24 | 2 | 2 |
| 1:A:83:ARG:HG2 | 1:A:85:ILE:HD11 | 0.46 | 1.86 | 20 | 1 |
| 1:A:59:TRP:CZ2 | 1:A:61:LEU:HD22 | 0.46 | 2.45 | 9 | 1 |
| 1:A:36:ILE:HD12 | 1:A:69:VAL:CG1 | 0.45 | 2.41 | 13 | 1 |
| 1:A:8:VAL:HG11 | 1:A:82:PHE:CZ | 0.45 | 2.47 | 18 | 1 |
| 1:A:47:ARG:NH1 | 1:A:85:ILE:HD11 | 0.44 | 2.27 | 16 | 1 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|---|---|---|---|---|---|
| | | | | Worst | Total |
| 1:A:12:VAL:HG21 | 1:A:14:PHE:CE1 | 0.44 | 2.48 | 1 | 4 |
| 1:A:8:VAL:HG11 | 1:A:82:PHE:CE2 | 0.44 | 2.48 | 18 | 1 |
| 1:A:52:ILE:HD12 | 1:A:54:TYR:CZ | 0.44 | 2.48 | 8 | 1 |
| 1:A:12:VAL:HG11 | 1:A:14:PHE:CE1 | 0.43 | 2.48 | 11 | 1 |
| 1:A:39:ILE:O | 1:A:39:ILE:HG22 | 0.43 | 2.12 | 11 | 4 |
| 1:A:69:VAL:O | 1:A:69:VAL:HG22 | 0.43 | 2.14 | 4 | 8 |
| 1:A:35:LYS:O | 1:A:36:ILE:C | 0.43 | 2.57 | 12 | 2 |
| 1:A:6:VAL:HA | 1:A:41:VAL:HG13 | 0.43 | 1.90 | 10 | 1 |
| 1:A:35:LYS:HE2 | 1:A:74:ILE:HG21 | 0.42 | 1.91 | 19 | 1 |
| 1:A:69:VAL:HG22 | 1:A:69:VAL:O | 0.41 | 2.15 | 12 | 4 |
| 1:A:52:ILE:HD12 | 1:A:54:TYR:HD1 | 0.41 | 1.75 | 3 | 2 |
| 1:A:6:VAL:HG13 | 1:A:41:VAL:CG2 | 0.41 | 2.46 | 4 | 1 |
| 1:A:7:ILE:HD11 | 1:A:18:SER:HB2 | 0.41 | 1.91 | 20 | 1 |
| 1:A:74:ILE:N | 1:A:75:PRO:HD3 | 0.41 | 2.31 | 18 | 1 |
| 1:A:12:VAL:HG11 | 1:A:14:PHE:CD1 | 0.41 | 2.51 | 11 | 1 |
| 1:A:52:ILE:HG23 | 1:A:55:GLY:N | 0.41 | 2.30 | 19 | 1 |
| 1:A:38:SER:O | 1:A:39:ILE:HD12 | 0.40 | 2.17 | 7 | 1 |
| 1:A:52:ILE:HD11 | 1:A:55:GLY:HA3 | 0.40 | 1.93 | 17 | 1 |

## 6.3 Torsion angles (i)

### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|---|---|---|---|---|---|---|---|
| 1 | A | 66/85 (78%) | 47±2 (72±3%) | 13±2 (19±3%) | 6±1 (9±2%) | 1 | 12 |
| All | All | 1320/1700 (78%) | 946 (72%) | 255 (19%) | 119 (9%) | 1 | 12 |

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 70 | GLU | 20 |
| 1 | A | 75 | PRO | 20 |
| 1 | A | 15 | GLY | 17 |
| 1 | A | 73 | GLY | 16 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 69 | VAL | 15 |
| 1 | A | 55 | GLY | 13 |
| 1 | A | 52 | ILE | 5 |
| 1 | A | 36 | ILE | 3 |
| 1 | A | 54 | TYR | 3 |
| 1 | A | 14 | PHE | 3 |
| 1 | A | 63 | PRO | 2 |
| 1 | A | 51 | TYR | 2 |

### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|---|---|---|---|---|---|
| 1 | A | 58/70 (83%) | 40±2 (69±3%) | 18±2 (31±3%) | 1 15 |
| All | All | 1160/1400 (83%) | 803 (69%) | 357 (31%) | 1 15 |

All 36 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 46 | TRP | 20 |
| 1 | A | 51 | TYR | 20 |
| 1 | A | 61 | LEU | 20 |
| 1 | A | 81 | SER | 20 |
| 1 | A | 11 | HIS | 19 |
| 1 | A | 17 | LYS | 18 |
| 1 | A | 19 | PHE | 17 |
| 1 | A | 37 | SER | 17 |
| 1 | A | 77 | ASN | 16 |
| 1 | A | 35 | LYS | 15 |
| 1 | A | 42 | LYS | 15 |
| 1 | A | 38 | SER | 14 |
| 1 | A | 66 | TYR | 13 |
| 1 | A | 76 | ASP | 13 |
| 1 | A | 40 | LYS | 12 |
| 1 | A | 47 | ARG | 10 |
| 1 | A | 54 | TYR | 10 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 57 | ARG | 9 |
| 1 | A | 84 | GLN | 9 |
| 1 | A | 65 | GLU | 8 |
| 1 | A | 43 | SER | 7 |
| 1 | A | 50 | GLU | 7 |
| 1 | A | 14 | PHE | 6 |
| 1 | A | 10 | GLU | 6 |
| 1 | A | 70 | GLU | 6 |
| 1 | A | 83 | ARG | 5 |
| 1 | A | 53 | ASN | 5 |
| 1 | A | 80 | SER | 5 |
| 1 | A | 18 | SER | 3 |
| 1 | A | 12 | VAL | 3 |
| 1 | A | 78 | SER | 2 |
| 1 | A | 34 | ASP | 2 |
| 1 | A | 7 | ILE | 2 |
| 1 | A | 13 | ASN | 1 |
| 1 | A | 74 | ILE | 1 |
| 1 | A | 5 | GLU | 1 |

### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 7 Chemical shift validation (i)

No chemical shift data were provided