



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 28, 2023 – 02:51 PM EDT

PDB ID : 3K4M
Title : Pyranose 2-oxidase Y456W mutant in complex with 2FG
Authors : Divne, C.; Tan, T.C.
Deposited on : 2009-10-05
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	 4% 71% 19% • 8%
1	B	623	 4% 75% 16% • 7%
1	C	623	 7% 74% 17% • 8%
1	D	623	 4% 75% 15% • 8%
1	E	623	 7% 73% 18% • 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	623	
1	G	623	
1	H	623	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FAD	A	801	X	-	-	-
2	FAD	B	801	X	-	-	-
2	FAD	C	801	X	-	-	-
2	FAD	E	801	X	-	-	-
2	FAD	F	801	X	-	-	-
2	FAD	H	801	X	-	-	-
3	SHG	B	901	X	-	-	-
3	SHG	D	901	X	-	-	-
3	SHG	E	901	X	-	X	-
3	SHG	G	901	X	-	-	-
3	SHG	H	901	X	-	-	-
4	MES	B	902	-	-	X	-
4	MES	C	624	-	-	X	-
4	MES	C	902	-	-	X	-
4	MES	E	902	-	-	X	-
4	MES	H	624	-	-	X	-
4	MES	H	902	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 38766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	574	4528	2861	776	867	24	0	0	0
1	B	577	4551	2874	779	873	25	0	0	0
1	C	574	4527	2859	776	868	24	0	0	0
1	D	574	4527	2859	776	868	24	0	0	0
1	E	576	4544	2870	778	871	25	0	0	0
1	F	573	4520	2855	775	866	24	0	0	0
1	G	573	4520	2855	775	866	24	0	0	0
1	H	573	4520	2855	775	866	24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	456	TRP	TYR	engineered mutation	UNP Q7ZA32
B	456	TRP	TYR	engineered mutation	UNP Q7ZA32
C	456	TRP	TYR	engineered mutation	UNP Q7ZA32
D	456	TRP	TYR	engineered mutation	UNP Q7ZA32
E	456	TRP	TYR	engineered mutation	UNP Q7ZA32
F	456	TRP	TYR	engineered mutation	UNP Q7ZA32
G	456	TRP	TYR	engineered mutation	UNP Q7ZA32
H	456	TRP	TYR	engineered mutation	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 2-deoxy-2-fluoro-beta-D-glucopyranose (three-letter code: SHG) (formula: $C_6H_{11}FO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	O		
3	A	1	Total 12	C 6	F 1	O 5	0	0
3	B	1	Total 12	C 6	F 1	O 5	0	0
3	C	1	Total 12	C 6	F 1	O 5	0	0
3	D	1	Total 12	C 6	F 1	O 5	0	0
3	E	1	Total 12	C 6	F 1	O 5	0	0
3	F	1	Total 12	C 6	F 1	O 5	0	0
3	G	1	Total 12	C 6	F 1	O 5	0	0
3	H	1	Total 12	C 6	F 1	O 5	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	B	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	C	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	C	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	E	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	F	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	H	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	H	1	Total 12	C 6	N 1	O 4	S 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	303	Total 303	O 303	0	0
5	B	277	Total 277	O 277	0	0
5	C	191	Total 191	O 191	0	0
5	D	235	Total 235	O 235	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	204	Total 204	O 204	0	0
5	F	199	Total 199	O 199	0	0
5	G	230	Total 230	O 230	0	0
5	H	274	Total 274	O 274	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyranose 2-oxidase






- Molecule 1: Pyranose 2-oxidase







• Molecule 1: Pyranose 2-oxidase




• Molecule 1: Pyranose 2-oxidase





• Molecule 1: Pyranose 2-oxidase








• Molecule 1: Pyranose 2-oxidase





• Molecule 1: Pyranose 2-oxidase





• Molecule 1: Pyranose 2-oxidase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	168.19Å 103.14Å 168.71Å 90.00° 106.47° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.98 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-2.20) 97.4 (29.98-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5	Depositor
R, R_{free}	0.207 , 0.266 0.211 , 0.266	Depositor DCC
R_{free} test set	2798 reflections (1.02%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.022 for l,-k,h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	38766	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, SHG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.09	11/4645 (0.2%)	0.92	6/6317 (0.1%)
1	B	1.04	4/4668 (0.1%)	0.94	10/6348 (0.2%)
1	C	0.94	3/4644 (0.1%)	0.87	4/6316 (0.1%)
1	D	0.99	4/4644 (0.1%)	0.91	3/6316 (0.0%)
1	E	0.94	5/4661 (0.1%)	0.86	1/6338 (0.0%)
1	F	0.95	6/4637 (0.1%)	0.89	3/6306 (0.0%)
1	G	0.98	2/4637 (0.0%)	0.90	5/6306 (0.1%)
1	H	1.03	4/4637 (0.1%)	0.91	3/6306 (0.0%)
All	All	1.00	39/37173 (0.1%)	0.90	35/50553 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	H	0	1
All	All	0	2

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	432	GLU	CG-CD	8.23	1.64	1.51
1	D	478	GLU	CG-CD	7.61	1.63	1.51
1	E	542	GLU	CG-CD	7.35	1.62	1.51
1	A	481	GLU	CG-CD	7.33	1.62	1.51
1	D	310	GLU	CG-CD	7.32	1.62	1.51

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	139	ARG	NE-CZ-NH2	-9.79	115.41	120.30
1	F	139	ARG	NE-CZ-NH2	-8.90	115.85	120.30
1	F	180	ARG	NE-CZ-NH1	-7.73	116.44	120.30
1	H	81	ASP	CB-CG-OD1	-7.07	111.94	118.30
1	G	139	ARG	NE-CZ-NH1	6.69	123.64	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	449	ILE	Peptide
1	H	449	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4528	0	4376	91	0
1	B	4551	0	4396	82	0
1	C	4527	0	4372	93	0
1	D	4527	0	4372	73	0
1	E	4544	0	4389	105	0
1	F	4520	0	4365	86	0
1	G	4520	0	4365	98	0
1	H	4520	0	4365	97	0
2	A	53	0	28	2	0
2	B	53	0	29	5	0
2	C	53	0	26	5	0
2	D	53	0	27	6	0
2	E	53	0	27	9	0
2	F	53	0	26	1	0
2	G	53	0	27	3	0
2	H	53	0	26	3	0
3	A	12	0	10	2	0
3	B	12	0	10	3	0
3	C	12	0	11	5	0
3	D	12	0	9	4	0
3	E	12	0	10	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	12	0	11	1	0
3	G	12	0	10	3	0
3	H	12	0	10	1	0
4	A	12	0	12	4	0
4	B	12	0	12	7	0
4	C	24	0	24	21	0
4	E	12	0	12	12	0
4	F	12	0	12	1	0
4	H	24	0	24	23	0
5	A	303	0	0	9	0
5	B	277	0	0	10	0
5	C	191	0	0	2	0
5	D	235	0	0	6	0
5	E	204	0	0	4	0
5	F	199	0	0	2	0
5	G	230	0	0	6	0
5	H	274	0	0	12	0
All	All	38766	0	35393	708	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 708 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:HIS:HE2	2:B:801:FAD:C8M	1.09	1.59
1:B:167:HIS:NE2	2:B:801:FAD:HM82	1.06	1.38
1:C:133:ALA:CB	4:C:902:MES:H71	1.58	1.32
1:E:490:LYS:NZ	1:E:490:LYS:HB3	1.50	1.16
1:G:82:SER:HB2	1:G:83:GLY:CA	1.70	1.16

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	572/623 (92%)	548 (96%)	24 (4%)	0	100	100
1	B	575/623 (92%)	544 (95%)	31 (5%)	0	100	100
1	C	572/623 (92%)	545 (95%)	26 (4%)	1 (0%)	47	55
1	D	572/623 (92%)	548 (96%)	24 (4%)	0	100	100
1	E	574/623 (92%)	547 (95%)	26 (4%)	1 (0%)	47	55
1	F	571/623 (92%)	539 (94%)	31 (5%)	1 (0%)	47	55
1	G	571/623 (92%)	545 (95%)	26 (5%)	0	100	100
1	H	571/623 (92%)	555 (97%)	15 (3%)	1 (0%)	47	55
All	All	4578/4984 (92%)	4371 (96%)	203 (4%)	4 (0%)	51	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	343	ALA
1	H	187	ASP
1	C	81	ASP
1	F	344	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/542 (93%)	476 (95%)	26 (5%)	23	28
1	B	505/542 (93%)	489 (97%)	16 (3%)	39	50
1	C	502/542 (93%)	481 (96%)	21 (4%)	30	38
1	D	502/542 (93%)	478 (95%)	24 (5%)	25	32
1	E	504/542 (93%)	483 (96%)	21 (4%)	30	38
1	F	501/542 (92%)	479 (96%)	22 (4%)	28	35
1	G	501/542 (92%)	481 (96%)	20 (4%)	31	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	501/542 (92%)	478 (95%)	23 (5%)	27	34
All	All	4018/4336 (93%)	3845 (96%)	173 (4%)	29	36

5 of 173 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	296	GLU
1	G	385	THR
1	F	344	ASN
1	G	82	SER
1	H	100	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	341	ASN
1	G	461	GLN
1	F	460	GLN
1	G	341	ASN
1	H	263	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	801	1	53,58,58	1.29	4 (7%)	68,89,89	2.20	22 (32%)
3	SHG	E	901	-	12,12,12	1.37	2 (16%)	16,17,17	5.25	9 (56%)
3	SHG	C	901	-	12,12,12	0.73	0	16,17,17	3.04	6 (37%)
3	SHG	D	901	-	12,12,12	1.04	1 (8%)	16,17,17	4.88	8 (50%)
4	MES	B	902	-	12,12,12	1.94	1 (8%)	14,16,16	7.18	10 (71%)
3	SHG	H	901	-	12,12,12	0.98	1 (8%)	16,17,17	4.88	10 (62%)
4	MES	H	902	-	12,12,12	1.95	1 (8%)	14,16,16	6.90	8 (57%)
4	MES	F	902	-	12,12,12	1.54	1 (8%)	14,16,16	6.47	10 (71%)
4	MES	H	624	-	12,12,12	2.15	1 (8%)	14,16,16	8.23	9 (64%)
3	SHG	G	901	-	12,12,12	1.04	1 (8%)	16,17,17	4.93	10 (62%)
2	FAD	D	801	1	53,58,58	1.22	2 (3%)	68,89,89	2.83	28 (41%)
2	FAD	E	801	1	53,58,58	1.13	3 (5%)	68,89,89	2.15	18 (26%)
4	MES	A	902	-	12,12,12	1.96	1 (8%)	14,16,16	5.29	9 (64%)
3	SHG	A	901	-	12,12,12	0.65	0	16,17,17	3.73	9 (56%)
2	FAD	C	801	1	53,58,58	1.35	6 (11%)	68,89,89	2.49	22 (32%)
4	MES	E	902	-	12,12,12	2.09	1 (8%)	14,16,16	7.53	9 (64%)
2	FAD	H	801	1	53,58,58	1.22	6 (11%)	68,89,89	3.47	29 (42%)
4	MES	C	902	-	12,12,12	1.58	1 (8%)	14,16,16	6.81	10 (71%)
2	FAD	G	801	1	53,58,58	1.30	6 (11%)	68,89,89	2.22	20 (29%)
4	MES	C	624	-	12,12,12	2.17	2 (16%)	14,16,16	7.42	8 (57%)
3	SHG	B	901	-	12,12,12	1.18	1 (8%)	16,17,17	4.79	9 (56%)
2	FAD	F	801	1	53,58,58	1.42	9 (16%)	68,89,89	2.13	22 (32%)
2	FAD	A	801	1	53,58,58	1.34	9 (16%)	68,89,89	2.62	22 (32%)
3	SHG	F	901	-	12,12,12	0.81	0	16,17,17	3.31	10 (62%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	801	1	1/1/9/9	1/30/50/50	0/6/6/6
3	SHG	E	901	-	1/1/5/5	0/2/22/22	0/1/1/1
3	SHG	C	901	-	-	0/2/22/22	0/1/1/1
3	SHG	D	901	-	1/1/5/5	1/2/22/22	0/1/1/1
4	MES	B	902	-	-	5/6/14/14	0/1/1/1
3	SHG	H	901	-	1/1/5/5	2/2/22/22	0/1/1/1
4	MES	H	902	-	-	2/6/14/14	0/1/1/1
4	MES	F	902	-	-	2/6/14/14	0/1/1/1
4	MES	H	624	-	-	4/6/14/14	0/1/1/1
3	SHG	G	901	-	1/1/5/5	2/2/22/22	0/1/1/1
2	FAD	D	801	1	-	3/30/50/50	0/6/6/6
2	FAD	E	801	1	1/1/9/9	5/30/50/50	0/6/6/6
4	MES	A	902	-	-	3/6/14/14	0/1/1/1
3	SHG	A	901	-	-	0/2/22/22	0/1/1/1
2	FAD	C	801	1	2/2/9/9	1/30/50/50	0/6/6/6
4	MES	E	902	-	-	4/6/14/14	0/1/1/1
2	FAD	H	801	1	2/2/9/9	1/30/50/50	0/6/6/6
4	MES	C	902	-	-	4/6/14/14	0/1/1/1
2	FAD	G	801	1	-	4/30/50/50	0/6/6/6
4	MES	C	624	-	-	5/6/14/14	0/1/1/1
3	SHG	B	901	-	1/1/5/5	0/2/22/22	0/1/1/1
2	FAD	F	801	1	2/2/9/9	4/30/50/50	0/6/6/6
2	FAD	A	801	1	1/1/9/9	3/30/50/50	0/6/6/6
3	SHG	F	901	-	-	2/2/22/22	0/1/1/1

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	624	MES	C8-S	-7.09	1.67	1.77
4	C	624	MES	C8-S	-6.98	1.67	1.77
4	E	902	MES	C8-S	-6.80	1.67	1.77
4	A	902	MES	C8-S	-6.48	1.68	1.77
4	B	902	MES	C8-S	-6.36	1.68	1.77

The worst 5 of 327 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	624	MES	O1S-S-C8	-24.99	76.82	106.92
4	E	902	MES	O1S-S-C8	-19.17	83.83	106.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	902	MES	O3S-S-C8	-17.43	77.59	105.77
4	B	902	MES	O1S-S-C8	-17.39	85.98	106.92
4	C	902	MES	O3S-S-C8	-16.89	78.46	105.77

5 of 14 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	801	FAD	C4B
2	B	801	FAD	C4B
2	C	801	FAD	C1B
2	C	801	FAD	C2B
2	E	801	FAD	C4B

5 of 58 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	FAD	PA-O3P-P-O5'
2	D	801	FAD	PA-O3P-P-O5'
2	E	801	FAD	PA-O3P-P-O5'
2	F	801	FAD	PA-O3P-P-O5'
2	H	801	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

24 monomers are involved in 118 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	FAD	5	0
3	E	901	SHG	6	0
3	C	901	SHG	5	0
3	D	901	SHG	4	0
4	B	902	MES	7	0
3	H	901	SHG	1	0
4	H	902	MES	16	0
4	F	902	MES	1	0
4	H	624	MES	7	0
3	G	901	SHG	3	0
2	D	801	FAD	6	0
2	E	801	FAD	9	0
4	A	902	MES	4	0
3	A	901	SHG	2	0
2	C	801	FAD	5	0
4	E	902	MES	12	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	801	FAD	3	0
4	C	902	MES	10	0
2	G	801	FAD	3	0
4	C	624	MES	11	0
3	B	901	SHG	3	0
2	F	801	FAD	1	0
2	A	801	FAD	2	0
3	F	901	SHG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/623 (92%)	-0.06	22 (3%) 40 38	11, 18, 31, 52	0
1	B	577/623 (92%)	-0.06	23 (3%) 38 36	11, 19, 33, 45	0
1	C	574/623 (92%)	0.11	41 (7%) 16 14	13, 23, 39, 53	0
1	D	574/623 (92%)	0.00	26 (4%) 33 32	12, 21, 34, 48	0
1	E	576/623 (92%)	0.15	41 (7%) 16 14	15, 24, 40, 54	0
1	F	573/623 (91%)	0.10	29 (5%) 28 26	13, 23, 36, 49	0
1	G	573/623 (91%)	0.03	25 (4%) 34 32	14, 21, 36, 52	0
1	H	573/623 (91%)	-0.07	16 (2%) 53 51	12, 19, 32, 46	0
All	All	4594/4984 (92%)	0.02	223 (4%) 29 28	11, 21, 36, 54	0

The worst 5 of 223 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	619	THR	7.8
1	C	619	THR	6.6
1	C	389	LEU	6.4
1	C	343	ALA	5.6
1	B	619	THR	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	C	624	12/12	0.68	0.39	99,101,101,101	0
4	MES	H	902	12/12	0.73	0.31	70,77,78,78	0
4	MES	C	902	12/12	0.76	0.32	65,68,69,72	0
4	MES	E	902	12/12	0.83	0.29	76,78,79,81	0
4	MES	B	902	12/12	0.85	0.25	70,73,76,78	0
4	MES	H	624	12/12	0.89	0.21	68,71,73,73	0
3	SHG	G	901	12/12	0.93	0.20	41,44,48,49	0
3	SHG	E	901	12/12	0.94	0.20	43,46,48,50	0
3	SHG	A	901	12/12	0.94	0.22	32,35,40,43	0
4	MES	A	902	12/12	0.94	0.19	44,48,53,53	0
3	SHG	D	901	12/12	0.95	0.20	38,40,42,44	0
3	SHG	F	901	12/12	0.95	0.21	28,31,35,40	0
3	SHG	B	901	12/12	0.96	0.20	34,37,42,42	0
3	SHG	C	901	12/12	0.96	0.21	37,39,46,46	0
2	FAD	C	801	53/53	0.96	0.18	20,30,35,41	0
2	FAD	E	801	53/53	0.96	0.17	21,28,34,36	0
2	FAD	G	801	53/53	0.96	0.20	16,24,28,30	0
4	MES	F	902	12/12	0.96	0.13	38,40,41,43	0
2	FAD	B	801	53/53	0.96	0.20	16,26,31,33	0
3	SHG	H	901	12/12	0.96	0.23	39,41,45,49	0
2	FAD	A	801	53/53	0.97	0.19	16,21,26,27	0
2	FAD	F	801	53/53	0.97	0.19	13,27,36,41	0
2	FAD	D	801	53/53	0.97	0.18	16,25,29,34	0
2	FAD	H	801	53/53	0.97	0.20	17,25,31,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD E 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD G 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD F 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around FAD D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD H 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.