



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 10:26 am BST

PDB ID : 5K4U  
Title : Three-dimensional structure of L-threonine 3-dehydrogenase from *Trypanosoma brucei* showing different active site loop conformations between dimer subunits, refined to 1.9 angstroms  
Authors : Adjogatse, E.K.; Cooper, J.B.; Erskine, P.T.  
Deposited on : 2016-05-22  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	 3% 93% 7%
1	C	320	 7% 88% 11%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-threonine 3-dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	320	Total	C	N	O	S	0	0	0
			2501	1595	416	471	19			
1	C	320	Total	C	N	O	S	0	0	0
			2501	1595	416	471	19			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6002	MET	-	initiating methionine	UNP Q7YW97
C	6002	MET	-	initiating methionine	UNP Q7YW97

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	C	1	44	21	7	14	2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	C	1	6	3	3	0	0
3	C	1	6	3	3	0	0
3	C	1	6	3	3	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	125	Total O 125 125	0	0
5	C	103	Total O 103 103	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-threonine 3-dehydrogenase



- Molecule 1: L-threonine 3-dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.07Å 133.09Å 55.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.55 – 1.90 28.55 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (28.55-1.90) 97.7 (28.55-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.172 , 0.223 0.180 , 0.228	Depositor DCC
$R_{free}$ test set	2679 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.0	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 55.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5368	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.96	0/2557	1.01	6/3469 (0.2%)
1	C	0.91	0/2557	0.95	2/3469 (0.1%)
All	All	0.93	0/5114	0.98	8/6938 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6084	ARG	NE-CZ-NH2	-14.92	112.84	120.30
1	A	6084	ARG	NE-CZ-NH1	14.91	127.75	120.30
1	C	6084	ARG	NE-CZ-NH1	9.94	125.27	120.30
1	C	6084	ARG	NE-CZ-NH2	-8.41	116.10	120.30
1	A	6300	ASP	CB-CG-OD1	6.21	123.89	118.30
1	A	6282	ASP	CB-CG-OD1	5.99	123.69	118.30
1	A	6302	ASP	CB-CG-OD1	5.67	123.41	118.30
1	A	6058	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2501	0	2512	12	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2501	0	2512	23	0
2	A	44	0	26	1	0
2	C	44	0	26	0	0
3	A	12	0	16	0	0
3	C	18	0	24	3	0
4	A	16	0	12	1	0
4	C	4	0	3	0	0
5	A	125	0	0	0	0
5	C	103	0	0	0	0
All	All	5368	0	5131	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6084:ARG:HD2	1:A:6088:GLU:OE1	1.86	0.76
1:C:6177:ALA:HB3	3:C:6402:GOL:H12	1.75	0.68
1:C:6299:TYR:CE2	1:C:6303:MET:HE2	2.31	0.65
1:C:6306:GLU:OE2	1:C:6310:ARG:CZ	2.48	0.62
1:A:6198:LEU:HG	1:A:6318:LEU:HD13	1.81	0.61
1:A:6237:ARG:HA	4:A:6404:ACT:H2	1.85	0.59
1:A:6253:ARG:NH2	1:A:6264:ILE:O	2.39	0.55
1:C:6257:GLU:HG2	1:C:6262:ARG:O	2.09	0.53
1:C:6129:LYS:HE2	1:C:6280:TRP:O	2.09	0.52
1:A:6057:LEU:HD11	1:A:6095:ILE:HD13	1.92	0.50
1:C:6181:PRO:HB2	1:C:6191:HIS:CD2	2.46	0.49
1:C:6180:LEU:CD2	1:C:6316:HIS:CE1	2.96	0.48
1:A:6095:ILE:HD12	2:A:6401:NAD:H61A	1.78	0.48
1:C:6235:LEU:O	3:C:6403:GOL:H11	2.14	0.48
1:C:6177:ALA:HB3	3:C:6402:GOL:C1	2.43	0.47
1:A:6080:ILE:HG23	1:A:6084:ARG:HG2	1.96	0.47
1:A:6256:ILE:HG21	1:A:6264:ILE:HD11	1.97	0.47
1:C:6222:ASN:HD21	1:C:6226:LYS:NZ	2.15	0.45
1:C:6181:PRO:O	1:C:6191:HIS:CE1	2.69	0.45
1:C:6181:PRO:HB3	1:C:6190:ILE:HB	2.00	0.44
1:C:6312:ILE:N	1:C:6313:PRO:CD	2.81	0.44
1:A:6121:ALA:HB2	1:A:6280:TRP:CZ2	2.52	0.43
1:C:6156:THR:O	1:C:6160:GLN:HG3	2.18	0.43
1:C:6299:TYR:CD2	1:C:6303:MET:HE2	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6188:TYR:CZ	1:A:6249:PRO:HG3	2.54	0.43
1:C:6038:GLU:HG2	1:C:6053:LYS:HD2	2.01	0.43
1:C:6299:TYR:CG	1:C:6303:MET:HE1	2.55	0.42
1:C:6077:LEU:N	1:C:6078:PRO:CD	2.83	0.41
1:C:6252:LEU:HA	1:C:6301:ILE:HD11	2.03	0.41
1:A:6025:LYS:HE2	1:A:6229:GLU:OE2	2.21	0.41
1:C:6188:TYR:CZ	1:C:6249:PRO:HG3	2.56	0.41
1:A:6084:ARG:O	1:A:6084:ARG:HG3	2.21	0.40
1:C:6032:LEU:HD11	1:C:6052:GLU:HG2	2.02	0.40
1:C:6180:LEU:HD23	1:C:6180:LEU:N	2.37	0.40
1:C:6299:TYR:CD2	1:C:6303:MET:CE	3.05	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/320 (99%)	312 (98%)	6 (2%)	0	100	100
1	C	318/320 (99%)	310 (98%)	8 (2%)	0	100	100
All	All	636/640 (99%)	622 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/277 (100%)	274 (99%)	3 (1%)	73	73
1	C	277/277 (100%)	271 (98%)	6 (2%)	52	47
All	All	554/554 (100%)	545 (98%)	9 (2%)	62	60

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6062	PHE
1	A	6074	MET
1	A	6318	LEU
1	C	6062	PHE
1	C	6187	ASP
1	C	6235	LEU
1	C	6264	ILE
1	C	6275	LYS
1	C	6321	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6060	ASN
1	C	6060	ASN
1	C	6222	ASN
1	C	6278	ASN
1	C	6311	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ACT	A	6405	-	1,3,3	0.96	0	0,3,3	0.00	-
4	ACT	C	6405	-	1,3,3	3.39	1 (100%)	0,3,3	0.00	-
2	NAD	C	6401	-	42,48,48	2.20	7 (16%)	50,73,73	1.66	11 (22%)
4	ACT	A	6406	-	1,3,3	0.27	0	0,3,3	0.00	-
3	GOL	A	6402	-	5,5,5	0.44	0	5,5,5	1.08	0
3	GOL	C	6402	-	5,5,5	0.19	0	5,5,5	0.78	0
4	ACT	A	6407	-	1,3,3	0.25	0	0,3,3	0.00	-
2	NAD	A	6401	-	42,48,48	1.98	8 (19%)	50,73,73	1.33	7 (14%)
3	GOL	A	6403	-	5,5,5	0.15	0	5,5,5	1.58	1 (20%)
3	GOL	C	6404	-	5,5,5	0.63	0	5,5,5	1.16	1 (20%)
3	GOL	C	6403	-	5,5,5	0.49	0	5,5,5	1.07	0
4	ACT	A	6404	-	1,3,3	0.73	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	C	6401	-	-	1/26/62/62	0/5/5/5
3	GOL	A	6402	-	-	0/4/4/4	-
3	GOL	C	6402	-	-	2/4/4/4	-
2	NAD	A	6401	-	-	3/26/62/62	0/5/5/5
3	GOL	A	6403	-	-	2/4/4/4	-
3	GOL	C	6404	-	-	3/4/4/4	-
3	GOL	C	6403	-	-	1/4/4/4	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6401	NAD	C2N-N1N	8.52	1.45	1.35
2	A	6401	NAD	C2N-N1N	7.88	1.44	1.35
2	C	6401	NAD	O4D-C1D	6.42	1.50	1.41
2	C	6401	NAD	O7N-C7N	5.11	1.33	1.24
2	A	6401	NAD	O7N-C7N	4.27	1.32	1.24
2	A	6401	NAD	C7N-N7N	3.62	1.39	1.33
2	A	6401	NAD	C6N-N1N	3.43	1.43	1.35
4	C	6405	ACT	CH3-C	3.39	1.53	1.48
2	A	6401	NAD	C3N-C7N	2.99	1.55	1.50
2	C	6401	NAD	O3B-C3B	2.78	1.49	1.43
2	A	6401	NAD	PA-O2A	-2.48	1.43	1.55
2	A	6401	NAD	C2D-C1D	-2.32	1.50	1.53
2	C	6401	NAD	O2B-C2B	2.22	1.48	1.43
2	C	6401	NAD	C2A-N1A	-2.12	1.30	1.33
2	A	6401	NAD	O4B-C4B	2.09	1.49	1.45
2	C	6401	NAD	C3D-C4D	-2.00	1.47	1.53

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6401	NAD	O4D-C1D-C2D	-4.50	100.35	106.93
2	C	6401	NAD	N3A-C2A-N1A	-4.19	122.13	128.68
2	C	6401	NAD	C6N-N1N-C2N	-3.26	119.00	121.97
3	A	6403	GOL	O1-C1-C2	-3.02	95.72	110.20
2	C	6401	NAD	O7N-C7N-C3N	-2.95	116.11	119.63
2	A	6401	NAD	O4B-C1B-C2B	-2.92	102.65	106.93
2	C	6401	NAD	O5D-PN-O1N	2.82	120.08	109.07
2	A	6401	NAD	N3A-C2A-N1A	-2.81	124.29	128.68
2	C	6401	NAD	C4A-C5A-N7A	-2.75	106.53	109.40
2	C	6401	NAD	O4B-C4B-C3B	2.72	110.51	105.11
2	A	6401	NAD	O7N-C7N-C3N	-2.63	116.48	119.63
2	C	6401	NAD	O7N-C7N-N7N	2.63	126.32	122.58
2	A	6401	NAD	C5B-C4B-C3B	-2.49	105.85	115.18
2	C	6401	NAD	C2N-C3N-C4N	2.41	120.99	118.26
2	A	6401	NAD	C2A-N1A-C6A	2.32	122.72	118.75
2	A	6401	NAD	C3N-C7N-N7N	2.23	120.43	117.75
2	C	6401	NAD	O3B-C3B-C4B	-2.18	104.75	111.05
2	C	6401	NAD	O2N-PN-O1N	2.16	122.90	112.24
2	A	6401	NAD	C5N-C6N-N1N	-2.08	117.42	120.40
3	C	6404	GOL	O2-C2-C1	2.02	118.02	109.12

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	6401	NAD	C5D-O5D-PN-O1N
3	C	6402	GOL	C1-C2-C3-O3
3	A	6403	GOL	O1-C1-C2-C3
3	C	6404	GOL	O1-C1-C2-C3
3	C	6403	GOL	C1-C2-C3-O3
3	A	6403	GOL	O1-C1-C2-O2
3	C	6402	GOL	O2-C2-C3-O3
2	A	6401	NAD	C5D-O5D-PN-O3
3	C	6404	GOL	O2-C2-C3-O3
3	C	6404	GOL	O1-C1-C2-O2
2	C	6401	NAD	O4B-C4B-C5B-O5B
2	A	6401	NAD	O4B-C4B-C5B-O5B


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	6402	GOL	2	0
2	A	6401	NAD	1	0
3	C	6403	GOL	1	0
4	A	6404	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	320/320 (100%)	0.12	11 (3%) 45 48	11, 20, 41, 66	0
1	C	320/320 (100%)	0.29	21 (6%) 18 20	11, 21, 47, 66	0
All	All	640/640 (100%)	0.21	32 (5%) 28 32	11, 20, 45, 66	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	6321	LEU	11.8
1	C	6181	PRO	6.4
1	A	6320	SER	4.7
1	A	6321	LEU	4.5
1	C	6320	SER	4.0
1	C	6261	ASP	4.0
1	C	6182	GLY	3.8
1	C	6180	LEU	3.7
1	A	6318	LEU	3.2
1	C	6049	LYS	3.2
1	A	6265	GLU	3.0
1	C	6262	ARG	2.9
1	A	6269	VAL	2.8
1	A	6319	PRO	2.7
1	C	6259	CYS	2.7
1	A	6314	ILE	2.6
1	C	6318	LEU	2.6
1	A	6002	MET	2.5
1	C	6040	GLY	2.5
1	C	6258	ARG	2.5
1	C	6041	ALA	2.4
1	C	6042	LYS	2.4
1	A	6174	ILE	2.3
1	A	6116	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	6201	LYS	2.3
1	C	6199	GLN	2.2
1	C	6263	THR	2.2
1	C	6029	ASP	2.2
1	C	6116	ILE	2.1
1	C	6270	GLU	2.1
1	A	6317	GLY	2.1
1	C	6265	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	C	6404	6/6	0.76	0.19	39,43,44,45	0
3	GOL	C	6402	6/6	0.85	0.18	31,49,56,58	0
3	GOL	A	6403	6/6	0.88	0.16	26,34,35,43	0
3	GOL	C	6403	6/6	0.90	0.19	30,37,38,39	0
4	ACT	A	6404	4/4	0.91	0.13	28,35,42,43	0
4	ACT	A	6405	4/4	0.92	0.17	43,48,51,53	0
4	ACT	A	6407	4/4	0.92	0.17	49,49,52,59	0
4	ACT	A	6406	4/4	0.92	0.17	24,29,32,32	0
3	GOL	A	6402	6/6	0.95	0.09	20,25,28,29	0
2	NAD	C	6401	44/44	0.97	0.10	12,17,19,22	0
2	NAD	A	6401	44/44	0.97	0.10	12,15,18,18	0
4	ACT	C	6405	4/4	0.97	0.14	24,28,28,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers

There are no such residues in this entry.