



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 12:40 AM EDT

PDB ID : 4KCQ  
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with N-(4-(2-(ethyl(3-(thiophene-2-carboximidamido)benzyl)amino)ethyl)phenyl)thiophene-2-carboximidamide  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2013-04-24  
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	 4% 82% 8% • 9%
1	B	443	 3% 79% 11% •• 9%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7011 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	As	C	N	O				S
1	A	405	3223	1	2049	568	589	16	0	0	0
1	B	403	3212	1	2042	566	587	16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	conflict	UNP P29473
B	100	ARG	CYS	conflict	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	43	34	1	4	4	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is N-(4-{2-[ethyl(3-{{(E)-imino(thiophen-2-yl)methyl}amino}benzyl)amino]ethyl}phenyl)thiophene-2-carboximidamide (three-letter code: 1QF) (formula: C<sub>27</sub>H<sub>29</sub>N<sub>5</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N S 34 27 5 2	0	0
6	B	1	Total C N S 34 27 5 2	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Zn 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	192	Total O 192 192	0	0
8	B	167	Total O 167 167	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, endothelial



- Molecule 1: Nitric oxide synthase, endothelial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.74Å 106.41Å 156.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.33 – 2.03 48.28 – 2.03	Depositor EDS
% Data completeness (in resolution range)	98.2 (48.33-2.03) 98.2 (48.28-2.03)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 2.03Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.179 , 0.224 0.178 , 0.223	Depositor DCC
$R_{free}$ test set	3095 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.6	Xtrriage
Anisotropy	0.466	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7011	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: H4B, HEM, 1QF, GOL, ZN, CAS, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.65	0/3303	0.74	2/4497 (0.0%)
1	B	0.65	0/3291	0.75	4/4480 (0.1%)
All	All	0.65	0/6594	0.75	6/8977 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	240	ARG	NE-CZ-NH2	-6.27	117.16	120.30
1	B	240	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	B	240	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	A	240	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	B	330	LEU	CB-CG-CD1	5.55	120.43	111.00
1	B	378	LEU	CA-CB-CG	5.20	127.27	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3223	0	3127	26	0
1	B	3212	0	3117	33	0
2	A	43	0	30	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	43	0	30	6	0
3	A	17	0	15	1	0
3	B	17	0	15	1	0
4	A	8	0	6	0	0
4	B	8	0	6	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	34	0	27	1	0
6	B	34	0	27	3	0
7	A	1	0	0	0	0
8	A	192	0	0	3	0
8	B	167	0	0	4	0
All	All	7011	0	6416	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:384:CAS:SG	1:B:384:CAS:AS	2.46	1.34
1:A:384:CAS:SG	1:A:384:CAS:AS	2.50	1.29
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.56	0.87
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.64	0.80
1:B:107:LEU:HD21	6:B:503:1QF:H7	1.68	0.74
1:A:384:CAS:AS	1:A:384:CAS:CB	2.98	0.71
1:B:99:ARG:HG2	1:B:100:ARG:HD2	1.76	0.66
1:B:236:ARG:HG3	1:B:349:GLU:HB2	1.78	0.66
1:B:384:CAS:AS	1:B:384:CAS:CB	3.04	0.66
1:A:378:LEU:HB2	8:A:618:HOH:O	1.98	0.63
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.27	0.62
1:A:384:CAS:AS	1:A:384:CAS:HB2	2.61	0.60
1:B:186:CYS:HB2	2:B:501:HEM:ND	2.16	0.60
1:B:186:CYS:HB2	2:B:501:HEM:C4D	2.38	0.58
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.34	0.57
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.85	0.57
1:A:423:HIS:HB2	1:B:392:THR:HB	1.89	0.55
2:B:501:HEM:CMC	2:B:501:HEM:HBC2	2.37	0.54
1:A:240:ARG:HD2	1:A:241:GLY:O	2.08	0.54
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.48	0.53
1:B:378:LEU:HB2	8:B:624:HOH:O	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:501:HEM:HHC	2:B:501:HEM:HBB2	1.91	0.53
1:A:95:PRO:HG3	1:A:108:PRO:HG3	1.92	0.52
1:B:390:ARG:HB2	1:B:390:ARG:NH1	2.25	0.50
1:B:240:ARG:HD2	1:B:241:GLY:O	2.10	0.50
1:A:449:TRP:HA	3:A:502:H4B:N1	2.27	0.50
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.93	0.49
1:A:363:GLU:OE1	6:A:506:1QF:N27	2.41	0.47
1:B:340:ASN:HD22	1:B:340:ASN:H	1.63	0.47
2:B:501:HEM:HBC2	2:B:501:HEM:HMC1	1.98	0.46
1:B:384:CAS:AS	1:B:384:CAS:HB2	2.75	0.45
1:B:172:LEU:HD11	1:B:232:VAL:HG11	1.98	0.45
1:A:236:ARG:HH12	1:A:342:LEU:HD11	1.81	0.45
1:B:449:TRP:HA	3:B:502:H4B:N1	2.31	0.44
1:A:180:TRP:CE3	1:A:192:TRP:HA	2.53	0.44
1:A:72:ARG:HD3	8:A:744:HOH:O	2.16	0.44
1:A:138:SER:O	1:A:141:LYS:HE3	2.16	0.44
1:A:340:ASN:H	1:A:340:ASN:HD22	1.66	0.44
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.58	0.43
1:B:384:CAS:SG	1:B:384:CAS:CE1	3.05	0.43
1:A:392:THR:HB	1:B:423:HIS:HB2	1.99	0.43
1:A:455:SER:HB3	1:A:458:LEU:HD22	2.00	0.43
1:B:238:PRO:HG3	8:B:663:HOH:O	2.19	0.43
1:A:325:PHE:O	1:A:328:LEU:HB2	2.19	0.42
1:A:106:VAL:HG12	1:A:107:LEU:HG	2.01	0.42
2:A:501:HEM:CBB	2:A:501:HEM:HHC	2.50	0.42
1:A:126:LEU:HD12	1:A:126:LEU:HA	1.83	0.42
1:B:167:LEU:HG	1:B:348:LEU:HD12	2.02	0.42
1:B:146:GLN:OE1	1:B:146:GLN:HA	2.20	0.42
1:B:231:THR:O	1:B:353:ALA:HA	2.20	0.41
1:B:246:TRP:CZ2	1:B:302:PRO:HG3	2.55	0.41
2:B:501:HEM:HBD1	6:B:503:1QF:H22	2.03	0.41
6:B:503:1QF:H14	8:B:682:HOH:O	2.20	0.41
1:B:154:GLU:OE1	1:B:168:ARG:NH2	2.53	0.41
1:A:472:SER:HA	1:A:473:PRO:C	2.41	0.41
1:B:472:SER:HA	1:B:473:PRO:C	2.41	0.41
1:A:468:ASN:ND2	8:A:710:HOH:O	2.53	0.40
1:B:308:PRO:HD2	1:B:311:LEU:HD12	2.02	0.40
1:A:216:HIS:C	1:A:216:HIS:CD2	2.95	0.40
1:B:149:GLU:O	1:B:153:GLN:HG3	2.20	0.40
1:B:340:ASN:HD22	1:B:340:ASN:N	2.18	0.40
1:B:287:ARG:HD3	8:B:666:HOH:O	2.21	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:LEU:HD12	1:B:330:LEU:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/443 (90%)	387 (97%)	13 (3%)	0	100	100
1	B	398/443 (90%)	385 (97%)	12 (3%)	1 (0%)	41	36
All	All	798/886 (90%)	772 (97%)	25 (3%)	1 (0%)	51	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	261	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/375 (92%)	326 (95%)	17 (5%)	24	19
1	B	342/375 (91%)	325 (95%)	17 (5%)	24	19
All	All	685/750 (91%)	651 (95%)	34 (5%)	24	19

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	99	ARG
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	109	ARG
1	A	123	GLU
1	A	125	LEU
1	A	126	LEU
1	A	156	GLU
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	378	LEU
1	A	458	LEU
1	A	468	ASN
1	B	69	LYS
1	B	92	GLN
1	B	100	ARG
1	B	102	LEU
1	B	105	LEU
1	B	124	GLN
1	B	225	ASN
1	B	226	LEU
1	B	236	ARG
1	B	240	ARG
1	B	323	GLU
1	B	330	LEU
1	B	331	ARG
1	B	340	ASN
1	B	378	LEU
1	B	389	THR
1	B	458	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	376	ASN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CAS	B	384	1	5,8,9	0.99	0	1,9,11	0.64	0
1	CAS	A	384	1	5,8,9	0.86	0	1,9,11	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	B	384	1	-	0/0/7/9	-
1	CAS	A	384	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	384	CAS	4	0
1	A	384	CAS	3	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ACT	A	504	-	3,3,3	0.62	0	3,3,3	1.01	0
6	1QF	B	503	-	37,37,37	1.31	5 (13%)	33,49,49	1.78	3 (9%)
3	H4B	B	502	-	16,18,18	1.12	1 (6%)	11,26,26	2.51	7 (63%)
4	ACT	B	505	-	3,3,3	0.75	0	3,3,3	1.03	0
4	ACT	B	504	-	3,3,3	0.51	0	3,3,3	0.96	0
6	1QF	A	506	-	37,37,37	1.37	5 (13%)	33,49,49	2.18	3 (9%)
4	ACT	A	503	-	3,3,3	1.01	0	3,3,3	0.46	0
2	HEM	B	501	1	41,50,50	1.29	5 (12%)	45,82,82	1.91	11 (24%)
3	H4B	A	502	-	16,18,18	1.24	1 (6%)	11,26,26	2.75	7 (63%)
5	GOL	A	505	-	5,5,5	0.25	0	5,5,5	0.64	0
2	HEM	A	501	1	41,50,50	1.63	8 (19%)	45,82,82	1.95	12 (26%)
5	GOL	B	506	-	5,5,5	0.31	0	5,5,5	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	1QF	B	503	-	-	4/19/27/27	0/4/4/4
3	H4B	B	502	-	-	0/8/17/17	0/2/2/2
6	1QF	A	506	-	-	5/19/27/27	0/4/4/4
2	HEM	B	501	1	-	0/12/54/54	-
3	H4B	A	502	-	-	0/8/17/17	0/2/2/2
5	GOL	A	505	-	-	0/4/4/4	-
2	HEM	A	501	1	-	2/12/54/54	-
5	GOL	B	506	-	-	0/4/4/4	-

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C1B-NB	-4.67	1.32	1.40
6	A	506	1QF	C05-C06	3.69	1.50	1.45
6	B	503	1QF	C05-C06	3.63	1.50	1.45
2	B	501	HEM	C1B-NB	-3.54	1.34	1.40
2	A	501	HEM	C4D-C3D	3.53	1.51	1.45
6	B	503	1QF	C26-N27	-3.20	1.32	1.39
6	A	506	1QF	C26-N27	-3.09	1.32	1.39
2	A	501	HEM	FE-NB	3.04	2.11	1.96
2	A	501	HEM	C3B-C4B	3.00	1.50	1.44
6	A	506	1QF	C25-C26	2.98	1.49	1.45
2	A	501	HEM	C1D-C2D	2.86	1.50	1.44
6	B	503	1QF	C31-N27	-2.85	1.35	1.41
3	A	502	H4B	C4A-C4	-2.76	1.37	1.41
2	B	501	HEM	FE-NB	2.48	2.09	1.96
6	A	506	1QF	C11-N07	-2.47	1.36	1.41
2	B	501	HEM	CHB-C1B	2.39	1.41	1.35
6	B	503	1QF	C23-C22	2.30	1.41	1.34
3	B	502	H4B	C7-C6	2.28	1.54	1.52
2	A	501	HEM	C4B-NB	-2.27	1.34	1.38
6	B	503	1QF	C11-N07	-2.22	1.37	1.41
6	A	506	1QF	C06-N07	-2.14	1.34	1.39
2	A	501	HEM	C3C-C2C	2.12	1.43	1.40
2	A	501	HEM	C4A-NA	2.11	1.40	1.36
2	B	501	HEM	CHA-C4D	2.04	1.40	1.35
2	B	501	HEM	C3B-C4B	2.04	1.48	1.44

All (43) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	506	1QF	C23-C22-S21	-8.75	105.88	112.98
6	A	506	1QF	C03-C02-S01	-7.65	106.77	112.98
6	B	503	1QF	C03-C02-S01	-7.44	106.94	112.98
2	A	501	HEM	C1B-NB-C4B	5.78	111.05	105.07
2	A	501	HEM	CHC-C4B-NB	5.50	130.41	124.43
6	B	503	1QF	C23-C22-S21	-5.19	108.77	112.98
3	A	502	H4B	C8A-C4A-C4	4.53	118.60	114.57
2	B	501	HEM	CBA-CAA-C2A	-4.48	104.98	112.62
2	B	501	HEM	CHC-C4B-NB	3.94	128.71	124.43
3	A	502	H4B	C2-N3-C4	3.88	122.10	115.93
2	B	501	HEM	C1B-NB-C4B	3.87	109.08	105.07
3	A	502	H4B	N1-C2-N3	-3.67	119.66	125.42
2	B	501	HEM	CHD-C1D-C2D	-3.63	119.31	124.98
3	B	502	H4B	C2-N3-C4	3.62	121.69	115.93
2	B	501	HEM	CHA-C4D-ND	3.61	128.85	124.38
3	B	502	H4B	N1-C2-N3	-3.60	119.77	125.42
3	A	502	H4B	C4-C4A-N5	3.59	122.14	119.12
2	A	501	HEM	O2D-CGD-CBD	3.48	125.22	114.03
3	B	502	H4B	C4-C4A-N5	3.45	122.02	119.12
2	B	501	HEM	CHD-C1D-ND	3.35	128.07	124.43
2	A	501	HEM	CHA-C4D-ND	3.21	128.35	124.38
2	B	501	HEM	CHA-C4D-C3D	-3.15	119.42	125.33
2	A	501	HEM	CHA-C4D-C3D	-3.12	119.47	125.33
3	B	502	H4B	C8A-C4A-C4	2.97	117.20	114.57
3	B	502	H4B	N2-C2-N3	2.92	121.79	117.25
2	A	501	HEM	O2D-CGD-O1D	-2.78	116.37	123.30
2	A	501	HEM	CBA-CAA-C2A	-2.68	108.05	112.62
2	B	501	HEM	O2D-CGD-CBD	2.61	122.43	114.03
3	A	502	H4B	C2-N1-C8A	2.54	120.24	114.54
3	A	502	H4B	N2-C2-N3	2.53	121.19	117.25
2	A	501	HEM	CHD-C1D-C2D	-2.51	121.06	124.98
3	B	502	H4B	C2-N1-C8A	2.46	120.06	114.54
6	A	506	1QF	C17-N18-C38	-2.41	106.21	111.29
2	B	501	HEM	C4B-C3B-C2B	-2.35	105.25	107.11
2	A	501	HEM	O2A-CGA-O1A	-2.27	117.64	123.30
2	B	501	HEM	O2D-CGD-O1D	-2.17	117.89	123.30
3	B	502	H4B	C4A-N5-C6	-2.17	115.26	121.16
2	A	501	HEM	CHB-C1B-NB	2.16	127.05	124.38
2	A	501	HEM	C4B-C3B-C2B	-2.16	105.40	107.11
3	A	502	H4B	C4A-C4-N3	-2.08	118.09	124.01
2	A	501	HEM	CMD-C2D-C1D	2.04	128.14	125.04
6	B	503	1QF	C37-C38-N18	-2.03	106.29	114.38
2	B	501	HEM	C2D-C1D-ND	2.00	112.28	109.88

There are no chirality outliers.

All (11) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	A	506	1QF	C20-C19-N18-C17
6	B	503	1QF	C20-C19-N18-C17
6	A	506	1QF	C20-C19-N18-C38
6	B	503	1QF	C20-C19-N18-C38
6	A	506	1QF	C05-C06-N07-C11
6	A	506	1QF	C16-C11-N07-C06
6	A	506	1QF	C12-C11-N07-C06
6	B	503	1QF	C35-C34-C37-C38
6	B	503	1QF	C33-C34-C37-C38
2	A	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	503	1QF	3	0
3	B	502	H4B	1	0
6	A	506	1QF	1	0
2	B	501	HEM	6	0
3	A	502	H4B	1	0
2	A	501	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/443 (91%)	0.04	18 (4%) 33 32	25, 39, 66, 86	0
1	B	402/443 (90%)	-0.16	12 (2%) 50 50	25, 41, 69, 98	0
All	All	806/886 (90%)	-0.06	30 (3%) 41 41	25, 40, 68, 98	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	GLN	5.8
1	B	261	GLY	4.8
1	A	239	GLY	4.0
1	A	259	GLN	3.9
1	A	67	GLY	3.9
1	B	146	GLN	3.4
1	A	109	ARG	3.3
1	A	123	GLU	3.2
1	A	91	GLN	2.9
1	B	239	GLY	2.8
1	A	69	LYS	2.8
1	A	257	ARG	2.7
1	B	142	ARG	2.6
1	A	450	ILE	2.6
1	A	99	ARG	2.5
1	A	160	ALA	2.5
1	A	238	PRO	2.3
1	A	122	ALA	2.3
1	B	122	ALA	2.3
1	A	153	GLN	2.2
1	B	258	GLN	2.2
1	A	144	GLY	2.2
1	B	223	ARG	2.2
1	A	449	TRP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	187	VAL	2.1
1	B	109	ARG	2.1
1	B	257	ARG	2.0
1	B	144	GLY	2.0
1	B	150	GLU	2.0
1	A	146	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CAS	A	384	9/10	0.98	0.07	39,40,50,54	0
1	CAS	B	384	9/10	0.98	0.06	42,44,55,57	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	505	6/6	0.86	0.19	52,63,66,76	0
5	GOL	B	506	6/6	0.88	0.17	43,53,63,77	0
6	1QF	A	506	34/34	0.88	0.15	33,66,95,98	0
6	1QF	B	503	34/34	0.91	0.16	33,64,120,128	0
4	ACT	B	504	4/4	0.95	0.14	45,47,49,51	0
4	ACT	B	505	4/4	0.95	0.10	42,44,47,49	0
4	ACT	A	504	4/4	0.96	0.10	41,43,43,45	0
3	H4B	B	502	17/17	0.97	0.17	29,34,38,44	0
4	ACT	A	503	4/4	0.97	0.21	44,44,46,50	0
2	HEM	A	501	43/43	0.97	0.19	28,31,39,43	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	H4B	A	502	17/17	0.98	0.19	29,34,37,38	0
2	HEM	B	501	43/43	0.98	0.13	28,31,41,47	0
7	ZN	A	507	1/1	0.99	0.06	47,47,47,47	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around 1QF B 503:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.