



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2023 – 04:23 PM EDT

PDB ID : 5KGA
Title : X-ray structure of a glucosamine N-Acetyltransferase from *Clostridium acetobutylicum*, mutant D287N, in complex with N-acetylglucosamine
Authors : Dopkins, B.J.; Thoden, J.B.; Tipton, P.A.; Holden, H.M.
Deposited on : 2016-06-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	
1	B	328	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Predicted acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2628	1725	432	461	10	0	3	0
1	B	316	2614	1714	431	459	10	0	2	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	287	ASN	ASP	engineered mutation	UNP Q97ML2
A	321	GLY	-	expression tag	UNP Q97ML2
A	322	GLY	-	expression tag	UNP Q97ML2
A	323	HIS	-	expression tag	UNP Q97ML2
A	324	HIS	-	expression tag	UNP Q97ML2
A	325	HIS	-	expression tag	UNP Q97ML2
A	326	HIS	-	expression tag	UNP Q97ML2
A	327	HIS	-	expression tag	UNP Q97ML2
A	328	HIS	-	expression tag	UNP Q97ML2
B	287	ASN	ASP	engineered mutation	UNP Q97ML2
B	321	GLY	-	expression tag	UNP Q97ML2
B	322	GLY	-	expression tag	UNP Q97ML2
B	323	HIS	-	expression tag	UNP Q97ML2
B	324	HIS	-	expression tag	UNP Q97ML2
B	325	HIS	-	expression tag	UNP Q97ML2
B	326	HIS	-	expression tag	UNP Q97ML2
B	327	HIS	-	expression tag	UNP Q97ML2
B	328	HIS	-	expression tag	UNP Q97ML2

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucofuranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	15	8	1	6	0	0
2	B	1	15	8	1	6	0	0

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	32	11	5	13	3	0	0
3	A	1	22	9	1	10	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	B	1	22	9	1	10	2	0	0

- Molecule 4 is ACETYL COENZYME *A (three-letter code: ACO) (formula: $C_{23}H_{38}N_7O_{17}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
4	A	1	51	23	7	17	3	1	0	0
4	B	1	51	23	7	17	3	1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is TETRAMETHYLAMMONIUM ION (three-letter code: TMA) (formula: C₄H₁₂N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	N	0	0
			5	4	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	327	Total	O	0	0
			327	327		
8	B	248	Total	O	0	0
			248	248		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Predicted acetyltransferase

Chain A:  90% 6% .



- Molecule 1: Predicted acetyltransferase

Chain B:  91% 2% 2%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.00Å 65.60Å 89.90Å 90.00° 107.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 28.81 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.5 (50.00-1.90) 98.5 (28.81-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.170 , 0.224 0.180 , 0.230	Depositor DCC
R_{free} test set	2714 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtrriage
Anisotropy	0.129	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6039	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ACO, NAG, COA, TMA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2705	0.87	0/3635
1	B	0.52	0/2687	0.85	5/3612 (0.1%)
All	All	0.52	0/5392	0.86	5/7247 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	162	PRO	N-CA-CB	6.35	110.92	103.30
1	B	208	ARG	NE-CZ-NH2	6.08	123.34	120.30
1	B	208	ARG	NE-CZ-NH1	-5.93	117.33	120.30
1	B	7	TYR	CA-CB-CG	-5.83	102.33	113.40
1	B	227	ASP	CB-CG-OD1	5.80	123.52	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2628	0	2659	13	0
1	B	2614	0	2619	6	0
2	A	15	0	15	0	0
2	B	15	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	54	0	26	0	0
3	B	22	0	15	1	0
4	A	51	0	34	0	0
4	B	51	0	34	1	0
5	A	4	0	6	0	0
5	B	4	0	6	0	0
6	A	1	0	0	1	0
7	B	5	0	12	1	0
8	A	327	0	0	7	0
8	B	248	0	0	4	1
All	All	6039	0	5441	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ILE:HG22	1:B:45:MET:CE	2.05	0.87
6:A:406:CL:CL	8:A:760:HOH:O	2.31	0.84
1:B:3:ILE:HG22	1:B:45:MET:HE3	1.74	0.69
1:A:43:LYS:NZ	8:A:501:HOH:O	2.31	0.63
4:B:403:ACO:N3A	8:B:504:HOH:O	2.30	0.63
1:A:266:ARG:NH1	8:A:502:HOH:O	2.32	0.54
1:A:41:ARG:NH2	8:A:507:HOH:O	2.40	0.52
1:A:44:ILE:HG12	1:A:58:ILE:HG12	1.94	0.50
1:A:170:VAL:O	1:A:171:ASN:OD1	2.33	0.47
1:B:5:GLU:HG2	1:B:7:TYR:HE2	1.79	0.47
3:B:404:COA:C5P	8:B:603:HOH:O	2.63	0.47
1:A:240:LYS:HD2	1:A:247:TYR:CZ	2.50	0.46
1:A:97:LEU:HD23	1:A:139:ARG:CD	2.46	0.46
1:B:157:VAL:HG11	1:B:159:PHE:CZ	2.51	0.45
1:B:178:ARG:NH1	1:B:207:GLU:OE2	2.48	0.44
1:A:43:LYS:HE2	1:A:45:MET:HE3	2.00	0.43
1:A:40:GLN:HB3	1:A:61:GLN:O	2.18	0.43
7:B:402:TMA:C1	8:B:611:HOH:O	2.67	0.43
1:A:171:ASN:HB2	8:A:759:HOH:O	2.18	0.42
1:A:39:ARG:NH2	8:A:504:HOH:O	2.36	0.42
1:A:169:LYS:NZ	8:A:514:HOH:O	2.48	0.42
1:B:90:GLN:NE2	8:B:503:HOH:O	2.30	0.41
1:A:205:MET:SD	1:A:224:HIS:CE1	3.15	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:B:504:HOH:O	8:B:722:HOH:O[2_444]	1.97	0.23

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/328 (97%)	308 (97%)	9 (3%)	0	100	100
1	B	316/328 (96%)	304 (96%)	12 (4%)	0	100	100
All	All	633/656 (96%)	612 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/291 (98%)	282 (99%)	3 (1%)	73	73
1	B	279/291 (96%)	274 (98%)	5 (2%)	59	55
All	All	564/582 (97%)	556 (99%)	8 (1%)	67	65

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	LYS
1	A	95	LYS
1	A	154	VAL
1	B	2	GLU
1	B	65	LYS
1	B	70	LYS
1	B	178	ARG
1	B	217	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	86	ASN
1	A	144	GLN
1	A	274	GLN
1	B	49	ASN
1	B	144	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	COA	A	404	-	16,21,50	1.97	5 (31%)	24,31,75	1.11	2 (8%)
3	COA	B	404	-	16,21,50	2.26	7 (43%)	24,31,75	1.15	2 (8%)
4	ACO	A	403	-	45,53,53	0.81	1 (2%)	56,79,79	1.52	8 (14%)
2	NAG	B	401	-	15,15,15	0.32	0	21,21,21	1.44	2 (9%)
5	EDO	B	405	-	3,3,3	0.27	0	2,2,2	1.07	0
4	ACO	B	403	-	45,53,53	1.01	3 (6%)	56,79,79	1.43	10 (17%)
3	COA	A	402	-	29,34,50	0.84	1 (3%)	34,53,75	1.57	7 (20%)
2	NAG	A	401	-	15,15,15	0.43	0	21,21,21	1.34	3 (14%)
7	TMA	B	402	-	4,4,4	0.43	0	6,6,6	0.34	0
5	EDO	A	405	-	3,3,3	0.64	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	404	-	-	1/27/28/64	-
3	COA	B	404	-	-	3/27/28/64	-
4	ACO	A	403	-	-	2/47/67/67	0/3/3/3
2	NAG	B	401	-	-	0/6/26/26	0/1/1/1
5	EDO	B	405	-	-	0/1/1/1	-
4	ACO	B	403	-	-	0/47/67/67	0/3/3/3
3	COA	A	402	-	-	0/20/40/64	0/3/3/3
2	NAG	A	401	-	-	0/6/26/26	0/1/1/1
5	EDO	A	405	-	-	0/1/1/1	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	404	COA	C7P-C6P	-4.02	1.48	1.53
3	A	404	COA	O9P-C9P	-3.81	1.15	1.23
3	B	404	COA	O9P-C9P	-3.57	1.16	1.23
3	B	404	COA	P2A-O4A	-3.51	1.38	1.50
4	B	403	ACO	O4B-C1B	3.48	1.45	1.41
3	B	404	COA	P1A-O5B	-3.33	1.42	1.54
3	A	404	COA	P1A-O5B	-2.92	1.43	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	404	COA	P2A-O5A	-2.89	1.41	1.55
3	B	404	COA	P1A-O1A	-2.82	1.41	1.50
4	A	403	ACO	C5A-C4A	2.78	1.48	1.40
3	A	404	COA	P2A-O4A	-2.73	1.41	1.50
3	A	404	COA	C7P-C6P	-2.63	1.50	1.53
4	B	403	ACO	C5A-C4A	2.55	1.47	1.40
3	B	404	COA	P2A-O5A	-2.30	1.44	1.55
3	A	402	COA	C5A-C4A	2.27	1.46	1.40
3	B	404	COA	C9P-N8P	-2.26	1.28	1.33
4	B	403	ACO	C2A-N3A	2.03	1.35	1.32

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	ACO	C1B-N9A-C4A	-4.58	118.59	126.64
2	A	401	NAG	C1-C2-N2	4.30	115.71	110.73
4	A	403	ACO	N3A-C2A-N1A	-4.22	122.09	128.68
4	B	403	ACO	N3A-C2A-N1A	-4.21	122.10	128.68
4	B	403	ACO	C1B-N9A-C4A	-3.91	119.77	126.64
2	B	401	NAG	C1-C2-N2	3.80	115.13	110.73
3	A	402	COA	N3A-C2A-N1A	-3.80	122.74	128.68
4	A	403	ACO	C2A-N1A-C6A	3.54	124.82	118.75
3	A	402	COA	C1B-N9A-C4A	-3.52	120.45	126.64
3	A	402	COA	O9A-P3B-O8A	3.23	119.97	107.64
4	B	403	ACO	O9A-P3B-O8A	3.11	119.53	107.64
4	B	403	ACO	N6A-C6A-N1A	2.77	124.33	118.57
4	A	403	ACO	C4A-C5A-N7A	-2.67	106.61	109.40
2	A	401	NAG	C1-O5-C5	-2.64	108.69	113.66
2	B	401	NAG	O7-C7-N2	2.60	126.72	121.95
3	A	402	COA	C2A-N1A-C6A	2.58	123.17	118.75
4	A	403	ACO	CDP-CBP-CAP	2.53	113.20	108.82
4	B	403	ACO	C5A-C6A-N6A	-2.49	116.58	120.35
4	B	403	ACO	O5P-C5P-C6P	-2.48	117.49	122.02
3	A	402	COA	O2A-P1A-O1A	2.45	124.37	112.24
3	B	404	COA	O5A-P2A-O4A	2.41	124.15	112.24
2	A	401	NAG	C3-C2-N2	-2.39	106.10	110.62
3	A	402	COA	N6A-C6A-N1A	2.32	123.39	118.57
3	A	404	COA	O5B-P1A-O1A	2.29	119.65	110.68
4	B	403	ACO	C2A-N1A-C6A	2.28	122.66	118.75
4	A	403	ACO	P2A-O3A-P1A	-2.28	125.00	132.83
3	A	404	COA	O6A-CCP-CBP	-2.28	106.89	110.55
4	B	403	ACO	C2P-C3P-N4P	-2.28	107.63	112.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	403	ACO	P2A-O3A-P1A	-2.22	125.21	132.83
4	A	403	ACO	C5A-C6A-N6A	2.20	123.69	120.35
3	B	404	COA	CEP-CBP-CDP	2.17	113.59	109.17
4	A	403	ACO	O2B-C2B-C1B	-2.11	103.05	110.85
3	A	402	COA	O2A-P1A-O5B	2.11	117.54	107.75
4	B	403	ACO	O2A-P1A-O1A	2.00	122.14	112.24

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	404	COA	C5P-C6P-C7P-N8P
3	B	404	COA	CAP-C9P-N8P-C7P
3	B	404	COA	O9P-C9P-N8P-C7P
4	A	403	ACO	C6P-C7P-N8P-C9P
4	A	403	ACO	C3B-O3B-P3B-O9A
3	B	404	COA	CCP-O6A-P2A-O4A


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	404	COA	1	0
4	B	403	ACO	1	0
7	B	402	TMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/328 (96%)	-0.46	1 (0%) 94 94	8, 15, 35, 48	0
1	B	316/328 (96%)	-0.27	5 (1%) 72 74	8, 18, 43, 55	0
All	All	632/656 (96%)	-0.36	6 (0%) 84 85	8, 16, 38, 55	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	204	GLY	5.8
1	B	159	PHE	3.2
1	B	179	LYS	2.5
1	B	164	ILE	2.4
1	A	204	GLY	2.2
1	B	316	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	B	405	4/4	0.91	0.12	34,36,36,40	0
6	CL	A	406	1/1	0.94	0.07	38,38,38,38	0
7	TMA	B	402	5/5	0.94	0.20	31,31,32,34	0
4	ACO	A	403	51/51	0.95	0.10	9,15,35,51	0
2	NAG	B	401	15/15	0.96	0.08	12,15,18,18	0
2	NAG	A	401	15/15	0.97	0.10	15,16,18,20	0
4	ACO	B	403	51/51	0.97	0.07	10,14,19,21	0
5	EDO	A	405	4/4	0.97	0.09	20,23,24,24	0
3	COA	A	402	32/48	0.97	0.12	15,21,43,46	0
3	COA	A	404	22/48	0.97	0.07	18,24,32,38	0
3	COA	B	404	22/48	0.97	0.08	20,26,37,50	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around COA A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COA A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around COA B 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.