

# Full wwPDB X-ray Structure Validation Report (i)

#### May 24, 2020 – 07:48 pm BST

:	3KH5
:	Crystal Structure of Protein MJ1225 from Methanocaldococcus jannaschii, a
	putative archaeal homolog of g-AMPK.
:	Gomez Garcia, I.; Oyenarte, I.; Martinez-Cruz, L.A.
:	2009-10-30
:	2.10  Å(reported)
	: : :

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
$R_{free}$	130704	5197(2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647(2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			7%		
1	А	280	76%	19%	••



#### $3 \mathrm{KH5}$

# 2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 2486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called protein MJ1225.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	278	Total 2211	C 1390	N 389	0 421	S 11	0	1	0


• Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
2	Δ	1	Total	С	Ν	Ο	Р	0	0
	Л	L	27	10	5	10	2	0	0
9	Λ	1	Total	С	Ν	Ο	Р	0	0
	Л	I	27	10	5	10	2	0	0
9	Λ	1	Total	С	Ν	Ο	Р	0	0
	Л	I	27	10	5	10	2	0	0

• Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
9	Δ	1	Total	С	Ν	Ο	Р	0	0
0	А	L	23	10	5	7	1	0	0
2	٨	1	Total	С	Ν	Ο	Р	0	0
0	А	L	23	10	5	7	1	0	0



Mol	Chain	Residues	Atom	.S	ZeroOcc	AltConf
4	А	1	Total C 5 4	) S 1	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	143	Total O 143 143	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: protein MJ1225



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	108.55Å $108.55$ Å $143.63$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	50.00 - 2.10	Depositor
Resolution (A)	39.33 - 2.10	EDS
% Data completeness	99.7(50.00-2.10)	Depositor
(in resolution range)	99.7(39.33-2.10)	EDS
R <sub>merge</sub>	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$< I/\sigma(I) > 1$	$7.17 (at 2.10 \text{\AA})$	Xtriage
Refinement program	REFMAC $5.5.0072$	Depositor
$R, R_{free}$	0.228 , $0.294$	Depositor
$\Pi, \Pi_{free}$	0.225 , $0.288$	DCC
$R_{free}$ test set	983 reflections $(5.12\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	41.1	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36 , $45.4$	EDS
L-test for twinning <sup>2</sup>	$ L  > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2486	wwPDB-VP
Average B, all atoms $(Å^2)$	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
IVIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.43	0/2233	0.64	$4/3011 \ (0.1\%)$

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	181	ARG	NE-CZ-NH2	-7.56	116.52	120.30
1	А	181	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	А	44	GLY	N-CA-C	-6.12	97.79	113.10
1	А	181	ARG	CG-CD-NE	-5.90	99.41	111.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	43	ALA	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2211	0	2299	45	0
2	А	81	0	36	4	0
3	А	46	0	24	0	0
4	А	5	0	0	1	0
5	А	143	0	0	5	1
All	All	2486	0	2359	47	1

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A.4.5.55. D	Interatomic	Clash
Atom-1	Atom-2	${ m distance}~({ m \AA})$	overlap (Å)
1:A:148:GLU:HG3	1:A:278:TYR:HE1	1.41	0.85
1:A:148:GLU:HG3	1:A:278:TYR:CE1	2.12	0.83
1:A:55:MET:HE3	1:A:196:SER:HB3	1.63	0.81
1:A:232:ASP:HB3	5:A:426:HOH:O	1.82	0.79
1:A:164:GLY:N	4:A:285:SO4:O3	2.13	0.79
1:A:55:MET:CE	1:A:196:SER:HB3	2.14	0.78
1:A:12:LYS:HG2	1:A:13:LYS:O	1.87	0.73
2:A:281:ADP:H3'	2:A:281:ADP:O2A	1.92	0.69
2:A:284:ADP:O1A	5:A:420:HOH:O	2.15	0.64
1:A:100:ASN:HB3	5:A:402:HOH:O	1.97	0.64
1:A:190:LEU:HB2	1:A:266:ILE:HD11	1.79	0.64
1:A:75:GLU:O	1:A:76:ARG:HB2	2.00	0.61
1:A:190:LEU:HB2	1:A:266:ILE:CD1	2.31	0.61
1:A:13:LYS:H	1:A:13:LYS:HD2	1.67	0.59
1:A:30:MET:HE2	1:A:36:ARG:HA	1.85	0.58
1:A:215:GLY:HA2	1:A:216:ASN:C	2.24	0.57
1:A:12:LYS:CG	1:A:13:LYS:N	2.68	0.56
1:A:181:ARG:HB3	1:A:195:THR:HG22	1.85	0.56
1:A:4:ARG:HB2	1:A:7:LYS:HD2	1.86	0.56
1:A:209:PHE:O	1:A:213:GLN:HG2	2.06	0.56
1:A:37:ARG:HB2	2:A:281:ADP:O3B	2.06	0.56
1:A:55:MET:HG3	2:A:283:ADP:O1A	2.06	0.55
1:A:78:PHE:HB3	1:A:217:VAL:HG11	1.89	0.53
1:A:43:ALA:N	1:A:44:GLY:HA3	2.28	0.48
1:A:79:LEU:CD2	1:A:212:MET:HB3	2.43	0.48
1:A:30:MET:HE1	1:A:54:SER:HB3	1.95	0.48

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:ARG:CZ	1:A:265:ARG:HD2	2.44	0.47
1:A:261:ASP:HB3	1:A:263:ASN:H	1.80	0.47
1:A:31:ASN:OD1	1:A:200:ILE:HG22	2.15	0.47
1:A:219:GLU:O	1:A:223:VAL:HG23	2.15	0.47
1:A:134:ARG:HG3	1:A:250:VAL:HG22	1.97	0.47
1:A:45:ASN:HD22	1:A:47:LYS:H	1.63	0.47
1:A:79:LEU:HD11	1:A:212:MET:SD	2.56	0.46
1:A:181:ARG:NH2	5:A:397:HOH:O	2.49	0.46
1:A:61:MET:HE1	1:A:199:PHE:HB3	1.98	0.46
1:A:27:LEU:HD21	1:A:57:ILE:HG21	1.97	0.46
1:A:209:PHE:CE2	1:A:213:GLN:NE2	2.84	0.46
1:A:75:GLU:O	1:A:76:ARG:CB	2.65	0.45
1:A:147:ASN:HA	1:A:241:LYS:HE2	1.97	0.44
1:A:189:ARG:NH1	1:A:265:ARG:HD2	2.33	0.44
1:A:5:VAL:HG22	1:A:101:ALA:O	2.18	0.43
1:A:79:LEU:HD22	1:A:212:MET:HB3	2.01	0.43
1:A:12:LYS:HG3	1:A:13:LYS:N	2.34	0.43
1:A:31:ASN:OD1	1:A:200:ILE:CG2	2.67	0.42
1:A:261:ASP:OD2	1:A:267:LYS:HD3	2.19	0.42
1:A:230:LYS:NZ	5:A:334:HOH:O	2.40	0.41
1:A:156:ARG:HD3	1:A:157:ASP:OD1	2.21	0.41

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	${f Interatomic} \ {f distance} \ ({ m \AA})$	Clash overlap (Å)
5:A:350:HOH:O	5:A:422:HOH:O[12_555]	2.09	0.11

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	277/280 (99%)	271 (98%)	6 (2%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	247/248~(100%)	227~(92%)	20 (8%)	11 8

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	4	ARG
1	А	10	GLN
1	А	13	LYS
1	А	36	ARG
1	А	45	ASN
1	А	79	LEU
1	А	124	GLU
1	А	130	LEU
1	А	148	GLU
1	А	149	VAL
1	А	156	ARG
1	А	160	VAL
1	А	172	ARG
1	А	181	ARG
1	А	187	GLU
1	А	210	ASN
1	А	237	LYS
1	А	250	VAL
1	А	262	GLU
1	А	265	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	А	11	ASN
1	А	42	ASN
1	А	45	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Res Link		Bo	Bond lengths			Bond angles		
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	#  Z  > 2	
3	AMP	А	1605	-	22,25,25	1.20	2 (9%)	$25,\!38,\!38$	1.28	2 (8%)	
3	AMP	А	282	-	22,25,25	0.95	2 (9%)	$25,\!38,\!38$	1.44	3 (12%)	
2	ADP	А	283	-	24,29,29	1.19	2 (8%)	$29,\!45,\!45$	2.20	<mark>5 (17%)</mark>	
2	ADP	А	284	-	24, 29, 29	1.23	2 (8%)	$29,\!45,\!45$	1.17	2 (6%)	
4	SO4	А	285	-	$4,\!4,\!4$	0.16	0	$^{6,6,6}$	0.08	0	
2	ADP	А	281	-	24, 29, 29	1.29	2 (8%)	$29,\!45,\!45$	2.30	<mark>6 (20%)</mark>	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	А	1605	-	-	3/6/26/26	0/3/3/3
2	ADP	А	283	-	-	0/12/32/32	0/3/3/3
2	ADP	А	284	-	-	1/12/32/32	0/3/3/3
3	AMP	A	282	-	-	3/6/26/26	0/3/3/3
2	ADP	А	281	-	-	4/12/32/32	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	А	281	ADP	O4'-C1'	4.16	1.46	1.41
3	А	1605	AMP	O4'-C1'	3.85	1.46	1.41
2	А	284	ADP	O4'-C1'	3.63	1.46	1.41
2	А	283	ADP	O4'-C1'	3.03	1.45	1.41
2	А	284	ADP	PB-O2B	2.71	1.65	1.54
2	А	283	ADP	PB-O2B	2.20	1.63	1.54
3	А	282	AMP	O4'-C1'	2.14	1.44	1.41
2	А	281	ADP	PB-O2B	2.07	1.62	1.54
3	А	1605	AMP	P-O2P	2.05	1.62	1.54
3	А	282	AMP	P-O3P	2.01	1.62	1.54

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	281	ADP	O2A-PA-O5'	-6.21	78.92	107.75
2	А	283	ADP	O5'-PA-O1A	-5.90	86.00	109.07
2	А	281	ADP	O5'-PA-O1A	-5.77	86.54	109.07
2	А	283	ADP	O2A-PA-O5'	-5.64	81.53	107.75
2	А	283	ADP	PA-O3A-PB	-5.61	113.59	132.83
2	А	283	ADP	N3-C2-N1	-4.77	121.22	128.68
3	А	282	AMP	N3-C2-N1	-4.75	121.25	128.68
3	А	1605	AMP	N3-C2-N1	-4.68	121.37	128.68
2	А	281	ADP	N3-C2-N1	-4.63	121.44	128.68
2	А	281	ADP	PA-O3A-PB	-4.57	117.15	132.83
2	А	284	ADP	N3-C2-N1	-4.38	121.83	128.68
2	А	281	ADP	C3'-C2'-C1'	3.72	106.58	100.98
3	А	282	AMP	O5'-P-O1P	2.71	114.08	106.47
3	А	1605	AMP	O5'-P-O1P	2.49	113.47	106.47
2	А	284	ADP	C4-C5-N7	-2.13	107.18	109.40
2	А	281	ADP	C4-C5-N7	-2.09	107.22	109.40
2	А	283	ADP	C1'-N9-C4	-2.05	123.04	126.64
3	А	282	AMP	C5'-C4'-C3'	-2.03	107.58	115.18



There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
3	А	282	AMP	C5'-O5'-P-O1P
3	А	282	AMP	C5'-O5'-P-O2P
3	А	282	AMP	C5'-O5'-P-O3P
2	А	281	ADP	C5'-O5'-PA-O2A
2	А	281	ADP	C5'-O5'-PA-O3A
3	А	1605	AMP	O4'-C4'-C5'-O5'
3	А	1605	AMP	C3'-C4'-C5'-O5'
2	А	281	ADP	C3'-C4'-C5'-O5'
2	А	281	ADP	O4'-C4'-C5'-O5'
3	А	1605	AMP	C5'-O5'-P-O2P
2	А	284	ADP	C5'-O5'-PA-O1A

All (11) torsion outliers are listed below:


There are no ring outliers.

4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	283	ADP	1	0
2	А	284	ADP	1	0
4	А	285	SO4	1	0
2	А	281	ADP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	$\#RSRZ{>}2$		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9	
1	А	278/280~(99%)	0.49	20 (7%)	15	19	28,  41,  71,  87	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	11	ASN	6.7
1	А	9	ALA	4.2
1	А	156	ARG	3.7
1	А	8	ILE	3.6
1	А	214	THR	3.6
1	А	10	GLN	3.5
1	А	7	LYS	3.5
1	А	215	GLY	3.4
1	А	218	ARG	3.4
1	А	216	ASN	3.4
1	А	217	VAL	3.1
1	А	261	ASP	2.7
1	А	123	ASP	2.5
1	А	36	ARG	2.4
1	А	280	ALA	2.3
1	А	159	ILE	2.3
1	А	262	GLU	2.3
1	А	34	LYS	2.2
1	А	35	TYR	2.1
1	А	3	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	$\mathbf{RSR}$	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	AMP	A	1605	23/23	0.78	0.22	62,70,78,78	0
2	ADP	А	281	27/27	0.79	0.17	$56,\!68,\!84,\!84$	0
2	ADP	А	284	27/27	0.90	0.18	$35,\!37,\!47,\!47$	0
3	AMP	А	282	23/23	0.90	0.14	$35,\!41,\!51,\!51$	0
4	SO4	А	285	5/5	0.95	0.16	53, 53, 53, 54	0
2	ADP	А	283	27/27	0.95	0.15	27,28,39,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
























## 6.5 Other polymers (i)

There are no such residues in this entry.

