



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2020 – 09:40 AM BST

PDB ID : 6KKO  
Title : The crystal structure of SiaB-SiaC complex from *Pseudomonas aeruginosa*  
Authors : Gan, J.H.; Yang, C.  
Deposited on : 2019-07-26  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	
1	B	181	
2	C	127	
2	D	127	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative serine phosphatase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	170	1295	813	237	240	2	3	0	1	0
1	B	168	1291	808	235	243	1	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP A0A485GYA3
B	0	SER	-	expression tag	UNP A0A485GYA3

- Molecule 2 is a protein called DUF1987 domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
			Total	C	N	O	P	S				Se
2	C	127	1030	642	179	203	1	1	4	0	0	0
2	D	126	1017	636	175	200	1	1	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP A0A072ZHB4
D	0	SER	-	expression tag	UNP A0A072ZHB4

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	27	10	5	10	2	0	0
3	B	1	27	10	5	10	2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	B	1	1	1	0	0
4	C	1	1	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	39	39	39	0	0
5	B	26	26	26	0	0
5	C	42	42	42	0	0
5	D	28	28	28	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative serine phosphatase




- Molecule 1: Putative serine phosphatase



- Molecule 2: DUF1987 domain-containing protein



- Molecule 2: DUF1987 domain-containing protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.62Å 42.02Å 96.58Å 90.00° 93.45° 90.00°	Depositor
Resolution (Å)	37.74 – 2.10 37.74 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (37.74-2.10) 99.2 (37.74-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.232 , 0.278 0.239 , 0.281	Depositor DCC
$R_{free}$ test set	1646 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.3	Xtrriage
Anisotropy	0.514	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, NA, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/1313	0.51	0/1767
1	B	0.33	0/1303	0.51	1/1749 (0.1%)
2	C	0.33	0/1038	0.50	0/1397
2	D	0.51	0/1025	0.65	2/1380 (0.1%)
All	All	0.38	0/4679	0.54	3/6293 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	26	MSE	O-C-N	-11.22	104.74	122.70
2	D	26	MSE	CA-C-N	7.42	133.51	117.20
1	B	28	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	26	MSE	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1295	0	1290	24	0
1	B	1291	0	1285	27	0
2	C	1030	0	965	11	0
2	D	1017	0	951	13	0
3	A	27	0	12	1	0
3	B	27	0	12	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	39	0	0	0	0
5	B	26	0	0	0	0
5	C	42	0	0	0	0
5	D	28	0	0	1	0
All	All	4824	0	4515	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:MSE:HE2	1:A:96:VAL:HG12	1.60	0.83
1:A:3:THR:HG22	1:B:16:GLN:HG2	1.63	0.81
1:B:130:LYS:HA	1:B:133:LEU:HD23	1.65	0.78
1:B:50:SER:HB2	2:D:103:ARG:HG2	1.65	0.77
2:D:75:MSE:HE1	2:D:114:ASP:OD2	1.89	0.73
1:A:95:VAL:HG11	1:B:2:GLU:HG2	1.69	0.73
1:B:48:LYS:HD3	1:B:51:GLU:H	1.55	0.70
1:A:62:MSE:CE	1:A:96:VAL:HG12	2.23	0.68
2:C:13:PRO:HD2	2:C:38:LEU:HD21	1.77	0.66
1:B:26:ILE:HD12	1:B:30:LEU:HB3	1.78	0.66
2:D:13:PRO:HD2	2:D:38:LEU:HD21	1.79	0.65
1:A:104:ARG:HD2	1:A:166:GLU:HB3	1.79	0.64
1:B:28:ARG:CA	1:B:28:ARG:HE	2.14	0.61
1:A:110:LEU:HD11	3:A:1001:ADP:C4	2.37	0.60
1:A:133:LEU:HD13	2:C:75:MSE:HE3	1.83	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:VAL:HG21	1:B:180:ILE:HD11	1.84	0.57
1:A:175:SER:HB2	1:B:6:LEU:HD23	1.88	0.56
1:B:134:ARG:NH1	2:D:114:ASP:HB3	2.21	0.56
1:A:28:ARG:NH2	2:C:33:GLU:H	2.03	0.55
1:B:50:SER:HB2	2:D:103:ARG:CG	2.36	0.55
1:A:117:ILE:HD11	1:A:145:ALA:HB1	1.88	0.54
1:A:152:ILE:HG22	1:A:160:LEU:HD11	1.89	0.53
1:B:163:SER:OG	1:B:175:SER:OG	2.27	0.53
1:B:16:GLN:HB2	1:B:18:ILE:HG13	1.90	0.53
1:A:75:TYR:HB3	1:A:79:GLU:HB3	1.89	0.52
1:A:21[A]:CYS:SG	1:B:23:ASN:HB3	2.50	0.51
1:A:153:ALA:HA	1:A:160:LEU:HD13	1.94	0.50
2:C:109:GLU:O	2:C:113:GLU:HG3	2.13	0.48
1:A:23:ASN:HB3	1:B:21:CYS:SG	2.54	0.48
1:A:31:ILE:CD1	1:A:63:THR:HB	2.45	0.47
2:C:49:PHE:CZ	2:C:55:ARG:HG3	2.49	0.47
2:D:15:ILE:HG12	2:D:26:MSE:HG2	1.98	0.46
1:A:58:VAL:HG21	1:A:180:ILE:HD11	1.99	0.45
2:D:82:GLU:OE2	2:D:116:SER:OG	2.33	0.45
2:C:23:ARG:HB3	2:C:23:ARG:NH1	2.31	0.45
1:A:26:ILE:HD12	1:A:30:LEU:HB3	1.99	0.44
2:C:20:GLN:HA	2:C:55:ARG:NH1	2.33	0.44
2:D:26:MSE:SE	2:D:74:MSE:HE1	2.67	0.44
2:C:63:LEU:HB3	2:C:66:LEU:HD11	1.99	0.44
2:D:109:GLU:O	2:D:113:GLU:HG3	2.18	0.43
1:B:134:ARG:HH12	2:D:114:ASP:HB3	1.82	0.43
1:A:114:ILE:HD13	1:A:176:LEU:HD13	2.00	0.43
2:D:62:ARG:HG2	2:D:96:HIS:HB2	2.00	0.43
2:C:97:TYR:CZ	2:C:105:ALA:HB2	2.54	0.42
1:B:12:SER:O	1:B:16:GLN:HG3	2.18	0.42
1:B:19:LEU:HD22	1:B:41:TYR:OH	2.19	0.42
1:B:114:ILE:HD13	1:B:176:LEU:HD13	2.00	0.42
1:A:31:ILE:HD12	1:A:63:THR:HB	2.02	0.42
1:A:28:ARG:HH22	2:C:33:GLU:H	1.67	0.42
1:A:117:ILE:HD11	1:A:145:ALA:CB	2.50	0.42
2:D:15:ILE:HD13	2:D:42:VAL:HG21	2.02	0.42
1:B:129:TYR:CE1	1:B:133:LEU:HD21	2.54	0.42
1:B:19:LEU:HD13	1:B:41:TYR:CE1	2.54	0.42
1:B:152:ILE:HG22	1:B:160:LEU:HD11	2.01	0.42
1:B:28:ARG:HE	1:B:28:ARG:HA	1.83	0.42
2:D:99:ARG:NH1	5:D:202:HOH:O	2.45	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:TYR:HB3	1:B:79:GLU:HB3	2.01	0.42
1:A:50:SER:HB2	2:C:103:ARG:HG2	2.02	0.42
1:B:153:ALA:HA	1:B:160:LEU:HD13	2.02	0.41
1:B:48:LYS:HD2	1:B:51:GLU:HB3	2.03	0.40
1:B:28:ARG:C	1:B:28:ARG:HE	2.23	0.40
1:A:19:LEU:HG	1:A:86:ILE:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/181 (92%)	166 (99%)	1 (1%)	0	100	100
1	B	162/181 (90%)	161 (99%)	1 (1%)	0	100	100
2	C	124/127 (98%)	123 (99%)	1 (1%)	0	100	100
2	D	123/127 (97%)	122 (99%)	1 (1%)	0	100	100
All	All	576/616 (94%)	572 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/139 (92%)	125 (98%)	3 (2%)	50	55
1	B	129/139 (93%)	126 (98%)	3 (2%)	50	55
2	C	108/105 (103%)	106 (98%)	2 (2%)	57	63
2	D	106/105 (101%)	103 (97%)	3 (3%)	43	47
All	All	471/488 (96%)	460 (98%)	11 (2%)	50	55

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	LEU
1	A	132	GLN
1	A	177	ARG
1	B	28	ARG
1	B	36	HIS
1	B	48	LYS
2	C	20	GLN
2	C	119	PHE
2	D	23	ARG
2	D	102	GLU
2	D	119	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TPO	D	68	2,4	8,10,11	0.90	0	10,14,16	1.74	2 (20%)
2	TPO	C	68	2,4	8,10,11	0.91	0	10,14,16	1.60	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	D	68	2,4	-	1/9/11/13	-
2	TPO	C	68	2,4	-	2/9/11/13	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	68	TPO	P-OG1-CB	-4.46	109.73	123.21
2	C	68	TPO	P-OG1-CB	-4.46	109.74	123.21
2	D	68	TPO	O-C-CA	-2.16	119.12	124.78

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	68	TPO	O-C-CA-CB
2	C	68	TPO	O-C-CA-CB
2	C	68	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ADP	A	1001	4	24,29,29	1.32	3 (12%)	29,45,45	1.50	5 (17%)
3	ADP	B	2001	4	24,29,29	1.19	3 (12%)	29,45,45	1.56	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1001	4	-	3/12/32/32	0/3/3/3
3	ADP	B	2001	4	-	4/12/32/32	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	ADP	C2'-C1'	-3.04	1.49	1.53
3	A	1001	ADP	PB-O3B	-2.43	1.45	1.54
3	B	2001	ADP	C5-N7	-2.20	1.31	1.39
3	B	2001	ADP	PB-O2B	-2.12	1.46	1.54
3	B	2001	ADP	C5-C4	2.10	1.46	1.40
3	A	1001	ADP	C5-C4	2.08	1.46	1.40

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	ADP	O2'-C2'-C1'	-3.36	98.44	110.85
3	B	2001	ADP	C2-N1-C6	2.93	123.77	118.75
3	B	2001	ADP	C3'-C2'-C1'	2.92	105.38	100.98
3	B	2001	ADP	N3-C2-N1	-2.89	124.16	128.68
3	B	2001	ADP	C4-C5-N7	-2.83	106.45	109.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	ADP	N3-C2-N1	-2.81	124.28	128.68
3	A	1001	ADP	C4-C5-N7	-2.47	106.82	109.40
3	A	1001	ADP	O5'-C5'-C4'	2.37	117.14	108.99
3	A	1001	ADP	C2-N1-C6	2.36	122.80	118.75
3	B	2001	ADP	O3B-PB-O1B	2.24	119.45	110.68
3	B	2001	ADP	C5-C6-N1	-2.18	115.42	120.35
3	B	2001	ADP	O3A-PB-O1B	-2.11	99.48	111.19

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	2001	ADP	C5'-O5'-PA-O1A
3	B	2001	ADP	C5'-O5'-PA-O3A
3	B	2001	ADP	C5'-O5'-PA-O2A
3	A	1001	ADP	PB-O3A-PA-O1A
3	A	1001	ADP	PB-O3A-PA-O2A
3	B	2001	ADP	PB-O3A-PA-O1A
3	A	1001	ADP	C5'-O5'-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	167/181 (92%)	0.93	26 (15%) <b>2</b> <b>2</b>	19, 41, 78, 111	0
1	B	164/181 (90%)	1.03	32 (19%) <b>1</b> <b>1</b>	19, 44, 73, 90	0
2	C	122/127 (96%)	0.54	4 (3%) 46 53	23, 37, 58, 69	0
2	D	121/127 (95%)	0.43	4 (3%) 46 53	26, 36, 56, 80	0
All	All	574/616 (93%)	0.77	66 (11%) <b>4</b> <b>6</b>	19, 39, 69, 111	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLU	7.7
1	A	80	ALA	5.0
1	A	170	GLY	4.6
1	B	133	LEU	4.4
1	B	80	ALA	4.2
1	A	3	THR	4.2
1	B	144	GLY	4.1
1	B	75	TYR	3.7
1	B	19	LEU	3.6
1	A	21[A]	CYS	3.5
1	B	118	ALA	3.4
1	B	120	LEU	3.4
1	B	125	LEU	3.3
1	B	90	GLU	3.3
1	A	146	GLY	3.2
1	A	117	ILE	3.1
1	A	144	GLY	3.1
1	A	81	ALA	3.1
1	B	34	ILE	3.1
1	B	86	ILE	3.0
1	A	168	PRO	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	20	LEU	3.0
1	B	124	ALA	2.9
1	B	119	ASN	2.9
2	D	52	ASP	2.9
1	A	22	PHE	2.8
1	B	72	LEU	2.8
1	B	31	ILE	2.8
2	C	37	GLU	2.7
1	B	84	VAL	2.7
1	B	22	PHE	2.7
1	A	105	ASP	2.6
1	B	105	ASP	2.6
1	A	75	TYR	2.6
1	A	19	LEU	2.6
2	D	115	CYS	2.5
2	C	36	TYR	2.5
1	A	30	LEU	2.4
1	A	4	LEU	2.4
1	A	20	LEU	2.4
2	C	0	SER	2.4
2	D	121	ILE	2.4
1	B	171	ARG	2.4
1	B	145	ALA	2.4
1	A	116	ALA	2.4
1	B	83	THR	2.4
1	B	21	CYS	2.4
2	C	51	ALA	2.4
1	A	104	ARG	2.4
1	B	117	ILE	2.3
1	A	78	HIS	2.3
1	B	112	ARG	2.3
1	A	18	ILE	2.3
1	A	121	ASP	2.2
1	B	159	PRO	2.2
1	A	112	ARG	2.2
1	B	116	ALA	2.2
1	A	145	ALA	2.2
1	B	26	ILE	2.1
1	A	34	ILE	2.1
2	D	93	LEU	2.1
1	B	146	GLY	2.1
1	B	121	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	83	THR	2.0
1	B	134	ARG	2.0
1	B	74	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	TPO	D	68	11/12	0.98	0.12	25,33,37,40	0
2	TPO	C	68	11/12	0.98	0.10	24,27,40,42	0

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ADP	B	2001	27/27	0.74	0.24	37,65,76,80	0
4	NA	B	2002	1/1	0.75	0.13	74,74,74,74	0
4	NA	C	1101	1/1	0.78	0.15	53,53,53,53	0
3	ADP	A	1001	27/27	0.81	0.24	31,66,71,73	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around NA B 2002:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.