



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2022 – 07:10 PM EST

PDB ID : 2KLQ
Title : The solution structure of CBD of human MCM6
Authors : Liu, C.
Deposited on : 2009-07-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.27
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	114	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:717-A:789, A:810-A:817 (81)	0.50	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 7 single-model clusters were found.

Cluster number	Models
1	4, 5, 9, 12, 13, 19
2	7, 10, 15
3	11, 17
4	8, 14
Single-model clusters	1; 2; 3; 6; 16; 18; 20

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1866 atoms, of which 928 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	114	1866	592	928	155	191	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	721	SER	CYS	engineered mutation	UNP Q14566

4 Residue-property plots i

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA replication licensing factor MCM6




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA replication licensing factor MCM6




4.2.2 Score per residue for model 2

- Molecule 1: DNA replication licensing factor MCM6




4.2.3 Score per residue for model 3

- Molecule 1: DNA replication licensing factor MCM6




4.2.4 Score per residue for model 4

- Molecule 1: DNA replication licensing factor MCM6




4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: DNA replication licensing factor MCM6




4.2.6 Score per residue for model 6

- Molecule 1: DNA replication licensing factor MCM6




4.2.7 Score per residue for model 7

- Molecule 1: DNA replication licensing factor MCM6




4.2.8 Score per residue for model 8

- Molecule 1: DNA replication licensing factor MCM6




4.2.9 Score per residue for model 9

- Molecule 1: DNA replication licensing factor MCM6



4.2.10 Score per residue for model 10

- Molecule 1: DNA replication licensing factor MCM6



4.2.11 Score per residue for model 11

- Molecule 1: DNA replication licensing factor MCM6





4.2.12 Score per residue for model 12

- Molecule 1: DNA replication licensing factor MCM6



4.2.13 Score per residue for model 13

- Molecule 1: DNA replication licensing factor MCM6



4.2.14 Score per residue for model 14

- Molecule 1: DNA replication licensing factor MCM6



4.2.15 Score per residue for model 15

- Molecule 1: DNA replication licensing factor MCM6




4.2.16 Score per residue for model 16

- Molecule 1: DNA replication licensing factor MCM6




4.2.17 Score per residue for model 17

- Molecule 1: DNA replication licensing factor MCM6




4.2.18 Score per residue for model 18

- Molecule 1: DNA replication licensing factor MCM6



4.2.19 Score per residue for model 19

- Molecule 1: DNA replication licensing factor MCM6



4.2.20 Score per residue for model 20

- Molecule 1: DNA replication licensing factor MCM6



A708	F709	K710	A711	S712	L713	R714	L715	G716	F717	S718	R722	L726	L731	M749	M750	M751	Y752	L753	K754	T781	H782	L787	L788	E789	L790	I791	Q792	A793	G794	L795	K796	G797	S798	I799	E800	G801	S802	E803	S804	Y805	E806	E807	D808	F809	V812	P815	M816	L817	L818	L819	E820
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

D821

5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *QUALITY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.2±0.4
All	All	0	4

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	722	ARG	Sidechain	3
1	A	745	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	695	693	693	6±2
All	All	13900	13860	13860	120

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:726:LEU:HD12	1:A:752:TYR:CD1	0.63	2.28	20	2
1:A:749:VAL:O	1:A:753:LEU:HG	0.63	1.94	8	20
1:A:780:LEU:CD2	1:A:787:LEU:HB2	0.61	2.26	13	1
1:A:779:ARG:HA	1:A:783:TYR:CD1	0.59	2.33	3	5
1:A:756:ILE:O	1:A:759:GLU:HG2	0.56	2.01	15	7
1:A:742:ALA:O	1:A:810:TYR:HA	0.54	2.02	8	1
1:A:756:ILE:O	1:A:760:ILE:HG12	0.53	2.03	11	9
1:A:788:ILE:HB	1:A:812:VAL:HG13	0.52	1.79	5	7
1:A:752:TYR:O	1:A:756:ILE:HG12	0.51	2.05	17	5
1:A:753:LEU:O	1:A:757:GLU:HG3	0.51	2.06	7	4
1:A:750:ASN:O	1:A:754:LYS:HD3	0.50	2.07	11	6
1:A:750:ASN:O	1:A:754:LYS:HG2	0.49	2.06	9	2
1:A:728:VAL:O	1:A:732:ARG:HG3	0.49	2.07	8	1
1:A:728:VAL:O	1:A:732:ARG:HG2	0.49	2.07	5	2
1:A:754:LYS:HA	1:A:757:GLU:OE1	0.49	2.08	12	1
1:A:769:LYS:O	1:A:773:ILE:HG22	0.49	2.07	15	1
1:A:787:LEU:HD12	1:A:812:VAL:O	0.49	2.08	20	1
1:A:781:THR:HG22	1:A:787:LEU:HD22	0.48	1.84	4	2
1:A:734:VAL:HG13	1:A:741:SER:HB2	0.48	1.85	8	1
1:A:726:LEU:HD12	1:A:752:TYR:CG	0.48	2.43	17	1
1:A:722:ARG:O	1:A:725:ASN:HB3	0.48	2.09	13	1
1:A:786:VAL:HG11	1:A:817:TYR:CD1	0.47	2.44	17	1
1:A:752:TYR:O	1:A:756:ILE:HG13	0.46	2.11	1	1
1:A:770:LYS:O	1:A:774:GLU:HG2	0.46	2.10	6	1
1:A:781:THR:HG22	1:A:787:LEU:HD13	0.46	1.86	19	1
1:A:760:ILE:HG21	1:A:766:LEU:HB2	0.46	1.87	8	1
1:A:768:ASN:O	1:A:772:ILE:HG12	0.45	2.10	12	4
1:A:731:LEU:HD21	1:A:815:PRO:HA	0.45	1.88	20	1
1:A:734:VAL:HG11	1:A:741:SER:HA	0.45	1.88	11	1
1:A:723:ILE:CG2	1:A:776:VAL:HG21	0.45	2.41	17	1
1:A:718:SER:O	1:A:722:ARG:HG2	0.45	2.12	17	4
1:A:788:ILE:HD11	1:A:814:ASN:ND2	0.44	2.27	4	1
1:A:788:ILE:HB	1:A:812:VAL:HB	0.44	1.88	18	1
1:A:770:LYS:HA	1:A:773:ILE:HG22	0.44	1.89	3	1
1:A:729:LEU:O	1:A:732:ARG:HG2	0.44	2.12	2	1
1:A:780:LEU:HG	1:A:787:LEU:HB2	0.44	1.89	9	1
1:A:745:ARG:HD3	1:A:746:SER:N	0.44	2.27	8	1
1:A:723:ILE:O	1:A:726:LEU:HB3	0.44	2.12	17	1
1:A:718:SER:O	1:A:722:ARG:HG3	0.44	2.11	20	1
1:A:730:HIS:O	1:A:733:LYS:HG2	0.43	2.13	6	1
1:A:786:VAL:HG11	1:A:817:TYR:HB2	0.43	1.89	11	1
1:A:766:LEU:O	1:A:770:LYS:HG3	0.42	2.14	15	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:763:GLU:O	1:A:767:ILE:HG13	0.42	2.15	3	1
1:A:781:THR:HA	1:A:787:LEU:CD1	0.42	2.44	11	1
1:A:767:ILE:O	1:A:771:ARG:HG3	0.42	2.15	2	1
1:A:726:LEU:HD23	1:A:727:ILE:N	0.42	2.30	16	2
1:A:781:THR:HG23	1:A:782:HIS:CD2	0.41	2.50	20	1
1:A:744:LYS:O	1:A:748:LEU:HG	0.41	2.14	7	1
1:A:776:VAL:O	1:A:780:LEU:HB2	0.41	2.15	13	1
1:A:765:GLU:O	1:A:769:LYS:HB2	0.41	2.15	15	1
1:A:723:ILE:HG21	1:A:776:VAL:HG21	0.41	1.93	17	1
1:A:780:LEU:CD2	1:A:787:LEU:HB3	0.41	2.46	17	1
1:A:731:LEU:O	1:A:735:GLU:HG2	0.40	2.15	4	1
1:A:733:LYS:O	1:A:737:GLU:HG2	0.40	2.17	16	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/114 (71%)	76±2 (94±3%)	5±2 (6±3%)	0±1 (1±1%)	29	74
All	All	1620/2280 (71%)	1518 (94%)	92 (6%)	10 (1%)	29	74

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	742	ALA	3
1	A	816	ASN	2
1	A	739	ASP	2
1	A	815	PRO	2
1	A	810	TYR	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/106 (75%)	77±1 (96±1%)	3±1 (4±1%)	36	84
All	All	1600/2120 (75%)	1537 (96%)	63 (4%)	36	84

All 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	726	LEU	19
1	A	812	VAL	15
1	A	719	GLU	4
1	A	787	LEU	3
1	A	789	GLU	3
1	A	754	LYS	2
1	A	722	ARG	2
1	A	745	ARG	2
1	A	769	LYS	2
1	A	817	TYR	2
1	A	765	GLU	1
1	A	770	LYS	1
1	A	814	ASN	1
1	A	783	TYR	1
1	A	816	ASN	1
1	A	717	PHE	1
1	A	731	LEU	1
1	A	810	TYR	1
1	A	735	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided