



Full wwPDB X-ray Structure Validation Report i

Aug 28, 2023 – 03:40 AM EDT

PDB ID : 3KL5
Title : Structure Analysis of a Xylanase From Glycosyl Hydrolase Family Thirty: Carbohydrate Ligand Complexes Reveal this Family of Enzymes Unique Mechanism of Substrate Specificity and Recognition
Authors : St John, F.J.; Hurlbert, J.C.; Pozharski, E.
Deposited on : 2009-11-06
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.


We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35

Continued from previous page...



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYP	H	1	-	-	-	X

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	10	Total O 10 10	0	0
3	B	17	Total O 17 17	0	0
3	C	19	Total O 19 19	0	0
3	D	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucuronoxylanase xynC



- Molecule 1: Glucuronoxylanase xynC



- Molecule 1: Glucuronoxylanase xynC



- Molecule 1: Glucuronoxylanase xynC

XYP1
XYP2
GCY3

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	137.72Å 194.01Å 65.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.59 44.67 – 2.59	Depositor EDS
% Data completeness (in resolution range)	97.5 (50.00-2.59) 97.6 (44.67-2.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.32 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R , R_{free}	0.241 , 0.290 0.237 , 0.284	Depositor DCC
R_{free} test set	2778 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	65.9	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.3	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12477	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Mol	Chain	Res	Type
1	A	58	GLU
1	A	60	ARG
1	A	178	TYR
1	A	209	SER
1	A	213	TYR
1	A	230	VAL
1	A	324	LYS
1	A	335	THR
1	A	337	VAL
1	A	369	THR
1	B	58	GLU
1	B	60	ARG
1	B	178	TYR
1	B	209	SER
1	B	230	VAL
1	B	251	GLN
1	B	324	LYS
1	B	335	THR
1	B	337	VAL
1	B	353[A]	ARG
1	B	353[B]	ARG
1	B	369	THR
1	B	392	LEU
1	C	58	GLU
1	C	60	ARG
1	C	89	SER
1	C	178	TYR
1	C	209	SER
1	C	213	TYR
1	C	230	VAL
1	C	324	LYS
1	C	335	THR
1	C	337	VAL
1	C	369	THR
1	D	46	GLN
1	D	60	ARG
1	D	71	LYS
1	D	76	HIS
1	D	123	THR
1	D	149	TRP
1	D	153	GLN
1	D	167	ASN
1	D	169	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	178	TYR
1	D	213	TYR
1	D	230	VAL
1	D	241	ASP
1	D	245	GLU
1	D	261	ASP
1	D	307	THR
1	D	309	ASN
1	D	337	VAL
1	D	363	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	345	ASN
1	C	162	ASN
1	C	345	ASN
1	D	24	HIS
1	D	61	ASN
1	D	76	HIS
1	D	145	HIS
1	D	153	GLN
1	D	309	ASN
1	D	313	ASN
1	D	339	GLN
1	D	363	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

12 monosaccharides are modelled in this entry.

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	XYP	O2-C2-C3	3.23	116.62	110.14
2	G	3	GCV	O4-C4-C3	-3.10	102.87	110.29
2	H	3	GCV	C4-C5-C6	-2.93	108.11	112.15
2	F	1	XYP	O1-C1-O5	2.32	115.77	109.72
2	G	1	XYP	C5-O5-C1	2.32	115.09	111.52
2	G	3	GCV	C4-C5-C6	2.22	115.22	112.15
2	E	2	XYP	C5-O5-C1	2.20	114.91	111.52
2	E	1	XYP	C5-C4-C3	-2.08	107.10	109.67
2	G	3	GCV	C1-C2-C3	2.05	112.19	109.67

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	3	GCV	O5-C5-C6-O6B


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1	XYP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '*#RSRZ > 2*' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	389/401 (97%)	0.41	25 (6%)	19  14 	42, 58, 72, 80	0
1	B	390/401 (97%)	0.42	6 (1%)	73  70 	42, 58, 73, 79	1 (0%)
1	C	389/401 (97%)	0.49	8 (2%)	63  58 	42, 58, 72, 79	0
1	D	367/401 (91%)	2.75	215 (58%)	0  0 	74, 120, 155, 179	364 (99%)
All	All	1535/1604 (95%)	0.99	254 (16%)	1  1 	42, 62, 140, 179	365 (23%)

All (254) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	102	SER	10.0
1	D	343	LEU	9.6
1	D	100	ASP	9.2
1	D	103	ALA	8.9
1	D	319	TYR	8.0
1	D	337	VAL	7.8
1	D	38	THR	7.7
1	D	11	ALA	7.7
1	D	387	PHE	7.7
1	D	302	VAL	7.6
1	D	306	ALA	7.5
1	D	183	SER	7.5
1	D	101	THR	7.4
1	D	95	PHE	7.1
1	D	307	THR	6.9
1	D	17	ARG	6.9
1	D	355	ILE	6.9
1	D	8	ASN	6.8
1	D	108	TYR	6.7
1	D	192	ALA	6.4
1	D	389	VAL	6.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	345	ASN	6.4
1	D	304	ILE	6.3
1	D	112	ALA	6.2
1	D	363	GLN	6.2
1	D	280	ASP	6.1
1	D	354	TRP	6.0
1	D	33	ALA	5.8
1	D	149	TRP	5.7
1	D	388	VAL	5.7
1	D	341	PHE	5.6
1	D	106	LEU	5.5
1	D	366	THR	5.5
1	D	7	VAL	5.4
1	D	61	ASN	5.4
1	D	27	TRP	5.3
1	D	91	MET	5.3
1	D	68	GLU	5.3
1	D	325	VAL	5.2
1	D	157	ARG	5.2
1	D	143	TYR	5.1
1	D	140	GLU	5.1
1	D	37	GLU	5.1
1	D	339	GLN	5.0
1	D	9	VAL	5.0
1	D	109	ASN	4.9
1	D	148	THR	4.9
1	D	190	PRO	4.9
1	D	318	ALA	4.9
1	D	67	VAL	4.8
1	D	146	GLU	4.8
1	D	342	VAL	4.8
1	D	43	GLY	4.8
1	D	336	GLY	4.7
1	D	386	THR	4.7
1	D	105	ARG	4.6
1	D	10	SER	4.6
1	D	4	ASP	4.6
1	D	281	GLY	4.6
1	D	375	PHE	4.5
1	D	322	ASP	4.5
1	D	376	TRP	4.5
1	D	311	ASN	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	3	SER	4.5
1	D	310	PRO	4.4
1	D	83	SER	4.4
1	A	104	LYS	4.4
1	D	113	ALA	4.4
1	D	94	THR	4.4
1	D	182	LEU	4.3
1	D	107	LYS	4.3
1	D	358	SER	4.3
1	D	5	VAL	4.3
1	D	135	ILE	4.2
1	D	116	GLN	4.2
1	D	115	ALA	4.2
1	A	391	ARG	4.2
1	D	32	THR	4.2
1	D	156	LEU	4.1
1	D	62	ASN	4.1
1	D	124	PHE	4.1
1	D	320	LYS	4.1
1	D	128	ASN	4.1
1	A	95	PHE	4.1
1	D	364	PRO	4.0
1	D	298	ARG	4.0
1	D	222	GLY	4.0
1	D	141	PRO	4.0
1	D	89	SER	4.0
1	D	64	TYR	4.0
1	D	42	ASN	4.0
1	D	305	ASP	4.0
1	D	104	LYS	3.9
1	D	391	ARG	3.9
1	D	203	LEU	3.9
1	D	47	LEU	3.9
1	D	76	HIS	3.9
1	D	36	ARG	3.9
1	D	59	ASN	3.8
1	D	221	ALA	3.8
1	D	380	PRO	3.8
1	A	145	HIS	3.8
1	D	147	TRP	3.8
1	D	159	MET	3.8
1	D	120	ASP	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	209	SER	3.6
1	D	297	VAL	3.6
1	D	153	GLN	3.6
1	A	103	ALA	3.6
1	D	338	ASN	3.6
1	A	101	THR	3.6
1	D	145	HIS	3.5
1	D	142	ASP	3.5
1	D	212	PRO	3.5
1	D	177	GLN	3.5
1	D	288	TYR	3.5
1	D	198	ILE	3.5
1	D	326	VAL	3.5
1	D	131	ASN	3.5
1	D	362	LEU	3.4
1	D	84	PRO	3.4
1	D	85	TRP	3.4
1	D	110	LYS	3.4
1	D	390	ASN	3.4
1	A	274	TYR	3.4
1	A	96	ASN	3.4
1	D	46	GLN	3.4
1	D	88	PRO	3.4
1	A	143	TYR	3.3
1	D	35	GLN	3.3
1	D	172	ALA	3.3
1	D	16	ILE	3.3
1	D	12	GLU	3.3
1	D	30	ASP	3.3
1	D	44	GLN	3.3
1	D	365	GLY	3.2
1	D	290	MET	3.2
1	D	152	PRO	3.2
1	D	117	HIS	3.2
1	D	151	THR	3.2
1	D	150	TRP	3.2
1	D	344	GLN	3.2
1	D	330	ILE	3.2
1	D	170	VAL	3.2
1	D	163	ALA	3.1
1	D	317	SER	3.1
1	D	138	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	171	ILE	3.1
1	D	186	ILE	3.1
1	A	99	GLY	3.1
1	D	200	GLY	3.1
1	A	344	GLN	3.1
1	D	60	ARG	3.1
1	D	176	PHE	3.1
1	D	28	ALA	3.1
1	D	31	LEU	3.1
1	D	258	VAL	3.1
1	B	193	LEU	3.1
1	D	79	ILE	3.0
1	A	144	ALA	3.0
1	D	168	ALA	3.0
1	D	296	PHE	3.0
1	A	98	ASN	3.0
1	D	111	TYR	3.0
1	D	213	TYR	3.0
1	D	87	PRO	3.0
1	D	121	PHE	2.9
1	D	218	GLN	2.9
1	D	41	GLY	2.9
1	D	15	VAL	2.9
1	D	316	VAL	2.9
1	D	196	MET	2.9
1	D	321	GLY	2.8
1	D	217	LYS	2.8
1	C	348	ALA	2.8
1	D	374	HIS	2.8
1	D	324	LYS	2.8
1	D	243	TRP	2.8
1	D	118	LEU	2.8
1	D	45	ASN	2.8
1	D	189	ASP	2.8
1	D	332	LYS	2.7
1	D	49	PHE	2.7
1	A	237	THR	2.7
1	D	327	ILE	2.7
1	D	216	PHE	2.7
1	D	314	VAL	2.6
1	D	194	ALA	2.6
1	D	224	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	201	THR	2.6
1	D	93	GLU	2.6
1	D	367	ASN	2.6
1	B	218	GLN	2.6
1	D	160	ARG	2.6
1	D	40	PHE	2.6
1	D	166	ILE	2.6
1	D	382	GLN	2.6
1	D	86	ASN	2.6
1	A	348	ALA	2.5
1	A	102	SER	2.5
1	A	149	TRP	2.5
1	A	373	ASN	2.5
1	D	381	ALA	2.5
1	D	34	ALA	2.5
1	D	237	THR	2.5
1	D	191	GLN	2.4
1	A	273	SER	2.4
1	A	106	LEU	2.4
1	D	328	VAL	2.4
1	D	301	TYR	2.4
1	C	373	ASN	2.4
1	D	71	LYS	2.4
1	B	3	SER	2.4
1	D	57	ASP	2.4
1	D	72	SER	2.4
1	A	157	ARG	2.4
1	D	139	ASN	2.3
1	C	226	TRP	2.3
1	D	90	ASP	2.3
1	B	312	ALA	2.3
1	D	48	GLY	2.3
1	D	309	ASN	2.3
1	C	15	VAL	2.3
1	D	187	LEU	2.3
1	D	129	GLY	2.2
1	D	127	ASN	2.2
1	D	383	SER	2.2
1	D	313	ASN	2.2
1	C	124	PHE	2.2
1	A	100	ASP	2.2
1	B	362	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	300	GLY	2.2
1	D	50	SER	2.2
1	D	58	GLU	2.2
1	C	265	TYR	2.1
1	A	127	ASN	2.1
1	D	299	PRO	2.1
1	D	225	LEU	2.1
1	A	358	SER	2.1
1	D	126	LYS	2.1
1	D	377	ALA	2.1
1	D	378	HIS	2.1
1	D	340	ASN	2.1
1	D	236	ASP	2.1
1	D	75	LYS	2.1
1	B	355	ILE	2.1
1	D	158	PHE	2.1
1	C	99	GLY	2.1
1	D	161	GLU	2.1
1	C	22	MET	2.0
1	D	14	GLN	2.0
1	D	21	GLY	2.0
1	D	262	PHE	2.0
1	D	205	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYP	H	1	10/10	0.39	0.41	118,118,118,119	0
2	XYP	H	2	9/10	0.55	0.38	115,116,117,117	0
2	XYP	G	2	9/10	0.79	0.17	88,90,91,91	0
2	XYP	E	2	9/10	0.80	0.23	92,93,94,95	0
2	XYP	F	1	10/10	0.83	0.20	80,82,83,83	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GCV	G	3	13/14	0.85	0.19	86,88,89,90	0
2	XYP	E	1	10/10	0.87	0.35	92,94,94,95	0
2	GCV	H	3	13/14	0.87	0.28	112,114,115,115	0
2	GCV	E	3	13/14	0.88	0.30	91,94,95,95	0
2	XYP	G	1	9/10	0.90	0.17	91,92,92,93	0
2	XYP	F	2	9/10	0.94	0.10	76,77,78,78	0
2	GCV	F	3	13/14	0.94	0.17	73,76,78,79	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain F:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
 $mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
 $mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.