

wwPDB X-ray Structure Validation Summary Report (i)

Oct 17, 2021 – 08:48 AM EDT

PDB ID	:	1KM1
Title	:	Orotidine monophosphate decarboxylase mutant S127A crystal structure
Authors	:	Wu, N.; Gillon, W.; Pai, E.F.
Deposited on	:	2001-12-13
Resolution	:	1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	3665(1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	247	78%	8%	14%
1	В	247	80%	5%	15%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3634 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	212	Total	С	Ν	0	\mathbf{S}	0	9	0
	A	212	1637	1030	291	305	11	0	Э	0
1	В	211	Total	С	Ν	0	S	0	2	0
	D	211	1618	1018	286	303	11		2	0

• Molecule 1 is a protein called OROTIDINE 5'-PHOSPHATE DECARBOXYLASE.

Chain	Residue	Modelled	Actual	Comment	Reference
А	101	PRO	ARG	engineered mutation	UNP O26232
А	127	ALA	SER	engineered mutation	UNP O26232
А	227	ILE	ASN	cloning artifact	UNP O26232
А	229	GLU	-	cloning artifact	UNP O26232
А	230	ASP	-	cloning artifact	UNP O26232
А	231	PRO	-	cloning artifact	UNP O26232
А	232	ALA	-	cloning artifact	UNP O26232
А	233	ALA	-	cloning artifact	UNP O26232
А	234	ASN	-	cloning artifact	UNP O26232
А	235	LYS	-	cloning artifact	UNP O26232
А	236	ALA	-	cloning artifact	UNP O26232
А	237	ARG	-	cloning artifact	UNP O26232
А	238	LYS	-	cloning artifact	UNP O26232
А	239	GLU	-	cloning artifact	UNP O26232
А	240	ALA	-	cloning artifact	UNP O26232
А	241	GLU	-	cloning artifact	UNP O26232
А	242	LEU	-	cloning artifact	UNP O26232
А	243	ALA	-	cloning artifact	UNP O26232
А	244	ALA	-	cloning artifact	UNP O26232
А	245	ALA	-	cloning artifact	UNP O26232
А	246	THR	-	cloning artifact	UNP O26232
А	247	ALA	-	cloning artifact	UNP O26232
В	101	PRO	ARG	engineered mutation	UNP O26232
В	127	ALA	SER	engineered mutation	UNP O26232
В	227	ILE	ASN	cloning artifact	UNP O26232


There are 44 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
В	229	GLU	-	cloning artifact	UNP O26232
В	230	ASP	-	cloning artifact	UNP O26232
В	231	PRO	-	cloning artifact	UNP O26232
В	232	ALA	-	cloning artifact	UNP O26232
В	233	ALA	-	cloning artifact	UNP O26232
В	234	ASN	-	cloning artifact	UNP O26232
В	235	LYS	-	cloning artifact	UNP O26232
В	236	ALA	-	cloning artifact	UNP O26232
В	237	ARG	-	cloning artifact	UNP O26232
В	238	LYS	-	cloning artifact	UNP O26232
В	239	GLU	-	cloning artifact	UNP O26232
В	240	ALA	-	cloning artifact	UNP O26232
В	241	GLU	-	cloning artifact	UNP O26232
В	242	LEU	-	cloning artifact	UNP O26232
В	243	ALA	-	cloning artifact	UNP O26232
В	244	ALA	-	cloning artifact	UNP O26232
В	245	ALA	-	cloning artifact	UNP O26232
В	246	THR	- cloning artifact		UNP O26232
В	247	ALA	-	cloning artifact	UNP O26232

Continued from previous page...

• Molecule 2 is 6-AZA URIDINE 5'-MONOPHOSPHATE (three-letter code: UP6) (formula: $C_8H_{12}N_3O_9P$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Λ	1	Total	С	Ν	Ο	Р	0	1
	Л	1	29	11	6	11	1	0	T



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	В	1	Total	С	Ν	0	Р	0	1
_	D	-	29	11	6	11	1	Ŭ	-

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	163	Total O 163 163	0	0
3	В	158	Total O 158 158	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: OROTIDINE 5'-PHOSPHATE DECARBOXYLASE



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	58.34Å 73.75Å 59.40Å	Depositor	
a, b, c, α , β , γ	90.00° 119.41° 90.00°	Depositor	
Resolution (Å)	29.85 - 1.60	Depositor	
% Data completeness	95.1 (29.85-1.60)	Depositor	
(in resolution range)	55.1 (25.05-1.00)	Depositor	
R_{merge}	(Not available)	Depositor	
R _{sym}	(Not available)	Depositor	
Refinement program	CNS	Depositor	
R, R_{free}	0.177 , 0.191	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3634	wwPDB-VP	
Average B, all atoms $(Å^2)$	17.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $\rm UP6$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.42	0/1661	0.65	0/2241	
1	В	0.44	0/1642	0.65	0/2216	
All	All	0.43	0/3303	0.65	0/4457	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1637	0	1652	18	0
1	В	1618	0	1629	11	0
2	А	29	0	6	0	0
2	В	29	0	6	0	0
3	А	163	0	0	3	0
3	В	158	0	0	1	0
All	All	3634	0	3293	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their



Atom-1	Atom-2	Interatomic	Clash
	1100m –	distance $(Å)$	overlap (Å)
1:B:216:ALA:O	1:B:220:GLU:HG3	1.86	0.75
1:A:144[A]:ARG:HH21	1:A:166:ARG:HH22	1.36	0.74
1:B:92:ALA:O	1:B:118:ARG:NH1	2.26	0.68
1:A:92:ALA:O	1:A:118:ARG:NH1	2.29	0.65
1:B:38:ILE:O	1:B:38:ILE:HD13	1.97	0.63

clash magnitude.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	213/247~(86%)	210 (99%)	3~(1%)	0	100	100
1	В	211/247~(85%)	208 (99%)	3 (1%)	0	100	100
All	All	424/494 (86%)	418 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	168/192~(88%)	166~(99%)	2(1%)	71 54
1	В	166/192~(86%)	165~(99%)	1 (1%)	86 77



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	334/384~(87%)	331~(99%)	3~(1%)	78 65

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	38	ILE
1	А	157	PRO
1	В	38	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
	Type	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	UP6	В	5002[A]	-	22,22,22	2.67	5 (22%)	32,33,33	<mark>3.23</mark>	8 (25%)



Mol	Turne	Chain	Res	Res Link B		ond leng	$_{\rm ths}$	В	ond ang	les
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	UP6	А	5001[B]	-	22,22,22	2.41	3 (13%)	32,33,33	<mark>3.18</mark>	7 (21%)
2	UP6	В	5002[B]	-	22,22,22	2.44	4 (18%)	32,33,33	3.16	6 (18%)
2	UP6	А	5001[A]	-	22,22,22	2.65	4 (18%)	32,33,33	3.22	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UP6	В	5002[A]	-	-	4/10/26/26	0/1/2/2
2	UP6	А	5001[B]	-	-	1/10/26/26	0/1/2/2
2	UP6	В	5002[B]	-	-	1/10/26/26	0/1/2/2
2	UP6	A	5001[A]	-	-	4/10/26/26	0/1/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	А	5001[A]	UP6	C5-C4	-9.88	1.43	1.50
2	В	5002[A]	UP6	C5-C4	-9.72	1.43	1.50
2	В	5002[B]	UP6	C5-C4	-9.40	1.43	1.50
2	А	5001[B]	UP6	C5-C4	-9.01	1.44	1.50
2	В	5002[A]	UP6	C5-N6	-4.94	1.31	1.46

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	5002[B]	UP6	O4-C4-C5	10.99	127.27	118.67
2	А	5001[B]	UP6	O4-C4-C5	10.97	127.26	118.67
2	А	5001[A]	UP6	O4-C4-C5	10.22	126.67	118.67
2	В	5002[A]	UP6	C5-N6-N1	10.19	123.27	110.27
2	А	5001[A]	UP6	C5-N6-N1	10.13	123.19	110.27

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	5001[A]	UP6	C2'-C1'-N1-N6
2	В	5002[A]	UP6	C2'-C1'-N1-N6
2	А	5001[A]	UP6	O4'-C1'-N1-C2




Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	В	5002[A]	UP6	O4'-C1'-N1-C2
2	А	5001[A]	UP6	O4'-C4'-C5'-O5'


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient must be highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

