



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 03:46 AM JST

PDB ID : 6KMV
Title : caspase-11 C254A P22/P10 in complex with mouse GSDMD-C domain
Authors : Ding, J.; Sun, Q.
Deposited on : 2019-08-01
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1558 (3.42-3.30)
Ramachandran outliers	138981	1599 (3.42-3.30)
Sidechain outliers	138945	1598 (3.42-3.30)
RSRZ outliers	127900	1507 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	99% .
2	B	167	99% .
2	V	167	10% 99% .
3	C	198	3% 100% .
3	G	198	2% 100% .
3	O	198	4% 97% .
4	D	197	5% 99% .

















Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	K	197	16% 99%
4	S	197	10% 100%
4	T	197	6% 100%
4	W	197	8% 100%
4	b	197	17% 91% 9%
4	f	197	12% 100%
5	E	166	% 99%
5	F	166	% 99%
5	Z	166	3% 99%
5	d	166	2% 99%
6	H	191	17% 89% 10%
7	I	165	100%
7	M	165	% 99%
7	N	165	2% 99%
7	Q	165	% 100%
7	Y	165	2% 99%
8	J	149	% 99%
9	L	214	5% 99%
9	P	214	99%
9	e	214	2% 99%
10	R	150	100%
10	c	150	% 100%
11	U	165	4% 99%
12	X	199	100%
13	a	196	14% 93% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
14	g	88	 99%
14	h	88	 98%
14	i	88	 97%
14	j	88	 99%
14	k	88	 98%
14	l	88	 99%
14	m	88	 98%
14	n	88	 99%
14	o	88	 98%
14	p	88	 99%
14	q	88	 99%
14	r	88	 99%
14	t	88	 98%
14	u	88	 98%
14	v	88	 99%
15	s	87	 99%

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 56064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	161	1258	788	217	242	11	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 2 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	167	1316	825	226	253	12	0	0	0
2	V	167	1316	825	226	253	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	254	ALA	CYS	engineered mutation	UNP P70343
V	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 3 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	198	1506	960	237	301	8	0	0	0
3	G	198	1506	960	237	301	8	0	0	0
3	O	198	1506	960	237	301	8	0	0	0

- Molecule 4 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			
4	K	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			
4	S	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			
4	T	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			
4	W	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			
4	b	180	Total	C	N	O	S	0	0	0
			1363	870	211	275	7			
4	f	197	Total	C	N	O	S	0	0	0
			1501	957	236	300	8			

- Molecule 5 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	166	Total	C	N	O	S	0	0	0
			1310	822	225	251	12			
5	F	166	Total	C	N	O	S	0	0	0
			1310	822	225	251	12			
5	Z	166	Total	C	N	O	S	0	0	0
			1310	822	225	251	12			
5	d	166	Total	C	N	O	S	0	0	0
			1310	822	225	251	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	254	ALA	CYS	engineered mutation	UNP P70343
F	254	ALA	CYS	engineered mutation	UNP P70343
Z	254	ALA	CYS	engineered mutation	UNP P70343
d	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 6 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	H	171	Total	C	N	O	S	0	0	0
			1304	830	206	261	7			

- Molecule 7 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	I	165	Total 1303	C 818	N 224	O 249	S 12	0	0	0
7	M	165	Total 1303	C 818	N 224	O 249	S 12	0	0	0
7	N	165	Total 1303	C 818	N 224	O 249	S 12	0	0	0
7	Q	165	Total 1303	C 818	N 224	O 249	S 12	0	0	0
7	Y	165	Total 1303	C 818	N 224	O 249	S 12	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	254	ALA	CYS	engineered mutation	UNP P70343
M	254	ALA	CYS	engineered mutation	UNP P70343
N	254	ALA	CYS	engineered mutation	UNP P70343
Q	254	ALA	CYS	engineered mutation	UNP P70343
Y	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 8 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	J	149	Total 1169	C 734	N 200	O 225	S 10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 9 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	L	214	Total 1627	C 1035	N 253	O 331	S 8	0	0	0
9	P	214	Total 1627	C 1035	N 253	O 331	S 8	0	0	0
9	e	214	Total 1627	C 1035	N 253	O 331	S 8	0	0	0

- Molecule 10 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	R	150	1175	737	201	227	10	0	0	0
10	c	150	1175	737	201	227	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	254	ALA	CYS	engineered mutation	UNP P70343
c	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 11 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	U	165	1301	817	224	248	12	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	254	ALA	CYS	engineered mutation	UNP P70343

- Molecule 12 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	X	199	1511	963	238	302	8	0	0	0

- Molecule 13 is a protein called Gasdermin-D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	a	183	1405	898	222	278	7	0	0	0

- Molecule 14 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	g	88	726	471	125	126	4	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	h	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	i	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	j	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	k	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	l	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	m	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	n	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	o	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	p	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	q	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	r	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	t	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	u	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			
14	v	88	Total	C	N	O	S	0	0	0
			726	471	125	126	4			

- Molecule 15 is a protein called Caspase-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	s	87	Total	C	N	O	S	0	0	0
			721	468	124	125	4			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Caspase-4

Chain A:  99%




- Molecule 2: Caspase-4

Chain B:  99%




- Molecule 2: Caspase-4

Chain V:  10% 99%




- Molecule 3: Gasdermin-D

Chain C:  3% 100%




- Molecule 3: Gasdermin-D


Chain G:  2% 100%




- Molecule 3: Gasdermin-D




● Molecule 4: Gasdermin-D




● Molecule 4: Gasdermin-D




● Molecule 4: Gasdermin-D





● Molecule 4: Gasdermin-D




● Molecule 4: Gasdermin-D




● Molecule 4: Gasdermin-D







• Molecule 4: Gasdermin-D




• Molecule 5: Caspase-4




• Molecule 5: Caspase-4






• Molecule 5: Caspase-4



• Molecule 5: Caspase-4



• Molecule 6: Gasdermin-D




- Molecule 7: Caspase-4

Chain I:  100%

There are no outlier residues recorded for this chain.

- Molecule 7: Caspase-4

Chain M:  99%



- Molecule 7: Caspase-4

Chain N:  99%



- Molecule 7: Caspase-4

Chain Q:  100%



- Molecule 7: Caspase-4

Chain Y:  99%




- Molecule 8: Caspase-4

Chain J:  99%



- Molecule 9: Gasdermin-D

Chain L:  99%




- Molecule 9: Gasdermin-D

Chain P:  99%



- Molecule 9: Gasdermin-D

Chain e:  99%



- Molecule 10: Caspase-4

Chain R:  100%

There are no outlier residues recorded for this chain.


- Molecule 10: Caspase-4

Chain c:  100%



- Molecule 11: Caspase-4

Chain U:  99%




- Molecule 12: Gasdermin-D

Chain X:  100%


There are no outlier residues recorded for this chain.

- Molecule 13: Gasdermin-D

Chain a:  93%



● Molecule 14: Caspase-4



● Molecule 14: Caspase-4



● Molecule 14: Caspase-4




● Molecule 14: Caspase-4



● Molecule 14: Caspase-4



● Molecule 14: Caspase-4



● Molecule 14: Caspase-4




- Molecule 14: Caspase-4

Chain n:  99%




- Molecule 14: Caspase-4

Chain o:  98%



- Molecule 14: Caspase-4

Chain p:  99%




- Molecule 14: Caspase-4

Chain q:  99%




- Molecule 14: Caspase-4

Chain r:  99%




- Molecule 14: Caspase-4

Chain t:  98%




- Molecule 14: Caspase-4

Chain u:  98%



- Molecule 14: Caspase-4



- Molecule 15: Caspase-4



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	89.80Å 139.21Å 175.94Å 92.43° 99.06° 96.35°	Depositor
Resolution (Å)	37.44 – 3.35 45.07 – 3.35	Depositor EDS
% Data completeness (in resolution range)	98.2 (37.44-3.35) 94.2 (45.07-3.35)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 3.32Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.219 , 0.252 0.220 , 0.252	Depositor DCC
R_{free} test set	2000 reflections (1.69%)	wwPDB-VP
Wilson B-factor (Å ²)	81.1	Xtrriage
Anisotropy	0.167	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	56064	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1280	0.50	0/1724
2	B	0.24	0/1339	0.43	0/1802
2	V	0.24	0/1339	0.43	0/1802
3	C	0.23	0/1532	0.40	0/2089
3	G	0.24	0/1532	0.39	0/2089
3	O	0.29	0/1532	0.46	0/2089
4	D	0.24	0/1527	0.39	0/2082
4	K	0.23	0/1527	0.39	0/2082
4	S	0.24	0/1527	0.39	0/2082
4	T	0.23	0/1527	0.39	0/2082
4	W	0.23	0/1527	0.39	0/2082
4	b	0.24	0/1382	0.39	0/1877
4	f	0.24	0/1527	0.39	0/2082
5	E	0.24	0/1333	0.43	0/1794
5	F	0.24	0/1333	0.43	0/1794
5	Z	0.24	0/1333	0.43	0/1794
5	d	0.24	0/1333	0.43	0/1794
6	H	0.23	0/1324	0.40	0/1801
7	I	0.24	0/1326	0.42	0/1784
7	M	0.24	0/1326	0.42	0/1784
7	N	0.24	0/1326	0.42	0/1784
7	Q	0.24	0/1326	0.42	0/1784
7	Y	0.24	0/1326	0.42	0/1784
8	J	0.24	0/1190	0.43	0/1603
9	L	0.24	0/1653	0.40	0/2253
9	P	0.31	0/1653	0.46	0/2253
9	e	0.24	0/1653	0.40	0/2253
10	R	0.24	0/1196	0.43	0/1611
10	c	0.24	0/1196	0.43	0/1611
11	U	0.24	0/1324	0.43	0/1782
12	X	0.25	0/1537	0.41	0/2096
13	a	0.24	0/1429	0.39	0/1945
14	g	0.24	0/748	0.44	0/1007
14	h	0.24	0/748	0.43	0/1007

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
14	i	0.24	0/748	0.44	0/1007
14	j	0.24	0/748	0.42	0/1007
14	k	0.24	0/748	0.44	0/1007
14	l	0.24	0/748	0.42	0/1007
14	m	0.24	0/748	0.43	0/1007
14	n	0.24	0/748	0.42	0/1007
14	o	0.24	0/748	0.44	0/1007
14	p	0.24	0/748	0.43	0/1007
14	q	0.24	0/748	0.44	0/1007
14	r	0.24	0/748	0.42	0/1007
14	t	0.24	0/748	0.42	0/1007
14	u	0.24	0/748	0.44	0/1007
14	v	0.25	0/748	0.42	0/1007
15	s	0.24	0/743	0.44	0/1000
All	All	0.24	0/57178	0.42	0/77373

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/161 (99%)	151 (95%)	8 (5%)	0	100	100
2	B	165/167 (99%)	157 (95%)	8 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	V	165/167 (99%)	156 (94%)	8 (5%)	1 (1%)	25	59
3	C	196/198 (99%)	195 (100%)	1 (0%)	0	100	100
3	G	196/198 (99%)	195 (100%)	1 (0%)	0	100	100
3	O	196/198 (99%)	193 (98%)	2 (1%)	1 (0%)	29	63
4	D	195/197 (99%)	193 (99%)	2 (1%)	0	100	100
4	K	195/197 (99%)	194 (100%)	1 (0%)	0	100	100
4	S	195/197 (99%)	194 (100%)	1 (0%)	0	100	100
4	T	195/197 (99%)	194 (100%)	1 (0%)	0	100	100
4	W	195/197 (99%)	194 (100%)	1 (0%)	0	100	100
4	b	170/197 (86%)	169 (99%)	1 (1%)	0	100	100
4	f	195/197 (99%)	194 (100%)	1 (0%)	0	100	100
5	E	164/166 (99%)	156 (95%)	8 (5%)	0	100	100
5	F	164/166 (99%)	157 (96%)	7 (4%)	0	100	100
5	Z	164/166 (99%)	155 (94%)	9 (6%)	0	100	100
5	d	164/166 (99%)	156 (95%)	8 (5%)	0	100	100
6	H	163/191 (85%)	163 (100%)	0	0	100	100
7	I	163/165 (99%)	155 (95%)	8 (5%)	0	100	100
7	M	163/165 (99%)	154 (94%)	9 (6%)	0	100	100
7	N	163/165 (99%)	155 (95%)	7 (4%)	1 (1%)	25	59
7	Q	163/165 (99%)	155 (95%)	8 (5%)	0	100	100
7	Y	163/165 (99%)	154 (94%)	9 (6%)	0	100	100
8	J	147/149 (99%)	141 (96%)	6 (4%)	0	100	100
9	L	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
9	P	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
9	e	212/214 (99%)	206 (97%)	5 (2%)	1 (0%)	29	63
10	R	148/150 (99%)	142 (96%)	6 (4%)	0	100	100
10	c	148/150 (99%)	140 (95%)	8 (5%)	0	100	100
11	U	163/165 (99%)	155 (95%)	8 (5%)	0	100	100
12	X	197/199 (99%)	196 (100%)	1 (0%)	0	100	100
13	a	179/196 (91%)	178 (99%)	1 (1%)	0	100	100
14	g	86/88 (98%)	83 (96%)	3 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
14	h	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	i	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	j	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	k	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	l	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	m	86/88 (98%)	82 (95%)	4 (5%)	0	100	100
14	n	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	o	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	p	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	q	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	r	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	t	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	u	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
14	v	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
15	s	85/87 (98%)	82 (96%)	3 (4%)	0	100	100
All	All	7044/7206 (98%)	6838 (97%)	202 (3%)	4 (0%)	51	82

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
9	e	282	GLU
2	V	265	ARG
3	O	398	THR
7	N	128	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/139 (100%)	137 (99%)	2 (1%)	67	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	146/146 (100%)	144 (99%)	2 (1%)	67	83
2	V	146/146 (100%)	145 (99%)	1 (1%)	84	92
3	C	171/171 (100%)	171 (100%)	0	100	100
3	G	171/171 (100%)	171 (100%)	0	100	100
3	O	171/171 (100%)	167 (98%)	4 (2%)	50	75
4	D	171/171 (100%)	170 (99%)	1 (1%)	86	93
4	K	171/171 (100%)	169 (99%)	2 (1%)	71	85
4	S	171/171 (100%)	171 (100%)	0	100	100
4	T	171/171 (100%)	171 (100%)	0	100	100
4	W	171/171 (100%)	171 (100%)	0	100	100
4	b	155/171 (91%)	155 (100%)	0	100	100
4	f	171/171 (100%)	171 (100%)	0	100	100
5	E	145/145 (100%)	144 (99%)	1 (1%)	84	92
5	F	145/145 (100%)	144 (99%)	1 (1%)	84	92
5	Z	145/145 (100%)	144 (99%)	1 (1%)	84	92
5	d	145/145 (100%)	144 (99%)	1 (1%)	84	92
6	H	150/166 (90%)	149 (99%)	1 (1%)	84	92
7	I	144/144 (100%)	144 (100%)	0	100	100
7	M	144/144 (100%)	143 (99%)	1 (1%)	84	92
7	N	144/144 (100%)	143 (99%)	1 (1%)	84	92
7	Q	144/144 (100%)	144 (100%)	0	100	100
7	Y	144/144 (100%)	143 (99%)	1 (1%)	84	92
8	J	128/128 (100%)	127 (99%)	1 (1%)	81	91
9	L	185/185 (100%)	182 (98%)	3 (2%)	62	81
9	P	185/185 (100%)	182 (98%)	3 (2%)	62	81
9	e	185/185 (100%)	183 (99%)	2 (1%)	73	86
10	R	129/129 (100%)	129 (100%)	0	100	100
10	c	129/129 (100%)	129 (100%)	0	100	100
11	U	144/144 (100%)	142 (99%)	2 (1%)	67	83
12	X	171/171 (100%)	171 (100%)	0	100	100
13	a	158/170 (93%)	157 (99%)	1 (1%)	86	93

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
14	g	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	h	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	i	80/80 (100%)	77 (96%)	3 (4%)	33	63
14	j	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	k	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	l	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	m	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	n	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	o	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	p	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	q	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	r	80/80 (100%)	79 (99%)	1 (1%)	69	84
14	t	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	u	80/80 (100%)	78 (98%)	2 (2%)	47	73
14	v	80/80 (100%)	79 (99%)	1 (1%)	69	84
15	s	80/80 (100%)	79 (99%)	1 (1%)	69	84
All	All	6269/6313 (99%)	6213 (99%)	56 (1%)	78	89

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	a	407	LEU
14	v	299	PHE
14	i	299	PHE
14	u	321	ARG
14	r	299	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
14	r	344	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/161 (100%)	-0.05	0 100 100	41, 65, 104, 154	0
2	B	167/167 (100%)	-0.01	0 100 100	41, 63, 95, 109	0
2	V	167/167 (100%)	0.48	17 (10%) 6 8	70, 116, 140, 159	0
3	C	198/198 (100%)	0.27	6 (3%) 50 53	60, 88, 125, 139	0
3	G	198/198 (100%)	0.29	3 (1%) 73 76	70, 97, 121, 137	0
3	O	198/198 (100%)	0.18	8 (4%) 38 40	30, 94, 119, 140	0
4	D	197/197 (100%)	0.30	9 (4%) 32 35	66, 98, 140, 149	0
4	K	197/197 (100%)	0.85	31 (15%) 2 2	77, 122, 147, 154	0
4	S	197/197 (100%)	0.57	20 (10%) 6 8	68, 107, 148, 162	0
4	T	197/197 (100%)	0.48	12 (6%) 21 23	81, 116, 150, 160	0
4	W	197/197 (100%)	0.64	16 (8%) 12 13	99, 126, 146, 153	0
4	b	180/197 (91%)	0.89	34 (18%) 1 1	96, 123, 145, 153	0
4	f	197/197 (100%)	0.69	23 (11%) 4 5	80, 111, 150, 163	0
5	E	166/166 (100%)	-0.03	2 (1%) 79 82	48, 75, 111, 129	0
5	F	166/166 (100%)	-0.07	2 (1%) 79 82	53, 77, 111, 120	0
5	Z	166/166 (100%)	0.17	5 (3%) 50 53	68, 93, 127, 147	0
5	d	166/166 (100%)	-0.08	3 (1%) 68 71	41, 64, 96, 123	0
6	H	171/191 (89%)	1.02	32 (18%) 1 1	86, 126, 156, 165	0
7	I	165/165 (100%)	-0.04	0 100 100	45, 66, 105, 120	0
7	M	165/165 (100%)	0.09	1 (0%) 89 92	61, 91, 117, 132	0
7	N	165/165 (100%)	0.09	3 (1%) 68 71	72, 100, 130, 138	0
7	Q	165/165 (100%)	0.09	1 (0%) 89 92	56, 82, 109, 123	0
7	Y	165/165 (100%)	0.09	3 (1%) 68 71	66, 94, 129, 144	0
8	J	149/149 (100%)	0.06	2 (1%) 77 80	44, 64, 88, 125	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
9	L	214/214 (100%)	0.30	10 (4%) 31 34	64, 100, 128, 138	0
9	P	214/214 (100%)	0.11	0 100 100	30, 77, 97, 114	0
9	e	214/214 (100%)	0.19	5 (2%) 60 63	51, 84, 121, 138	0
10	R	150/150 (100%)	-0.12	0 100 100	45, 61, 100, 118	0
10	c	150/150 (100%)	-0.01	1 (0%) 87 91	45, 74, 95, 136	0
11	U	165/165 (100%)	0.19	7 (4%) 36 38	66, 87, 121, 130	0
12	X	199/199 (100%)	0.12	0 100 100	30, 73, 97, 109	0
13	a	183/196 (93%)	0.84	28 (15%) 2 2	86, 120, 142, 148	0
14	g	88/88 (100%)	0.08	1 (1%) 80 84	43, 59, 86, 104	0
14	h	88/88 (100%)	-0.01	0 100 100	41, 58, 95, 118	0
14	i	88/88 (100%)	-0.02	0 100 100	52, 69, 93, 104	0
14	j	88/88 (100%)	0.01	0 100 100	50, 72, 92, 102	0
14	k	88/88 (100%)	0.07	0 100 100	44, 66, 87, 94	0
14	l	88/88 (100%)	0.07	1 (1%) 80 84	43, 64, 94, 109	0
14	m	88/88 (100%)	0.11	1 (1%) 80 84	67, 87, 110, 119	0
14	n	88/88 (100%)	0.12	0 100 100	69, 90, 124, 130	0
14	o	88/88 (100%)	-0.02	0 100 100	42, 68, 96, 105	0
14	p	88/88 (100%)	-0.00	1 (1%) 80 84	44, 61, 82, 112	0
14	q	88/88 (100%)	0.26	1 (1%) 80 84	69, 95, 114, 128	0
14	r	88/88 (100%)	0.36	5 (5%) 23 26	78, 107, 136, 144	0
14	t	88/88 (100%)	0.40	6 (6%) 17 20	71, 93, 122, 132	0
14	u	88/88 (100%)	0.11	1 (1%) 80 84	43, 63, 100, 105	0
14	v	88/88 (100%)	0.14	2 (2%) 60 63	43, 64, 99, 130	0
15	s	87/87 (100%)	0.08	1 (1%) 80 84	67, 88, 112, 128	0
All	All	7156/7206 (99%)	0.25	304 (4%) 36 38	30, 89, 137, 165	0

The worst 5 of 304 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
13	a	396	LEU	6.7
4	f	431	LEU	6.2
13	a	441	THR	6.0
4	K	450	LEU	6.0
4	b	398	THR	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.