

Full wwPDB X-ray Structure Validation Report (i)

Sep 20, 2023 – 12:19 PM EDT

PDB ID	:	5KNL
Title	:	Crystal structure of S. pombe ubiquitin E1 (Uba1) in complex with Ubc15 and
		ubiquitin
Authors	:	Olsen, S.K.; Lv, Z.; Yuan, L.; Williams, K.
Deposited on	:	2016-06-28
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	4.02b-467
:	1.8.5 (274361), CSD as541be (2020)
:	1.13
:	2.35.1
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	5.8.0158
:	7.0.044 (Gargrove)
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.35.1
	:::::::::::::::::::::::::::::::::::::::

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	$5231 \ (2.50-2.50)$
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	1001	95%	•••
1	D	1001	95%	•••
2	В	96	74% • •	21%
2	Е	96	^{2%} 78% •	21%
3	С	175	<u>5%</u> 91%	6% ••



Mol	Chain	Length	Quality of chain	
	1		5%	
3	\mathbf{F}	175	89%	•• 7%



5KNL

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 38561 atoms, of which 19138 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ubiquitin-activating enzyme E1 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	А	981	Total 15342	C 4026	H 7637	N 1250	0 1481	S 30	0	0	0
1		077	Total	4920 C	H	1259 N	0	<u> </u>	0	0	0
	I D	977	15280	4907	7607	1254	1473	39	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	12	SER	-	expression tag	UNP O94609
D	12	SER	-	expression tag	UNP O94609

• Molecule 2 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	В	76	Total 1236	C 375	Н 623	N 119	0 118	S 1	0	0	0
2	Е	76	Total 1236	C 375	Н 623	N 119	0 118	S 1	0	0	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-19	MET	-	initiating methionine	UNP A0A081CFF0
В	-18	GLY	-	expression tag	UNP A0A081CFF0
В	-17	SER	-	expression tag	UNP A0A081CFF0
В	-16	SER	-	expression tag	UNP A0A081CFF0
В	-15	HIS	-	expression tag	UNP A0A081CFF0
В	-14	HIS	-	expression tag	UNP A0A081CFF0
В	-13	HIS	-	expression tag	UNP A0A081CFF0
В	-12	HIS	-	expression tag	UNP A0A081CFF0
В	-11	HIS	-	expression tag	UNP A0A081CFF0
B	-10	HIS	-	expression tag	UNP A0A081CFF0



Chain	Residue	Modelled	Actual	Comment	Reference
В	-9	SER	-	expression tag	UNP A0A081CFF0
В	-8	SER	-	expression tag	UNP A0A081CFF0
В	-7	GLY	-	expression tag	UNP A0A081CFF0
В	-6	LEU	-	expression tag	UNP A0A081CFF0
В	-5	VAL	-	expression tag	UNP A0A081CFF0
В	-4	PRO	-	expression tag	UNP A0A081CFF0
В	-3	ARG	-	expression tag	UNP A0A081CFF0
В	-2	GLY	-	expression tag	UNP A0A081CFF0
В	-1	SER	-	expression tag	UNP A0A081CFF0
В	0	HIS	-	expression tag	UNP A0A081CFF0
В	6	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	11	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	27	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	29	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	33	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	48	ARG	LYS	engineered mutation	UNP A0A081CFF0
В	57	ALA	SER	engineered mutation	UNP A0A081CFF0
В	63	ARG	LYS	engineered mutation	UNP A0A081CFF0
E	-19	MET	-	initiating methionine	UNP A0A081CFF0
Ε	-18	GLY	-	expression tag	UNP A0A081CFF0
E	-17	SER	-	expression tag	UNP A0A081CFF0
E	-16	SER	-	expression tag	UNP A0A081CFF0
E	-15	HIS	-	expression tag	UNP A0A081CFF0
E	-14	HIS	-	expression tag	UNP A0A081CFF0
E	-13	HIS	-	expression tag	UNP A0A081CFF0
E	-12	HIS	-	expression tag	UNP A0A081CFF0
E	-11	HIS	-	expression tag	UNP A0A081CFF0
E	-10	HIS	-	expression tag	UNP A0A081CFF0
E	-9	SER	-	expression tag	UNP A0A081CFF0
E	-8	SER	-	expression tag	UNP A0A081CFF0
E	-7	GLY	-	expression tag	UNP A0A081CFF0
E	-6	LEU	-	expression tag	UNP A0A081CFF0
E	-5	VAL	-	expression tag	UNP A0A081CFF0
E	-4	PRO	-	expression tag	UNP A0A081CFF0
E	-3	ARG	-	expression tag	UNP A0A081CFF0
E	-2	GLY	-	expression tag	UNP A0A081CFF0
E	-1	SER	-	expression tag	UNP A0A081CFF0
E	0	HIS	-	expression tag	UNP A0A081CFF0
E	6	ARG	LYS	engineered mutation	UNP A0A081CFF0
E	11	ARG	LYS	engineered mutation	UNP A0A081CFF0
E	27	ARG	LYS	engineered mutation	UNP A0A081CFF0
E	29	ARG	LYS	engineered mutation	UNP A0A081CFF0



Chain	Residue	Modelled	Actual	Comment	Reference
Ε	33	ARG	LYS	engineered mutation	UNP A0A081CFF0
Е	48	ARG	LYS	engineered mutation	UNP A0A081CFF0
Е	57	ALA	SER	engineered mutation	UNP A0A081CFF0
Е	63	ARG	LYS	engineered mutation	UNP A0A081CFF0

• Molecule 3 is a protein called Ubiquitin-conjugating enzyme E2 15.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
3	3 C	171	Total	С	Н	Ν	0	S	0	Ο	0
0			2718	877	1346	230	258	7	0	0	0
2	3 F	162	Total	С	Η	Ν	0	S	0	1	0
0			2610	840	1302	222	239	7	0	L	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	168	LEU	-	expression tag	UNP Q9Y818
С	169	GLU	-	expression tag	UNP Q9Y818
С	170	HIS	-	expression tag	UNP Q9Y818
С	171	HIS	-	expression tag	UNP Q9Y818
С	172	HIS	-	expression tag	UNP Q9Y818
С	173	HIS	-	expression tag	UNP Q9Y818
С	174	HIS	-	expression tag	UNP Q9Y818
С	175	HIS	-	expression tag	UNP Q9Y818
F	168	LEU	-	expression tag	UNP Q9Y818
F	169	GLU	-	expression tag	UNP Q9Y818
F	170	HIS	-	expression tag	UNP Q9Y818
F	171	HIS	-	expression tag	UNP Q9Y818
F	172	HIS	-	expression tag	UNP Q9Y818
F	173	HIS	-	expression tag	UNP Q9Y818
F	174	HIS	-	expression tag	UNP Q9Y818
F	175	HIS	-	expression tag	UNP Q9Y818

• Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	F	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	32	$\begin{array}{cc} \text{Total} & \text{O} \\ 32 & 32 \end{array}$	0	0
5	В	7	Total O 7 7	0	0
5	С	6	Total O 6 6	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	46	Total O 46 46	0	0
5	Ε	2	Total O 2 2	0	0
5	F	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Ubiquitin-activating enzyme E1 1





Chain E:	78%	• 21%
MET GLY SER SER HIS HIS HIS HIS SER SER SER SER SER VAL VAL	PR0 ARG GLY BER MI MI MI M1 M1 C M1 C C C C C C C	
• Molecule 3: Ubiqu	itin-conjugating enzyme E2 15	
Chain C:	91%	6% ••
M1 K12 H41 P97 P97 P99 D100 D100 M100 R110	D133 E134 8135 8135 8135 8135 8135 8135 8145 81471 81471 81471 81473 81473 81473 81473 81473 81473 81473 81473 81474 81474 81476 814566 81456 81456 81456 814566 814566 814566 814566 8146	
• Molecule 3: Ubiqu	itin-conjugating enzyme E2 15	
Chain F:	89%	•• 7%
M1 F25 D32 D32 ASP ASP ASP TYR GU7	TYR GLU ASP ALA ASP ALA TIA B110 B115 B115 B115 P136 P136 P136 P136 P136 P136 P136 P136	SIH



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	77.11Å 82.24Å 135.38Å	Depositor
a, b, c, α , β , γ	102.10° 95.78° 90.86°	Depositor
Bosolution(A)	28.31 - 2.50	Depositor
Resolution (A)	28.41 - 2.50	EDS
% Data completeness	96.9 (28.31-2.50)	Depositor
(in resolution range)	96.9 (28.41-2.50)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.08 (at 2.51 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
B B.	0.216 , 0.251	Depositor
n, n_{free}	0.219 , 0.252	DCC
R_{free} test set	10829 reflections (10.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	29.5	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 35.8	EDS
L-test for $twinning^2$	$ < L >=0.44, < L^2>=0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	38561	wwPDB-VP
Average B, all atoms $(Å^2)$	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.22	0/7876	0.37	0/10660
1	D	0.23	0/7844	0.37	0/10618
2	В	0.22	0/618	0.43	0/831
2	Е	0.22	0/618	0.40	0/831
3	С	0.23	0/1411	0.40	0/1915
3	F	0.24	0/1348	0.46	1/1829~(0.1%)
All	All	0.23	0/19715	0.39	1/26684~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	F	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	F	135	SER	C-N-CD	-5.85	107.73	120.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	F	135	SER	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	7705	7637	7634	10	0
1	D	7673	7607	7607	11	0
2	В	613	623	625	3	0
2	Е	613	623	625	1	0
3	С	1372	1346	1341	5	0
3	F	1308	1302	1298	4	0
4	А	20	0	0	0	0
4	В	5	0	0	1	0
4	D	15	0	0	0	0
4	F	5	0	0	0	0
5	А	32	0	0	0	0
5	В	7	0	0	0	0
5	С	6	0	0	0	0
5	D	46	0	0	0	0
5	E	2	0	0	1	0
5	F	1	0	0	0	0
All	All	19423	19138	19130	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:684:GLN:O	1:D:688:ASN:ND2	2.32	0.61
2:B:33:ARG:NH1	4:B:101:SO4:O3	2.36	0.59
1:D:1009:CYS:SG	1:D:1011:LYS:NZ	2.68	0.57
1:D:13:ASN:OD1	1:D:14:THR:N	2.37	0.57
1:A:997:ASP:O	1:A:999:ASN:N	2.40	0.55
1:D:997:ASP:O	1:D:999:ASN:N	2.40	0.54
1:D:225:THR:O	1:D:227:ARG:NH1	2.41	0.54
1:A:684:GLN:O	1:A:688:ASN:ND2	2.41	0.51
3:F:135:SER:HB3	3:F:136:PRO:HA	1.94	0.49
1:D:994:CYS:SG	3:F:12:LYS:NZ	2.81	0.48
1:A:785:GLN:O	1:A:787:LEU:N	2.45	0.48



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
3:C:98:GLY:O	3:C:110:ARG:NH1	2.47	0.48
1:A:783:ASP:OD1	1:A:783:ASP:N	2.47	0.47
1:D:957:GLN:NE2	3:F:32:ASP:OD2	2.48	0.47
3:F:25:PHE:HZ	3:F:125:ILE:HG12	1.81	0.46
2:B:27:ARG:NH2	2:B:50:LEU:O	2.48	0.45
1:A:994:CYS:SG	3:C:12:LYS:NZ	2.81	0.45
1:D:881:GLU:O	1:D:884:LYS:NZ	2.50	0.44
2:E:6:ARG:NH1	5:E:101:HOH:O	2.46	0.44
3:C:169:GLU:O	3:C:171:HIS:N	2.50	0.44
1:A:645:ARG:HG3	1:A:788:LYS:CE	2.48	0.44
1:D:995:CYS:O	1:D:1003:VAL:N	2.47	0.43
1:A:476:PHE:O	1:A:479:ARG:NH1	2.49	0.42
2:B:74:ARG:HD3	2:B:75:GLY:N	2.34	0.41
3:C:100:ASP:HB3	3:C:101:LYS:HA	2.02	0.41
1:A:959:PRO:HA	1:A:962:LEU:HB2	2.02	0.41
3:C:97:PRO:HA	3:C:98:GLY:HA2	1.87	0.41
1:A:174:ASP:O	1:A:377:PHE:N	2.42	0.40
1:D:957:GLN:HB2	1:D:962:LEU:HD11	2.03	0.40
1:A:881:GLU:O	1:A:884:LYS:NZ	2.55	0.40
1:D:205:GLU:N	1:D:208:ASP:OD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	977/1001~(98%)	923 (94%)	52 (5%)	2(0%)	47	68
1	D	973/1001~(97%)	923~(95%)	49 (5%)	1 (0%)	51	73
2	В	74/96~(77%)	71 (96%)	3 (4%)	0	100	100
2	Е	74/96~(77%)	71 (96%)	3 (4%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
3	С	169/175~(97%)	160 (95%)	6 (4%)	3 (2%)	8	14
3	F	159/175~(91%)	152 (96%)	5(3%)	2(1%)	12	21
All	All	2426/2544~(95%)	2300~(95%)	118 (5%)	8 (0%)	41	61

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	170	HIS
1	А	786	GLU
1	А	998	ALA
3	С	170	HIS
1	D	998	ALA
3	F	135	SER
3	С	99	ASP
3	С	98	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	860/877~(98%)	853~(99%)	7 (1%)	81	93
1	D	857/877~(98%)	853 (100%)	4 (0%)	88	96
2	В	67/84~(80%)	66~(98%)	1 (2%)	65	85
2	Е	67/84~(80%)	67 (100%)	0	100	100
3	С	154/160~(96%)	153~(99%)	1 (1%)	86	95
3	F	149/160~(93%)	149 (100%)	0	100	100
All	All	2154/2242~(96%)	2141 (99%)	13 (1%)	86	95

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	80	GLN
1	А	218	MET



Mol	Chain	Res	Type
1	А	464	MET
1	А	691	LYS
1	А	841	ARG
1	А	999	ASN
1	А	1002	ASP
2	В	74	ARG
3	С	41	MET
1	D	80	GLN
1	D	999	ASN
1	D	1002	ASP
1	D	1006	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Turne	Chain	Dec	Tink	B	ond leng	gths	B	ond ang	gles
IVIOI	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	SO4	А	1102	-	4,4,4	0.15	0	$6,\!6,\!6$	0.05	0
4	SO4	D	1102	-	4,4,4	0.14	0	$6,\!6,\!6$	0.05	0
4	SO4	В	101	-	4,4,4	0.14	0	$6,\!6,\!6$	0.05	0
4	SO4	А	1101	-	4,4,4	0.15	0	$6,\!6,\!6$	0.06	0
4	SO4	А	1104	-	4,4,4	0.14	0	$6,\!6,\!6$	0.06	0
4	SO4	D	1103	-	4,4,4	0.14	0	$6,\!6,\!6$	0.06	0
4	SO4	D	1101	-	4,4,4	0.13	0	$6,\!6,\!6$	0.07	0
4	SO4	F	201	-	4,4,4	0.13	0	$6,\!6,\!6$	0.05	0
4	SO4	А	1103	-	4,4,4	0.14	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	101	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ >	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	981/1001 (98%)	0.35	68 (6%) 16 17	11, 44, 103, 165	0
1	D	977/1001~(97%)	0.40	75 (7%) 13 13	12, 43, 111, 179	0
2	В	76/96~(79%)	0.05	0 100 100	22, 45, 88, 108	0
2	Е	76/96~(79%)	0.35	2 (2%) 56 59	21, 61, 104, 146	0
3	С	171/175~(97%)	0.28	9 (5%) 26 28	22, 49, 98, 140	0
3	F	162/175~(92%)	0.41	8 (4%) 29 31	30, 57, 102, 146	1 (0%)
All	All	2443/2544 (96%)	0.36	162 (6%) 18 19	11, 45, 104, 179	1 (0%)

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	199	GLU	8.2
1	D	998	ALA	7.6
1	А	1000	GLY	7.2
1	А	198	GLU	6.2
1	D	911	LYS	6.0
1	D	250	TYR	6.0
1	D	187	SER	5.7
1	А	998	ALA	5.5
1	D	228	LYS	5.3
1	D	201	ARG	5.2
1	D	909	ASN	5.2
1	D	235	TYR	4.9
1	А	911	LYS	4.8
1	D	198	GLU	4.8
1	D	638	LEU	4.7
1	A	908	VAL	4.6
1	А	199	GLU	4.6
1	D	245	LEU	4.5
1	А	999	ASN	4.4



Mol	Chain	Res	Type	RSRZ
1	D	910	GLY	4.4
1	D	640	THR	4.3
1	А	1001	GLU	4.3
1	D	1000	GLY	4.2
1	D	999	ASN	4.2
1	А	638	LEU	4.2
1	D	936	LYS	4.2
1	А	782	LYS	4.2
1	D	232	LYS	4.1
1	D	251	ASN	4.1
1	А	907	LYS	4.0
1	А	209	PHE	4.0
1	D	244	ASP	4.0
1	А	909	ASN	4.0
1	D	639	LYS	3.9
1	А	783	ASP	3.9
1	А	784	LYS	3.8
1	А	230	GLU	3.8
1	А	935	GLN	3.7
1	D	762	LYS	3.7
1	D	745	ALA	3.7
1	D	1001	GLU	3.6
2	Ε	17	VAL	3.6
1	А	232	LYS	3.6
1	А	220	GLY	3.6
1	А	742	THR	3.5
3	F	109	GLU	3.5
1	D	802	GLY	3.5
1	А	244	ASP	3.4
1	D	935	GLN	3.4
1	D	741	GLU	3.4
1	А	407	GLY	3.3
1	A	14	THR	3.3
1	A	243	LYS	3.3
1	D	239	ILE	3.3
1	D	335	ASP	3.3
1	D	963	ALA	3.3
1	D	907	LYS	3.2
1	A	936	LYS	3.2
1	D	908	VAL	3.2
1	А	223	ASP	3.1
1	А	987	LYS	3.1



Mol	Chain	Res	Type	RSRZ
1	А	997	ASP	3.1
1	D	200	THR	3.1
3	С	150	GLN	3.1
1	D	219	PRO	3.1
1	D	696	SER	3.1
1	А	749	ARG	3.1
1	D	185	ILE	3.0
1	А	228	LYS	3.0
1	А	192	GLY	3.0
1	А	191	ASP	3.0
2	Е	32	ASP	3.0
1	A	404	LYS	3.0
1	D	912	GLU	2.9
1	A	758	PRO	2.9
1	А	789	SER	2.9
1	D	242	VAL	2.9
3	С	171	HIS	2.9
1	А	635	GLU	2.8
1	А	719	ARG	2.8
1	А	239	ILE	2.8
1	А	963	ALA	2.8
1	А	241	SER	2.8
1	D	236	THR	2.8
1	D	697	THR	2.8
1	D	184	MET	2.7
3	С	135	SER	2.7
1	D	197	LEU	2.7
1	А	785	GLN	2.7
1	А	980	LYS	2.7
1	D	247	SER	2.7
1	A	937	GLU	2.7
1	D	243	LYS	2.7
3	F	25	PHE	2.7
1	A	910	GLY	2.6
1	D	937	GLU	2.6
1	A	231	VAL	2.6
1	A	639	LYS	2.6
1	D	631	PRO	2.6
1	D	759	PHE	2.6
3	С	161	ARG	2.6
1	A	640	THR	2.5
1	А	968	LEU	2.5



Mol	Chain	Res	Type	RSRZ	
1	А	240	GLY	2.5	
1	D	755	ASN	2.5	
1	D	241	SER	2.5	
1	D	209	PHE	2.5	
1	А	912	GLU	2.5	
1	D	997	ASP	2.4	
3	F	47	ASP	2.4	
1	А	802	GLY	2.4	
1	А	788	LYS	2.4	
1	D	248	ALA	2.4	
1	А	527	GLU	2.4	
3	С	169	GLU	2.4	
3	F	162	ARG	2.4	
1	А	157	GLY	2.3	
1	D	246	GLY	2.3	
1	D	906	MET	2.3	
1	D	195	THR	2.3	
1	А	634	LEU	2.3	
1	А	757	PRO	2.3	
1	D	748	GLU	2.3	
1	D	698	GLY	2.3	
1	D	980	LYS	2.3	
1	А	224	GLY	2.3	
3	С	134	GLU	2.3	
1	А	913	ILE	2.3	
1	А	225	THR	2.2	
1	D	661	LEU	2.2	
1	А	801	VAL	2.2	
1	А	941	GLU	2.2	
1	D	224	GLY	2.2	
1	D	987	LYS	2.2	
1	D	738	LEU	2.2	
1	A	455	GLU	2.2	
1	D	632	ASN	2.2	
1	D	230	GLU	2.2	
3	F	169	169 GLU 2		
1	D	216	LYS	2.1	
3	F	150	GLN	2.1	
3	С	157	ARG	2.1	
3	F	110	ARG	2.1	
1	D	188	ILE	2.1	
1	D 800 LEU 2.1		2.1		



Mol	Chain	Res	Type	RSRZ
3	С	133	ASP	2.1
1	D	993	ILE	2.1
1	D	221	LEU	2.1
1	А	687	PHE	2.1
1	А	979	LYS	2.1
1	А	879	LYS	2.1
1	D	183	GLY	2.1
1	А	759	PHE	2.0
1	А	985	PHE	2.0
1	D	214	GLU	2.0
1	D	455	GLU	2.0
3	F	146	ARG	2.0
1	D	763	SER	2.0
3	С	106	ASP	2.0
1	A	756	PRO	2.0
1	D	202	HIS	2.0
1	D	262	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
4	SO4	F	201	5/5	0.86	0.16	129,129,130,131	0
4	SO4	В	101	5/5	0.90	0.17	99,100,101,101	0
4	SO4	D	1102	5/5	0.93	0.18	92,92,94,95	0
4	SO4	D	1101	5/5	0.95	0.13	55,56,60,61	0
4	SO4	А	1101	5/5	0.96	0.16	46,49,50,51	0
4	SO4	А	1103	5/5	0.98	0.09	49,51,51,53	0



Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	B -factors $(Å^2)$	Q<0.9
4	SO4	А	1104	5/5	0.98	0.09	$34,\!36,\!39,\!47$	0
4	SO4	D	1103	5/5	0.98	0.09	37,39,44,44	0
4	SO4	А	1102	5/5	0.98	0.09	38,43,44,48	0

6.5 Other polymers (i)

There are no such residues in this entry.

