

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 16, 2023 – 03:09 AM JST

PDB ID	:	6KP2
Title	:	Quadruple mutant plasmodium falciparum dihydrofolate reductase complexed
		with B10042
Authors	:	Vanichtanankul, J.; Vitsupakorn, D.
Deposited on	:	2019-08-14
Resolution	:	1.97  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	608	4% 79%	6%	14%
1	В	608	76%	9% •	12%



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9649 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bifunctional dihydrofolate reductase-thymidylate synthase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	520	Total 4341	C 2819	N 715	0 782	S 25	0	2	0
1	В	532	Total 4424	C 2863	N 729	O 807	$\begin{array}{c} \mathrm{S} \\ \mathrm{25} \end{array}$	0	0	0

• Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
0	Δ	1	Total	С	Ν	0	Р	0	0
	2 A	L	48	21	7	17	3	0	0
0	Р	1	Total	С	Ν	0	Р	0	0
	D	L	48	21	7	17	3	0	0

• Molecule 3 is 5-(3-chlorophenyl)-6-(3-phenoxypropyl)pyrimidine-2,4-diamine (three-letter code: DQ3) (formula: C<sub>19</sub>H<sub>19</sub>ClN<sub>4</sub>O) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
3	Δ	1	Total	С	Cl	Ν	0	0	0	
5	3 A	1	25	19	1	4	1	0	0	
2	В	1	Total	С	Cl	Ν	Ο	0	0	
J	D	1	25	19	1	4	1	0	0	



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0



• Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	381	Total O 381 381	0	0
6	В	335	Total O 335 335	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase



• Molecule 1: Bifunctional dihydrofolate reductase-thymidylate synthase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	56.99Å 155.94Å 165.34Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution (Å)	32.20 - 1.97	Depositor
Resolution (A)	32.18 - 1.97	EDS
% Data completeness	99.4 (32.20-1.97)	Depositor
(in resolution range)	99.4 (32.18-1.97)	EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.15 (at 1.97 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
B B.	0.173 , $0.213$	Depositor
$n, n_{free}$	0.183 , $0.222$	DCC
$R_{free}$ test set	5079 reflections $(4.89%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	22.8	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , $50.0$	EDS
L-test for $twinning^2$	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9649	wwPDB-VP
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, DQ3, PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
1VIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.79	0/4449	0.90	1/6007~(0.0%)	
1	В	0.78	0/4527	0.90	2/6116~(0.0%)	
All	All	0.78	0/8976	0.90	3/12123~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	421	ARG	NE-CZ-NH2	-8.97	115.82	120.30
1	А	421	ARG	NE-CZ-NH2	-8.01	116.30	120.30
1	В	455	GLN	CB-CA-C	-6.38	97.64	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	165	GLY	Peptide
1	В	304	LYS	Peptide



#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4341	0	4316	28	0
1	В	4424	0	4373	38	0
2	А	48	0	26	2	0
2	В	48	0	26	1	0
3	А	25	0	0	0	0
3	В	25	0	0	0	0
4	А	6	0	8	0	0
4	В	6	0	8	0	0
5	А	5	0	0	0	0
5	В	5	0	0	0	0
6	А	381	0	0	6	0
6	В	335	0	0	12	0
All	All	9649	0	8757	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom_1	Atom_2	Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
1:B:82:ASN:HB3	6:B:1052:HOH:O	1.62	0.98
1:B:4:GLN:HE21	1:B:4:GLN:HA	1.23	0.97
1:A:349:GLY:HA3	1:A:554:ASN:ND2	1.81	0.95
1:B:573:THR:HG22	6:B:1079:HOH:O	1.70	0.91
1:B:4:GLN:HE21	1:B:4:GLN:CA	1.92	0.83
1:A:573:THR:HG22	6:A:1126:HOH:O	1.86	0.76
1:B:603:SER:H	1:B:608:ALA:HB3	1.50	0.75
1:A:573:THR:HG23	6:A:1021:HOH:O	1.86	0.75
1:B:4:GLN:HA	1:B:4:GLN:NE2	1.99	0.74
1:B:50:CYS:SG	6:B:1088:HOH:O	2.45	0.74
1:A:349:GLY:HA3	1:A:554:ASN:HD21	1.54	0.69
1:B:139:ASP:N	1:B:139:ASP:OD1	2.25	0.67
1:B:210:VAL:HB	6:B:953:HOH:O	1.95	0.66
1:B:109:TRP:CZ2	1:B:117:LYS:HD2	2.32	0.65
1:B:166:GLY:HA3	2:B:701:NDP:PA	2.39	0.63



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:375:PHE:HD1	6:A:1093:HOH:O	1.82	0.62	
1:B:407:ASN:ND2	6:B:805:HOH:O	2.32	0.62	
1:B:294:ASN:ND2	6:B:807:HOH:O	2.34	0.61	
1:A:109:TRP:CZ2	1:A:117:LYS:HD2	2.36	0.60	
1:A:332:ILE:CD1	1:A:514:LEU:HB3	2.34	0.57	
1:B:136:PHE:CD2	1:B:142:ILE:HD11	2.41	0.56	
1:B:165:GLY:HA3	1:B:170:TYR:CZ	2.43	0.54	
1:B:461:ASN:OD1	6:B:801:HOH:O	2.19	0.53	
1:A:102:VAL:HB	1:A:164:LEU:HD11	1.91	0.52	
1:A:344:ASP:HA	6:B:1065:HOH:O	2.08	0.52	
1:B:51:ILE:HD13	1:B:187:ILE:HD12	1.92	0.52	
1:A:210:VAL:HB	6:A:1043:HOH:O	2.10	0.52	
1:A:487:LEU:HD23	1:A:487:LEU:N	2.26	0.51	
1:B:345:ARG:HG2	1:B:345:ARG:HH11	1.76	0.51	
1:B:297:LYS:HA	1:B:297:LYS:HE2	1.92	0.50	
1:B:582:ASN:HB3	6:B:948:HOH:O	2.12	0.49	
1:B:297:LYS:HA	1:B:297:LYS:CE	2.43	0.49	
1:B:147:GLU:N	1:B:147:GLU:OE1	2.45	0.49	
1:A:308:HIS:O	1:A:311:ASP:HB2	2.14	0.48	
1:A:206:GLN:HE21	1:A:206:GLN:CA	2.28	0.47	
1:B:321:LYS:CE	6:B:818:HOH:O	2.62	0.47	
1:B:109:TRP:CE2	1:B:117:LYS:HD2	2.51	0.46	
1:B:171:GLN:NE2	1:B:175:GLU:HG3	2.31	0.46	
1:B:59:ARG:O	1:B:63:THR:HG23	2.16	0.46	
1:A:174:LEU:HD12	1:A:198:PRO:HG2	1.97	0.45	
1:A:600:GLU:HG3	6:A:812:HOH:O	2.16	0.45	
1:A:195:VAL:HG21	2:A:701:NDP:H4D	1.97	0.45	
1:B:133:LYS:O	1:B:133:LYS:HD3	2.17	0.45	
1:A:349:GLY:CA	1:A:554:ASN:ND2	2.67	0.44	
1:A:40:LEU:O	2:A:701:NDP:H2N	2.17	0.44	
1:B:311:ASP:OD2	1:B:561:LYS:NZ	2.44	0.44	
1:B:20:VAL:HG21	1:B:38:ARG:NH2	2.32	0.44	
1:A:17:CYS:HA	1:A:39:GLY:O	2.18	0.44	
1:A:308:HIS:C	6:A:818:HOH:O	2.56	0.43	
1:B:20:VAL:HG22	6:B:876:HOH:O	2.18	0.43	
1:B:197:PHE:CD1	1:B:198:PRO:HD2	2.54	0.43	
1:B:345:ARG:HH11	1:B:345:ARG:CG	2.31	0.43	
1:B:379:ILE:HD13	1:B:520:PHE:CD1	2.54	0.43	
1:A:109:TRP:CE2	1:A:117:LYS:HD2	2.54	0.43	
1:A:287:GLU:HB2	1:B:69:LYS:NZ	2.34	0.42	
1:B:102:VAL:HB	1:B:164:LEU:HD11	2.00	0.42	



6KP $2$
---------

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:487:LEU:N	1:A:487:LEU:CD2	2.83	0.42
1:A:516:LEU:HD23	1:A:516:LEU:HA	1.93	0.42
1:A:197:PHE:CG	1:A:198:PRO:HD2	2.55	0.42
1:B:124:ASN:N	1:B:124:ASN:HD22	2.18	0.42
1:A:487:LEU:HD23	1:A:487:LEU:H	1.86	0.41
1:B:165:GLY:HA3	1:B:170:TYR:CE2	2.56	0.41
1:A:143:ILE:CD1	1:A:149:LEU:HB2	2.51	0.40
1:A:214:TYR:O	1:A:220:THR:HA	2.21	0.40
1:B:309:PRO:HA	6:B:851:HOH:O	2.22	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erce	ntile	es
1	А	510/608~(84%)	494 (97%)	16 (3%)	0	1	.00	100	)
1	В	522/608~(86%)	496 (95%)	22 (4%)	4 (1%)		19	9	
All	All	1032/1216~(85%)	990 (96%)	38 (4%)	4 (0%)		34	22	

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	50	CYS
1	В	607	ALA
1	В	309	PRO
1	В	347	GLY



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	488/570~(86%)	478 (98%)	10 (2%)	55 48
1	В	496/570~(87%)	469 (95%)	27 (5%)	22 10
All	All	984/1140~(86%)	947~(96%)	37 (4%)	33 21

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	49	LYS
1	А	67	GLU
1	А	72	LYS
1	А	79	LYS
1	А	81	LEU
1	А	97	LYS
1	А	114	LYS
1	А	145	LYS
1	А	206	GLN
1	А	440	GLU
1	В	4	GLN
1	В	8	VAL
1	В	49	LYS
1	В	50	CYS
1	В	74	LYS
1	В	79	LYS
1	В	97	LYS
1	В	107	THR
1	В	132	LYS
1	В	137	ASP
1	В	139	ASP
1	В	144	ASN
1	В	147	GLU
1	В	151	VAL
1	В	201	ASN
1	В	202	GLU
1	В	285	GLU



Mol	Chain	Res	Type
1	В	294	ASN
1	В	297	LYS
1	В	298	GLU
1	В	303	ASN
1	В	307	ILE
1	В	345	ARG
1	В	346	THR
1	В	348	VAL
1	В	379	ILE
1	В	606	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	29	ASN
1	А	99	GLN
1	А	157	ASN
1	А	206	GLN
1	А	394	ASN
1	А	400	ASN
1	А	407	ASN
1	А	424	ASN
1	А	554	ASN
1	В	4	GLN
1	В	88	ASN
1	В	99	GLN
1	В	144	ASN
1	В	303	ASN
1	В	398	ASN
1	В	407	ASN
1	В	415	ASN
1	В	424	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tiple	B	ond leng	gths	E	ond ang	gles
INIOI	Type	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	NDP	В	701	-	45,52,52	1.96	16 (35%)	53,80,80	1.56	10 (18%)
5	PO4	А	704	-	$4,\!4,\!4$	1.57	1 (25%)	6,6,6	0.49	0
3	DQ3	А	702	-	27,27,27	1.48	4 (14%)	36,36,36	2.23	12 (33%)
4	GOL	А	703	-	$5,\!5,\!5$	0.09	0	5,5,5	0.18	0
3	DQ3	В	702	-	27,27,27	1.32	3 (11%)	36,36,36	1.95	7 (19%)
5	PO4	В	704	-	4,4,4	0.91	0	6,6,6	0.78	0
2	NDP	А	701	-	45,52,52	2.08	9 (20%)	53,80,80	1.65	10 (18%)
4	GOL	В	703	-	$5,\!5,\!5$	0.13	0	5,5,5	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	В	701	-	-	7/30/77/77	0/5/5/5
3	DQ3	А	702	-	-	2/11/11/11	0/3/3/3
4	GOL	А	703	-	-	0/4/4/4	-
3	DQ3	В	702	-	-	0/11/11/11	0/3/3/3
2	NDP	А	701	-	-	3/30/77/77	0/5/5/5
4	GOL	В	703	-	-	3/4/4/4	-

All (33) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	А	701	NDP	C2A-N3A	5.90	1.41	1.32
2	А	701	NDP	O4D-C1D	4.54	1.52	1.42
2	А	701	NDP	C3B-C4B	4.46	1.64	1.53
2	А	701	NDP	O4B-C1B	4.26	1.47	1.41
3	А	702	DQ3	C5-CAI	-4.23	1.42	1.50
2	А	701	NDP	C2A-N1A	4.21	1.41	1.33
2	В	701	NDP	PA-O5B	4.13	1.76	1.59
2	В	701	NDP	O4B-C1B	4.12	1.46	1.41
2	А	701	NDP	P2B-O2B	3.92	1.66	1.59
2	В	701	NDP	C2A-N3A	3.90	1.38	1.32
2	В	701	NDP	C2A-N1A	3.74	1.40	1.33
2	В	701	NDP	C6N-C5N	3.71	1.40	1.33
2	А	701	NDP	O3D-C3D	3.59	1.51	1.43
3	В	702	DQ3	C5-CAI	-3.55	1.43	1.50
2	В	701	NDP	C4N-C3N	-3.39	1.43	1.49
3	А	702	DQ3	C4-N3	3.32	1.40	1.34
3	А	702	DQ3	CAL-CAK	2.98	1.43	1.38
2	В	701	NDP	PN-O5D	2.96	1.71	1.59
2	А	701	NDP	C4N-C3N	-2.96	1.44	1.49
2	В	701	NDP	O4D-C4D	2.81	1.51	1.45
3	В	702	DQ3	CAJ-C4	-2.66	1.44	1.51
3	А	702	DQ3	C6-N1	2.61	1.38	1.35
2	В	701	NDP	C4N-C5N	-2.56	1.42	1.48
3	В	702	DQ3	CAK-CL	2.41	1.79	1.74
2	В	701	NDP	C2N-C3N	2.39	1.41	1.34
5	А	704	PO4	P-01	2.22	1.56	1.50
2	А	701	NDP	P2B-O3X	-2.22	1.46	1.54
2	В	701	NDP	C5A-C4A	-2.20	1.35	1.40
2	В	701	NDP	O3D-C3D	2.18	1.48	1.43
2	В	701	NDP	$\overline{C6A-C5A}$	-2.17	1.35	1.43
2	В	701	NDP	P2B-O3X	-2.15	1.46	1.54
2	В	701	NDP	PN-O2N	-2.14	1.45	1.55
2	В	701	NDP	O4D-C1D	2.09	1.47	1.42

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	702	DQ3	C2-N3-C4	7.79	122.73	116.24
3	В	702	DQ3	C2-N3-C4	7.31	122.33	116.24
2	В	701	NDP	N3A-C2A-N1A	-5.38	120.27	128.68
3	А	702	DQ3	N3-C2-N1	-4.04	119.08	125.42
2	А	701	NDP	O2N-PN-O1N	4.01	132.07	112.24
3	В	702	DQ3	C2-N1-C6	3.94	121.34	116.99



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	701	NDP	O4D-C1D-N1N	-3.93	100.37	108.06
3	А	702	DQ3	C5-C6-N1	-3.91	120.32	122.52
2	А	701	NDP	C1B-N9A-C4A	-3.83	119.92	126.64
2	А	701	NDP	N3A-C2A-N1A	-3.78	122.78	128.68
3	А	702	DQ3	C5-C4-N3	-3.70	119.04	123.61
3	В	702	DQ3	C5-C4-N3	-3.69	119.06	123.61
3	В	702	DQ3	N3-C2-N1	-3.49	119.94	125.42
3	А	702	DQ3	C2-N1-C6	3.31	120.64	116.99
2	В	701	NDP	O2A-PA-O1A	3.29	128.52	112.24
2	В	701	NDP	O2N-PN-O5D	3.28	122.98	107.75
2	В	701	NDP	O3X-P2B-O2X	3.14	119.63	107.64
2	А	701	NDP	O3X-P2B-O2X	2.90	118.71	107.64
2	А	701	NDP	O4B-C4B-C5B	2.86	118.77	109.37
3	А	702	DQ3	NAG-C6-N1	2.83	121.03	117.03
3	В	702	DQ3	C5-C6-N1	-2.77	120.97	122.52
2	В	701	NDP	C1B-N9A-C4A	-2.74	121.83	126.64
3	А	702	DQ3	CAU-CAV-CAT	2.71	123.30	118.96
2	В	701	NDP	O2A-PA-O5B	-2.65	95.43	107.75
3	А	702	DQ3	C6-C5-C4	2.65	118.10	115.91
3	А	702	DQ3	CAR-OAS-CAT	-2.62	111.07	117.93
2	В	701	NDP	O2N-PN-O1N	2.60	125.10	112.24
3	А	702	DQ3	CAI-C5-C4	-2.48	121.45	123.46
2	А	701	NDP	C2B-C3B-C4B	-2.37	96.84	101.99
2	А	701	NDP	C4A-C5A-N7A	-2.35	106.95	109.40
3	В	702	DQ3	CAI-CAL-CAK	2.32	122.44	119.24
2	В	701	NDP	C4A-C5A-N7A	-2.32	106.98	109.40
3	А	702	DQ3	NAH-C2-N3	2.21	120.70	117.25
2	А	701	NDP	O2A-PA-O5B	-2.13	97.87	107.75
2	В	701	NDP	O2B-P2B-O1X	-2.12	101.22	109.39
3	В	702	DQ3	CAI-C5-C4	-2.09	121.77	123.46
2	В	701	NDP	C5D-C4D-C3D	-2.07	107.42	115.18
2	А	701	NDP	O3X-P2B-O2B	-2.05	96.82	105.99
3	A	702	DQ3	CAX-CAT-CAV	-2.02	117.06	120.18

Continued from previous page...

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	701	NDP	C5D-O5D-PN-O1N
2	В	701	NDP	C5D-O5D-PN-O1N
4	В	703	GOL	C1-C2-C3-O3
4	В	703	GOL	O2-C2-C3-O3



Mol	Chain	Res	Type	Atoms
3	А	702	DQ3	CAV-CAT-OAS-CAR
3	А	702	DQ3	CAX-CAT-OAS-CAR
4	В	703	GOL	O1-C1-C2-O2
2	В	701	NDP	C4D-C5D-O5D-PN
2	В	701	NDP	O4D-C1D-N1N-C2N
2	А	701	NDP	O4D-C1D-N1N-C2N
2	А	701	NDP	C2D-C1D-N1N-C2N
2	В	701	NDP	C2D-C1D-N1N-C2N
2	В	701	NDP	C2B-O2B-P2B-O2X
2	В	701	NDP	C5D-O5D-PN-O3
2	В	701	NDP	C2N-C3N-C7N-N7N

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	701	NDP	1	0
2	А	701	NDP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	520/608~(85%)	0.06	23 (4%) 34 36	13, 24, 54, 82	0
1	В	532/608~(87%)	0.51	72 (13%) 3 3	13, 28, 73, 118	0
All	All	1052/1216~(86%)	0.29	95 (9%) 9 10	13, 25, 67, 118	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	136	PHE	9.3
1	В	607	ALA	8.3
1	В	608	ALA	6.7
1	В	134	GLU	6.3
1	В	346	THR	6.1
1	В	345	ARG	5.8
1	В	130	THR	5.3
1	В	116	PHE	5.3
1	В	131	LEU	5.0
1	В	303	ASN	4.7
1	В	137	ASP	4.5
1	В	164	LEU	4.5
1	В	138	GLU	4.5
1	В	135	ASP	4.2
1	В	14	ILE	4.0
1	В	75	TYR	4.0
1	В	67	GLU	3.9
1	В	348	VAL	3.8
1	В	49	LYS	3.7
1	В	163	ILE	3.7
1	В	70	TYR	3.6
1	В	284	ASP	3.6
1	В	347	GLY	3.6
1	В	283	ASP	3.5



6KP $2$
---------

Mol	Chain	Res	Type	RSRZ	
1	А	96 LYS		3.5	
1	В	5	VAL	3.4	
1	В	115	LYS	3.4	
1	В	166	GLY	3.4	
1	В	73 LEU		3.4	
1	В	151	VAL	3.3	
1	А	29	ASN	3.3	
1	А	309	PRO	3.3	
1	В	310	ASN	3.2	
1	А	79	LYS	3.2	
1	В	80	TYR	3.2	
1	А	75	TYR	3.2	
1	В	86	VAL	3.1	
1	В	9	PHE	3.1	
1	В	8	VAL	3.0	
1	А	290	PHE	3.0	
1	В	35	TYR	2.9	
1	В	308	HIS	2.9	
1	В	165	GLY	2.9	
1	В	144	ASN	2.8	
1	В	68	SER	2.8	
1	В	129	ARG	2.8	
1	В	286	GLU	2.8	
1	А	14	ILE	2.7	
1	В	132	LYS	2.7	
1	А	82	ASN	2.7	
1	В	103	VAL	2.7	
1	В	127	LEU	2.7	
1	В	306	SER	2.6	
1	А	182	ILE	2.6	
1	А	306	SER	2.6	
1	А	305	ASN	2.6	
1	В	203	ASN	2.5	
1	В	229	THR	2.5	
1	В	285	GLU	2.5	
1	В	606	MET	2.5	
1	А	81	LEU	2.5	
1	А	164	LEU	2.4	
1	В	59	ARG	2.4	
1	В	114	LYS	2.4	
1	А	607	ALA	2.4	
1	В	15	CYS	2.4	



Mol	Chain	$\mathbf{Res}$	Type	RSRZ	
1	В	148	ASP	2.4	
1	В	85	THR	2.3	
1	А	31	VAL	2.3	
1	В	71	GLU	2.3	
1	В	66	ASN	2.3	
1	А	71	GLU	2.3	
1	В	102	VAL	2.3	
1	В	298	GLU	2.3	
1	В	139	ASP	2.3	
1	В	113	PRO	2.2	
1	В	72	LYS	2.2	
1	А	287	GLU	2.2	
1	В	69	LYS	2.2	
1	В	287	GLU	2.2	
1	В	307	ILE	2.1	
1	А	286	GLU	2.1	
1	А	78	CYS	2.1	
1	В	106	ARG	2.1	
1	А	4	GLN	2.1	
1	В	170	TYR	2.1	
1	А	493	LEU	2.1	
1	В	605	ASP	2.1	
1	В	202	GLU	2.1	
1	В	309	PRO	2.1	
1	В	125	VAL	2.1	
1	А	308	HIS	2.0	
1	В	126	ILE	2.0	
1	В	473	LEU	2.0	
1	А	162	PHE	2.0	

Continued from previous page...

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	NDP	В	701	48/48	0.87	0.17	30,50,73,76	0
3	DQ3	В	702	25/25	0.90	0.16	30,39,70,76	0
4	GOL	В	703	6/6	0.94	0.10	24,29,32,35	0
5	PO4	В	704	5/5	0.94	0.17	44,45,52,55	0
3	DQ3	А	702	25/25	0.95	0.13	15,20,48,49	0
5	PO4	А	704	5/5	0.95	0.11	34,37,47,56	0
4	GOL	А	703	6/6	0.95	0.14	22,27,30,32	0
2	NDP	А	701	48/48	0.98	0.07	18,23,28,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











## 6.5 Other polymers (i)

There are no such residues in this entry.

