

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 04:28 PM EDT

PDB ID	:	2KQ9
BMRB ID	:	16594
Title	:	Solution structure of DnaK suppressor protein from Agrobacterium tumefa-
		ciens C58. Northeast Structural Genomics Consortium target AtT12/Ontario
		Center for Structural Proteomics Target atc0888
Authors	:	Wu, B.; Yee, A.; Fares, C.; Lemak, A.; Semest, A.; Montelione, G.T.; Ar-
		rowsmith, C.; Northeast Structural Genomics Consortium (NESG); Ontario
		Centre for Structural Proteomics (OCSP)
Deposited on	:	2009-11-02

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	v_1n_11_5_13_A (Berjanski et al., 2005)
:	Wang et al. (2010)
:	v1.2
:	v1.2
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.33
	: : : : :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive
Methe	$(\# { m Entries})$	$(\# { m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of	chain	
1	А	112	63%	•	33%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues										
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model									
1 A:8-A:28, A:58-A:111 (75) 0.54 3										

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	2, 3, 7, 10, 11, 16, 17, 20
2	6, 12, 14, 18
3	13, 15, 19
4	1, 4
Single-model clusters	5; 8; 9



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1745 atoms, of which 861 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called DnaK suppressor protein.

Mol	Chain	Residues		Atoms								
1	Δ	119	Total	С	Н	Ν	0	S	0			
	A	112	1744	537	861	163	178	5	0			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference		
А	1	LEU	MET	engineered mutation	PDB 2KQ9		

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms
2	А	1	Total Zn 1 1



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: DnaK suppressor protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

• Molecule 1: DnaK suppressor protein

Chain A:						60%	, D								7	7%				339	%				-
L1 A2 G4 K5 K5 M7	R19 Q20	K29 I30	E31 A32	D33 F34	E35 E35	E30 P37	N39	P40 D41	D42	E43	D44 R45	A46	S47 E48	S50 N51		E53 V54	D56 E57	<u>662</u>	L66	T69		G82	R90	L112	



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1320
Number of shifts mapped to atoms	1320
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	591	595	594	3 ± 1
All	All	11840	11900	11880	68

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 21 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:105:CYS:SG	1:A:108:CYS:HB2	0.76	2.21	5	4
1:A:105:CYS:SG	1:A:108:CYS:CB	0.57	2.91	5	4
1:A:105:CYS:SG	1:A:107:GLU:HG3	0.54	2.42	6	1
1:A:27:LEU:HB2	1:A:58:LEU:HA	0.51	1.83	8	2
1:A:105:CYS:SG	1:A:107:GLU:OE1	0.50	2.69	17	2



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	75/112~(67%)	$72\pm1~(96\pm2\%)$	$3\pm1~(4\pm2\%)$	0±1 (0±1%)	38 78
All	All	1500/2240~(67%)	1439~(96%)	55 (4%)	6~(0%)	38 78

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	8	ASN	4
1	А	58	LEU	1
1	А	84	CYS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	61/93~(66%)	59 ± 1 (96 $\pm2\%$)	$2\pm1 (4\pm2\%)$	36 84
All	All	1220/1860~(66%)	1174 (96%)	46 (4%)	36 84

5 of 14 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	14	LYS	9
1	А	19	ARG	9
1	А	95	ARG	7
1	А	107	GLU	5
1	А	24	TYR	4



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1320
Number of shifts mapped to atoms	1320
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	110	-0.31 ± 0.11	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	105	0.13 ± 0.04	None needed (< 0.5 ppm)
$^{13}C'$	102	-0.27 ± 0.08	None needed (< 0.5 ppm)
¹⁵ N	103	-0.75 ± 0.30	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 916 atoms were assigned a chemical shift out of a possible 1044. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	372/376~(99%)	152/153~(99%)	148/150~(99%)	72/73~(99%)
Sidechain	513/613~(84%)	350/395~(89%)	158/182~(87%)	5/36~(14%)

Continued on next page...



Continucu	Continuca from pretious page						
	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$			
Aromatic	31/55~(56%)	20/26~(77%)	11/27~(41%)	0/2~(0%)			
Overall	916/1044 (88%)	522/574~(91%)	317/359~(88%)	77/111 (69%)			

Continued from previous page...


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (1)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3081
Intra-residue (i-j =0)	701
Sequential (i-j =1)	702
Medium range ($ i-j >1$ and $ i-j <5$)	815
Long range $(i-j \ge 5)$	710
Inter-chain	0
Hydrogen bond restraints	133
Disulfide bond restraints	20
Total dihedral-angle restraints	129
Number of unmapped restraints	3
Number of restraints per residue	28.7
Number of long range restraints per residue ¹	6.5

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	13.2	0.2
0.2-0.5 (Medium)	6.0	0.47
>0.5 (Large)	6.5	1.34



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max ($^{\circ}$)
1.0-10.0 (Small)	1.6	3.4
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints type	Count	$\%^1$	Vie	lated	3	Consis	tently	Violated ⁴
Restraints type	Count	70-	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	701	22.8	8	1.1	0.3	0	0.0	0.0
Backbone-Backbone	2	0.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	500	16.2	7	1.4	0.2	0	0.0	0.0
Sidechain-Sidechain	199	6.5	1	0.5	0.0	0	0.0	0.0
Sequential (i-j =1)	702	22.8	32	4.6	1.0	0	0.0	0.0
Backbone-Backbone	138	4.5	3	2.2	0.1	0	0.0	0.0
Backbone-Sidechain	455	14.8	19	4.2	0.6	0	0.0	0.0
Sidechain-Sidechain	109	3.5	10	9.2	0.3	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	815	26.5	50	6.1	1.6	0	0.0	0.0
Backbone-Backbone	214	6.9	3	1.4	0.1	0	0.0	0.0
Backbone-Sidechain	401	13.0	23	5.7	0.7	0	0.0	0.0
Sidechain-Sidechain	200	6.5	24	12.0	0.8	0	0.0	0.0
Long range $(i-j \ge 5)$	710	23.0	28	3.9	0.9	1	0.1	0.0
Backbone-Backbone	28	0.9	1	3.6	0.0	0	0.0	0.0
Backbone-Sidechain	305	9.9	16	5.2	0.5	0	0.0	0.0
Sidechain-Sidechain	377	12.2	11	2.9	0.4	1	0.3	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	133	4.3	6	4.5	0.2	0	0.0	0.0
Disulfide bond	20	0.6	9	45.0	0.3	7	35.0	0.2
Total	3081	100.0	133	4.3	4.3	8	0.3	0.3
Backbone-Backbone	515	16.7	13	2.5	0.4	0	0.0	0.0
Backbone-Sidechain	1661	53.9	65	3.9	2.1	0	0.0	0.0
Sidechain-Sidechain	905	29.4	55	6.1	1.8	8	0.9	0.3

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	;	Maan (Å)	Mor (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (Å)	Max (Å)	$SD^{*}(A)$	Median (Å)
1	2	3	8	9	0	22	0.44	1.02	0.37	0.24
2	1	7	16	9	0	33	0.33	1.18	0.32	0.15
3	1	5	9	8	0	23	0.44	1.17	0.35	0.23
4	1	4	8	11	0	24	0.41	0.97	0.34	0.26
5	1	7	11	9	0	28	0.29	1.34	0.34	0.16
6	1	3	10	9	0	23	0.44	1.19	0.35	0.25
7	1	7	8	10	0	26	0.39	1.1	0.35	0.16
8	2	6	12	8	0	28	0.26	1.2	0.28	0.16
9	1	2	9	12	0	24	0.41	1.26	0.35	0.24
10	1	8	11	13	0	33	0.35	1.0	0.33	0.19
11	3	2	7	8	0	20	0.3	1.21	0.32	0.19


Continued on next page...




Madal ID			nber o		ations	5	Mean (Å)	Mar (Å)		Median (Å)	
Model ID	IR^{1}	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (A)	Max (Å)	SD^{6} (Å)	Median (A)	
12	1	3	11	9	0	24	0.41	0.97	0.35	0.2	
13	1	6	10	9	0	26	0.38	0.95	0.32	0.23	
14	1	4	6	9	0	20	0.48	1.18	0.35	0.29	
15	1	4	12	9	0	26	0.4	0.99	0.35	0.21	
16	2	4	8	10	0	24	0.42	1.18	0.33	0.22	
17	1	6	9	10	0	26	0.41	1.16	0.32	0.29	
18	2	5	9	9	0	25	0.41	1.23	0.35	0.24	
19	1	4	10	13	0	28	0.36	0.95	0.33	0.2	
20	0	4	15	12	0	31	0.25	1.26	0.28	0.17	

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right




9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2810(IR:693, SQ:670, MR:765, LR:682, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restra	aints	Fractio	n of the ensemble
IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Count^6	%
6	17	31	13	0	67	1	5.0
0	2	7	8	0	17	2	10.0
1	3	4	3	0	11	3	15.0
0	3	2	0	0	5	4	20.0
0	2	1	1	0	4	5	25.0
0	1	0	1	0	2	6	30.0
0	2	0	0	0	2	7	35.0
0	0	1	1	0	2	8	40.0
0	0	1	0	0	1	9	45.0
0	1	1	0	0	2	10	50.0
0	0	0	0	0	0	11	55.0
0	1	0	0	0	1	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
1	0	0	0	0	1	16	80.0
0	0	1	0	0	1	17	85.0
0	0	1	0	0	1	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	1	0	1	20	100.0

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(4,3)	1:A:84:CYS:SG	1:A:108:CYS:SG	20	0.97	0.15	0.92
(4,6)	1:A:84:CYS:SG	1:A:108:CYS:SG	20	0.97	0.15	0.92
(4,2)	1:A:84:CYS:SG	1:A:105:CYS:SG	20	0.79	0.29	0.91
(4,5)	1:A:84:CYS:SG	1:A:105:CYS:SG	20	0.79	0.29	0.91
(4,1)	1:A:84:CYS:SG	1:A:87:CYS:SG	20	0.77	0.29	0.9
(4,4)	1:A:84:CYS:SG	1:A:87:CYS:SG	20	0.77	0.29	0.9
(1,2931)	1:A:84:CYS:SG	1:A:108:CYS:CB	20	0.69	0.2	0.69
(1,101)	1:A:9:VAL:HG21	1:A:76:ILE:HG21	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG21	1:A:76:ILE:HG22	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG21	1:A:76:ILE:HG23	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG22	1:A:76:ILE:HG21	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG22	1:A:76:ILE:HG22	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG22	1:A:76:ILE:HG23	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG23	1:A:76:ILE:HG21	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG23	1:A:76:ILE:HG22	20	0.33	0.08	0.34
(1,101)	1:A:9:VAL:HG23	1:A:76:ILE:HG23	20	0.33	0.08	0.34

Continued on next page...



Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,2929)	1:A:84:CYS:SG	1:A:87:CYS:CB	19	0.69	0.47	0.3
(1,1453)	1:A:61:VAL:HA	1:A:65:GLU:HG2	18	0.2	0.06	0.2


Continued from previous page...

 $^1\mathrm{Number}$ of violated models, $^2\mathrm{Standard}$ deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,6)	1:A:84:CYS:SG	1:A:108:CYS:SG	5	1.34
(4,3)	1:A:84:CYS:SG	1:A:108:CYS:SG	5	1.34
(4,6)	1:A:84:CYS:SG	1:A:108:CYS:SG	20	1.26

Continued on next page...



Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,3)	1:A:84:CYS:SG	1:A:108:CYS:SG	20	1.26
(1,2929)	1:A:84:CYS:SG	1:A:87:CYS:CB	9	1.26
(1,2929)	1:A:84:CYS:SG	1:A:87:CYS:CB	18	1.23
(4,6)	1:A:84:CYS:SG	1:A:108:CYS:SG	11	1.21
(4,3)	1:A:84:CYS:SG	1:A:108:CYS:SG	11	1.21
(4,6)	1:A:84:CYS:SG	1:A:108:CYS:SG	8	1.2
(4,3)	1:A:84:CYS:SG	1:A:108:CYS:SG	8	1.2

Continued from previous page...



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	$\%^1$	Vie				Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$	
PSI	64	49.6	11	17.2	8.5	0	0.0	0.0	
PHI	65	50.4	4	6.2	3.1	0	0.0	0.0	
Total	129	100.0	15	11.6	11.6	0	0.0	0.0	

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories





10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber c	of violations	Moon (°)	M_{ov} (°)	SD (°)	Median (°)
Model ID	PSI	PHI	Total	Mean $(^{\circ})$	Max (°)	$SD(^{\circ})$	Median ()
1	2	1	3	2.37	3.1	0.66	2.5
2	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0.0	0.0	0.0	0.0
5	2	0	2	1.75	1.9	0.15	1.75
6	1	0	1	2.2	2.2	0.0	2.2
7	1	0	1	1.6	1.6	0.0	1.6
8	0	1	1	1.5	1.5	0.0	1.5
9	3	1	4	1.5	1.8	0.25	1.5
10	2	1	3	1.77	2.1	0.24	1.6
11	1	1	2	2.65	3.4	0.75	2.65
12	2	1	3	1.67	2.4	0.52	1.4
13	1	1	2	1.7	1.8	0.1	1.7
14	2	0	2	2.35	2.8	0.45	2.35
15	1	1	2	1.6	1.8	0.2	1.6
16	2	1	3	1.77	2.8	0.74	1.4
17	2	0	2	1.2	1.3	0.1	1.2
18	1	0	1	1.4	1.4	0.0	1.4
19	1	0	1	1.3	1.3	0.0	1.3
20	0	0	0	0.0	0.0	0.0	0.0







The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count^1	%
5	2	7	1	5.0
2	1	3	2	10.0
2	0	2	3	15.0
1	0	1	4	20.0
1	1	2	5	25.0
0	0	0	6	30.0
0	0	0	7	35.0
0	0	0	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0

Continued on next page...




PSI


PHI

Continued from previous page								
Nun	nber o	f violated restraints	Fraction of the ensemble					
PSI	PHI	Total	Count^1	%				
0	0	0	12	60.0				
0	0	0	13	65.0				
0	0	0	14	70.0				
0	0	0	15	75.0				
0	0	0	16	80.0				
0	0	0	17	85.0				
0	0	0	18	90.0				
0	0	0	19	95.0				
0	0	0	20	100.0				

Continued from previous page...

 1 Number of models with violations







10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models







10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	$Models^1$	Mean	SD^2	Median
(1,88)	1:A:82:GLY:C	1:A:83:THR:N	1:A:83:THR:CA	1:A:83:THR:C	5	1.9	0.46	1.6
(1,111)	1:A:97:LYS:N	1:A:97:LYS:CA	1:A:97:LYS:C	1:A:98:ALA:N	5	1.56	0.21	1.6
(1,34)	1:A:25:ARG:N	1:A:25:ARG:CA	1:A:25:ARG:C	1:A:26:ARG:N	4	1.82	0.29	1.85
(1,99)	1:A:91:ILE:N	1:A:91:ILE:CA	1:A:91:ILE:C	1:A:92:SER:N	3	2.2	0.64	1.8
(1,123)	1:A:107:GLU:N	1:A:107:GLU:CA	1:A:107:GLU:C	1:A:108:CYS:N	3	1.4	0.16	1.4
(1,25)	1:A:20:GLN:C	1:A:21:ARG:N	1:A:21:ARG:CA	1:A:21:ARG:C	2	2.6	0.8	2.6
(1,97)	1:A:90:ARG:N	1:A:90:ARG:CA	1:A:90:ARG:C	1:A:91:ILE:N	2	2.45	0.35	2.45
(1,10)	1:A:13:GLU:N	1:A:13:GLU:CA	1:A:13:GLU:C	1:A:14:LYS:N	2	1.25	0.05	1.25


¹ Number of violated models, ²Standard deviation, All angle values are in degree ($^{\circ}$)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation ($^{\circ}$)
(1,25)	1:A:20:GLN:C	1:A:21:ARG:N	1:A:21:ARG:CA	1:A:21:ARG:C	11	3.4
(1,99)	1:A:91:ILE:N	1:A:91:ILE:CA	1:A:91:ILE:C	1:A:92:SER:N	1	3.1
(1,97)	1:A:90:ARG:N	1:A:90:ARG:CA	1:A:90:ARG:C	1:A:91:ILE:N	14	2.8
(1,117)	1:A:104:PHE:N	1:A:104:PHE:CA	1:A:104:PHE:C	1:A:105:CYS:N	16	2.8
(1,88)	1:A:82:GLY:C	1:A:83:THR:N	1:A:83:THR:CA	1:A:83:THR:C	1	2.5
(1,88)	1:A:82:GLY:C	1:A:83:THR:N	1:A:83:THR:CA	1:A:83:THR:C	12	2.4
(1,34)	1:A:25:ARG:N	1:A:25:ARG:CA	1:A:25:ARG:C	1:A:26:ARG:N	6	2.2
(1,97)	1:A:90:ARG:N	1:A:90:ARG:CA	1:A:90:ARG:C	1:A:91:ILE:N	10	2.1
(1,34)	1:A:25:ARG:N	1:A:25:ARG:CA	1:A:25:ARG:C	1:A:26:ARG:N	5	1.9
(1,121)	1:A:106:GLN:N	1:A:106:GLN:CA	1:A:106:GLN:C	1:A:107:GLU:N	11	1.9

