

# Full wwPDB X-ray Structure Validation Report (i)

#### Mar 9, 2024 – 06:50 PM EST

PDB ID	:	3KRU
Title	:	Crystal Structure of the Thermostable Old Yellow Enzyme from Thermoanaer-
		obacter pseudethanolicus E39
Authors	:	Adalbjornsson, B.V.; Toogood, H.S.; Leys, D.; Scrutton, N.S.
Deposited on		
Resolution	:	1.60  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolDrobity		4 021 467
5		4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665(1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	343	95%	• •
1	В	343	93%	•••
1	С	343	94%	
1	D	343	91%	7% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	А	402	-	-	Х	-
3	ACT	В	402	-	-	Х	-
3	ACT	С	402	-	-	Х	-
3	ACT	D	402	-	-	Х	-

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



#### 3KRU

# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	335	Total	С	Ν	0	$\mathbf{S}$	5	11	0
	А	000	2681	1711	455	497	18	0		0
1	В	335	Total	С	Ν	0	S	0	13	0
	D	000	2713	1727	464	505	17	0		
1	С	334	Total	С	Ν	0	S	0	8	0
	U		2690	1706	461	507	16	0	8	0
1	л	225	Total	С	Ν	0	S	0	9	0
	I D	335	2687	1710	463	497	17	0	9	U

• Molecule 1 is a protein called NADH:flavin oxidoreductase/NADH oxidase.


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	338	HIS	-	expression tag	UNP B0KAH1
A	339	HIS	-	expression tag	UNP B0KAH1
А	340	HIS	-	expression tag	UNP B0KAH1
А	341	HIS	-	expression tag	UNP B0KAH1
А	342	HIS	-	expression tag	UNP B0KAH1
А	343	HIS	-	expression tag	UNP B0KAH1
В	338	HIS	-	expression tag	UNP B0KAH1
В	339	HIS	-	expression tag	UNP B0KAH1
В	340	HIS	-	expression tag	UNP B0KAH1
В	341	HIS	-	expression tag	UNP B0KAH1
В	342	HIS	-	expression tag	UNP B0KAH1
В	343	HIS	-	expression tag	UNP B0KAH1
С	338	HIS	-	expression tag	UNP B0KAH1
С	339	HIS	-	expression tag	UNP B0KAH1
С	340	HIS	-	expression tag	UNP B0KAH1
С	341	HIS	-	expression tag	UNP B0KAH1
С	342	HIS	-	expression tag	UNP B0KAH1
С	343	HIS	-	expression tag	UNP B0KAH1
D	338	HIS	-	expression tag	UNP B0KAH1
D	339	HIS	-	expression tag	UNP B0KAH1
D	340	HIS	-	expression tag	UNP B0KAH1



Chain	Residue	Modelled	Actual	Comment	Reference
D	341	HIS	-	expression tag	UNP B0KAH1
D	342	HIS	-	expression tag	UNP B0KAH1
D	343	HIS	-	expression tag	UNP B0KAH1


 $\bullet \ \ \ Molecule \ 2 \ is \ FLAVIN \ MONONUCLEOTIDE \ (three-letter \ code: \ FMN) \ (formula: \ C_{17}H_{21}N_4O_9P).$ 



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	Δ	1	Total	С	Ν	0	Р	0	0	
	А	T	31	17	4	9	1	0	0	
2	B	1	Total	С	Ν	0	Р	0	0	
	D	T	31	17	4	9	1	0	0	
2	С	1	Total	С	Ν	Ο	Р	0	0	
	U	1	31	17	4	9	1	0	0	
2	Л	1	Total	С	Ν	0	Р	0	0	
	2 D	1	31	17	4	9	1	0	0	

 $\bullet\,$  Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  ${\rm C_2H_3O_2}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	488	Total O 488 488	0	0
4	В	493	Total         O           493         493	0	0
4	С	416	Total         O           416         416	0	0
4	D	434	Total O 434 434	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NADH:flavin oxidoreductase/NADH oxidase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	86.92Å 97.38Å 94.39Å	Denesiten
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $92.34^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{D}$	64.82 - 1.60	Depositor
Resolution (Å)	$64.82 \ - \ 1.60$	EDS
% Data completeness	98.8 (64.82-1.60)	Depositor
(in resolution range)	98.8 (64.82-1.60)	EDS
R <sub>merge</sub>	0.06	Depositor
$\mathrm{R}_{sym}$	0.06	Depositor
$< I/\sigma(I) > 1$	2.66 (at 1.60Å)	Xtriage
Refinement program	REFMAC	Depositor
D D	0.152 , $0.179$	Depositor
$R, R_{free}$	0.160 , $0.188$	DCC
$R_{free}$ test set	10251  reflections  (5.02%)	wwPDB-VP
Wilson B-factor $(Å^2)$	14.5	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35, 61.4	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.48, < L^2 > = 0.31$	Xtriage
	0.018 for -h,-l,-k	
Estimated twinning fraction	0.005 for -h,l,k	Xtriage
	0.024 for h,-k,-l	
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12742	wwPDB-VP
Average B, all atoms $(Å^2)$	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
	RMSZ	# Z  > 5	RMSZ	# Z  > 5		
1	А	0.51	0/2763	0.68	2/3741~(0.1%)	
1	В	0.50	0/2796	0.67	2/3780~(0.1%)	
1	С	0.47	0/2752	0.65	2/3718~(0.1%)	
1	D	0.46	0/2761	0.65	2/3732~(0.1%)	
All	All	0.49	0/11072	0.66	8/14971~(0.1%)	

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	312	ARG	NE-CZ-NH1	8.34	124.47	120.30
1	D	312	ARG	NE-CZ-NH1	7.41	124.00	120.30
1	С	312	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	А	312	ARG	NE-CZ-NH1	6.84	123.72	120.30
1	С	312	ARG	NE-CZ-NH2	-6.81	116.90	120.30
1	А	312	ARG	NE-CZ-NH2	-6.15	117.22	120.30
1	D	312	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	В	312	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2681	0	2664	6	0
1	В	2713	0	2718	10	0
1	С	2690	0	2676	6	0
1	D	2687	0	2698	18	0
2	А	31	0	19	5	0
2	В	31	0	19	5	0
2	С	31	0	19	5	0
2	D	31	0	19	3	0
3	А	4	0	3	4	0
3	В	4	0	3	4	0
3	С	4	0	3	3	0
3	D	4	0	3	2	0
4	А	488	0	0	1	0
4	В	493	0	0	1	0
4	С	416	0	0	1	0
4	D	434	0	0	2	0
All	All	12742	0	10844	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:C:401:FMN:C4	3:C:402:ACT:H3	1.99	0.93
2:A:401:FMN:C4	3:A:402:ACT:H3	2.01	0.91
1:D:270:GLU:OE1	4:D:1738:HOH:O	1.94	0.85
1:C:4:LEU:HD11	1:C:19:ILE:HD11	1.60	0.83
1:D:259[B]:ASN:OD1	4:D:979:HOH:O	2.14	0.66
2:C:401:FMN:O4	3:C:402:ACT:H3	2.00	0.61
2:B:401:FMN:C4	3:B:402:ACT:CH3	2.80	0.60
1:B:4[B]:LEU:HD22	1:B:19:ILE:HD11	1.82	0.60
1:B:259[B]:ASN:OD1	4:B:857:HOH:O	2.17	0.58
1:B:159:VAL:HG22	1:B:212:PRO:HG2	1.85	0.58
1:D:135:VAL:HG13	1:D:139:LYS:NZ	2.19	0.57
1:D:4:LEU:HD23	1:D:296:LEU:HD12	1.88	0.56
2:A:401:FMN:O4	3:A:402:ACT:H3	2.06	0.55
1:D:200:ILE:O	1:D:204[A]:ARG:HG3	2.06	0.54
1:D:4:LEU:HD23	1:D:296:LEU:CD1	2.39	0.53
1:C:24:MET:HA	2:C:401:FMN:C5A	2.39	0.52
1:D:4:LEU:HD22	1:D:19:ILE:HD11	1.91	0.52
1:C:24:MET:HA	2:C:401:FMN:N5	2.25	0.51



1:D:4:LEU:HD11

1:D:204[B]:ARG:NH2

1:D:163:HIS:HE1

Continued from previo	Continued from previous page								
Atom-1	Atom-2	Interatomic	Clash						
		distance (Å)	overlap (Å)						
1:B:53:LEU:HD13	1:B:96:VAL:HG12	1.94	0.49						
1:A:24:MET:HA	2:A:401:FMN:N5	2.27	0.49						
1:A:277:ASN:ND2	4:A:488:HOH:O	2.44	0.49						
1:D:139:LYS:CE	1:D:185:TYR:OH	2.60	0.48						
1:A:24:MET:HA	2:A:401:FMN:C5A	2.43	0.48						
1:D:24:MET:HA	2:D:401:FMN:N5	2.29	0.48						
1:B:24:MET:HA	2:B:401:FMN:N5	2.28	0.48						
1:A:267:LYS:HA	1:A:270[B]:GLU:HG2	1.97	0.46						
1:C:267:LYS:HA	1:C:270[B]:GLU:HG2	1.97	0.46						
1:A:247:ASP:HA	1:A:281[A]:SER:OG	2.15	0.46						
1:B:24:MET:HA	2:B:401:FMN:C5A	2.45	0.46						
1:D:335:PHE:O	1:D:336:LYS:C	2.53	0.46						
1:D:24:MET:HA	2:D:401:FMN:C5A	2.46	0.46						
1:D:160:VAL:HG13	1:D:207:TRP:CH2	2.52	0.45						
2:A:401:FMN:C4	3:A:402:ACT:CH3	2.87	0.45						
1:D:223:MET:HG3	1:D:256[B]:VAL:HG11	1.97	0.45						
1:D:204[B]:ARG:NH2	1:D:208:PRO:O	2.48	0.44						
1:D:204[B]:ARG:HD2	1:D:213:ILE:HD12	1.98	0.44						
1:C:224[B]:GLU:OE1	4:C:403:HOH:O	2.21	0.44						
1:C:223:MET:HG3	1:C:256:VAL:HG11	2.00	0.44						
1:B:67:ILE:CD1	3:B:402:ACT:H3	2.48	0.43						
1:B:247:ASP:HA	1:B:281[B]:SER:OG	2.18	0.43						
2:C:401:FMN:C4	3:C:402:ACT:CH3	2.85	0.43						
2:D:401:FMN:C4	3:D:402:ACT:CH3	2.96	0.43						
2:B:401:FMN:C4	3:B:402:ACT:H2	2.49	0.43						
1:A:163:HIS:HE1	3:A:402:ACT:O	2.02	0.42						
1:B:53:LEU:HD13	1:B:96:VAL:CG1	2.49	0.42						
1:B:4[B]:LEU:HD23	1:B:296:LEU:CD1	2.50	0.41						
2:B:401:FMN:C4A	3:B:402:ACT:H2	2.51	0.41						
			i						

There are no symmetry-related clashes.

1:D:289:GLN:HG3

1:D:209:GLU:HA

3:D:402:ACT:OXT

#### 5.3Torsion angles (i)

#### 5.3.1Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

2.03

2.36

2.04

0.41

0.40

0.40



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	344/343~(100%)	336~(98%)	7 (2%)	1 (0%)	41	21
1	В	346/343~(101%)	338~(98%)	7(2%)	1 (0%)	41	21
1	С	340/343~(99%)	331~(97%)	9~(3%)	0	100	100
1	D	342/343~(100%)	335~(98%)	7(2%)	0	100	100
All	All	1372/1372~(100%)	1340~(98%)	30~(2%)	2~(0%)	47	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	187	ASN
1	В	187	ASN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	283/296~(96%)	283 (100%)	0	100	100	
1	В	291/296~(98%)	290 (100%)	1 (0%)	92	87	
1	$\mathbf{C}$	287/296~(97%)	286 (100%)	1 (0%)	92	87	
1	D	287/296~(97%)	287 (100%)	0	100	100	
All	All	1148/1184~(97%)	1146 (100%)	2~(0%)	93	88	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	163	HIS
1	С	163	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:



Mol	Chain	Res	Type
1	А	163	HIS
1	А	265	GLN
1	А	277	ASN
1	В	93	ASN
1	В	255	ASN
1	В	298	ASN
1	С	93	ASN
1	С	187	ASN
1	С	265	GLN
1	С	298	ASN
1	D	93	ASN
1	D	163	HIS
1	D	298	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res Link		Bo	ond leng	ths	В	ond ang	gles
Moi Type (	Chain Res			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
2	FMN	С	401	-	33,33,33	1.22	4 (12%)	$48,\!50,\!50$	1.46	9 (18%)



Mol	Turne	Chain	hain Res		Bo	ond leng	$_{\rm ths}$	Bond angles		
NIOI	Type	Unam	nes	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	FMN	В	401	-	33,33,33	1.26	5 (15%)	48,50,50	1.50	10 (20%)
2	FMN	А	401	-	33,33,33	1.19	4 (12%)	48,50,50	1.37	7 (14%)
3	ACT	А	402	-	3,3,3	0.93	0	3,3,3	1.28	0
3	ACT	D	402	-	3,3,3	0.72	0	3, 3, 3	1.34	0
3	ACT	В	402	-	3,3,3	0.64	0	$3,\!3,\!3$	1.58	1 (33%)
2	FMN	D	401	-	33,33,33	1.15	2 (6%)	48,50,50	1.46	8 (16%)
3	ACT	С	402	-	3,3,3	1.01	0	3,3,3	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	С	401	-	-	1/18/18/18	0/3/3/3
2	FMN	D	401	-	-	1/18/18/18	0/3/3/3
2	FMN	В	401	-	-	1/18/18/18	0/3/3/3
2	FMN	А	401	-	-	1/18/18/18	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	А	401	FMN	C4A-N5	3.72	1.38	1.30
2	В	401	FMN	C4A-N5	3.64	1.37	1.30
2	D	401	FMN	C4A-N5	3.26	1.37	1.30
2	С	401	FMN	C4A-N5	3.24	1.37	1.30
2	D	401	FMN	C9A-N10	-2.46	1.36	1.41
2	С	401	FMN	C9A-N10	-2.41	1.36	1.41
2	А	401	FMN	C9A-N10	-2.30	1.37	1.41
2	С	401	FMN	C4-N3	-2.29	1.34	1.38
2	А	401	FMN	C5A-N5	-2.27	1.35	1.39
2	В	401	FMN	C9A-N10	-2.25	1.37	1.41
2	С	401	FMN	C5A-N5	-2.14	1.35	1.39
2	В	401	FMN	C2-N3	-2.13	1.34	1.39
2	В	401	FMN	C4-N3	-2.12	1.34	1.38
2	В	401	FMN	C5A-N5	-2.07	1.35	1.39
2	А	401	FMN	C2-N3	-2.04	1.34	1.39

All (35) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	С	401	FMN	C4A-C10-N10	3.92	122.21	116.48
2	С	401	FMN	C4-C4A-N5	3.88	123.76	118.23
2	В	401	FMN	C4-C4A-N5	3.78	123.61	118.23
2	D	401	FMN	C4-C4A-N5	3.75	123.56	118.23
2	А	401	FMN	C4-C4A-N5	3.53	123.25	118.23
2	D	401	FMN	C4A-C10-N10	3.41	121.47	116.48
2	В	401	FMN	C4A-C10-N10	3.25	121.23	116.48
2	В	401	FMN	O4-C4-N3	-3.07	114.24	120.12
2	D	401	FMN	C10-C4A-N5	-3.06	118.36	124.86
2	С	401	FMN	C10-C4A-N5	-3.04	118.41	124.86
2	А	401	FMN	C4A-C10-N10	2.81	120.58	116.48
2	В	401	FMN	C10-C4A-N5	-2.78	118.97	124.86
2	А	401	FMN	C4A-C4-N3	2.68	120.00	113.19
2	В	401	FMN	C4A-C4-N3	2.64	119.90	113.19
2	С	401	FMN	C4A-C4-N3	2.63	119.86	113.19
2	С	401	FMN	C4-N3-C2	-2.58	120.88	125.64
2	D	401	FMN	C4-N3-C2	-2.55	120.93	125.64
2	D	401	FMN	C4A-C4-N3	2.52	119.59	113.19
2	А	401	FMN	P-O5'-C5'	2.46	125.07	118.30
2	А	401	FMN	C10-C4A-N5	-2.37	119.83	124.86
2	С	401	FMN	O4-C4-N3	-2.34	115.64	120.12
2	В	401	FMN	P-O5'-C5'	2.30	124.62	118.30
2	D	401	FMN	C7M-C7-C8	2.29	125.43	120.74
2	С	401	FMN	C4A-C10-N1	-2.24	119.52	124.73
2	D	401	FMN	C7M-C7-C6	-2.24	115.35	119.49
2	В	401	FMN	C4-N3-C2	-2.21	121.55	125.64
2	В	401	FMN	C4A-C10-N1	-2.17	119.69	124.73
2	А	401	FMN	C4-N3-C2	-2.16	121.64	125.64
2	А	401	FMN	C9A-C5A-N5	-2.16	120.09	122.43
2	В	401	FMN	O3P-P-O2P	2.15	115.85	107.64
2	D	401	FMN	C4A-C10-N1	-2.10	119.85	124.73
3	В	402	ACT	OXT-C-CH3	2.09	123.83	115.18
2	С	401	FMN	C10-N1-C2	2.09	121.09	116.90
2	В	401	FMN	C7M-C7-C6	-2.07	115.66	119.49
2	С	401	FMN	P-O5'-C5'	2.00	123.81	118.30

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	401	FMN	C4'-C5'-O5'-P
2	D	401	FMN	C4'-C5'-O5'-P
2	В	401	FMN	C4'-C5'-O5'-P



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	С	401	FMN	C4'-C5'-O5'-P


There are no ring outliers.

8 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	401	FMN	5	0
2	В	401	FMN	5	0
2	А	401	FMN	5	0
3	А	402	ACT	4	0
3	D	402	ACT	2	0
3	В	402	ACT	4	0
2	D	401	FMN	3	0
3	С	402	ACT	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	А	335/343~(97%)	-0.51	0 100 100	7, 12, 21, 28	4 (1%)
1	В	335/343~(97%)	-0.54	0 100 100	8, 13, 22, 30	2(0%)
1	С	334/343~(97%)	-0.44	1 (0%) 94 93	10, 16, 26, 36	0
1	D	335/343~(97%)	-0.52	0 100 100	11, 17, 24, 32	0
All	All	1339/1372~(97%)	-0.50	1 (0%) 95 94	7, 14, 24, 36	6 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	125	ASP	2.4

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
3	ACT	А	402	4/4	0.90	0.15	20,21,22,23	0




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q < 0.9
3	ACT	С	402	4/4	0.92	0.14	$23,\!25,\!25,\!25$	0
3	ACT	D	402	4/4	0.92	0.10	$26,\!27,\!27,\!28$	0
3	ACT	В	402	4/4	0.93	0.12	$21,\!23,\!25,\!25$	0
2	FMN	С	401	31/31	0.94	0.16	$6,\!17,\!18,\!19$	31
2	FMN	D	401	31/31	0.94	0.14	$7,\!15,\!17,\!18$	31
2	FMN	В	401	31/31	0.94	0.13	6,14,16,17	31
2	FMN	А	401	31/31	0.95	0.13	$6,\!13,\!15,\!15$	31

Continued from previous page...


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















# 6.5 Other polymers (i)

There are no such residues in this entry.

