



Full wwPDB EM Validation Report ⓘ

Mar 20, 2024 – 01:54 AM JST

PDB ID : 6KRD
EMDB ID : EMD-0756
Title : TRiC at 0.05 mM ADP-AlFx, Conformation 4, 0.05-C4
Authors : Jin, M.; Cong, Y.
Deposited on : 2019-08-21
Resolution : 4.38 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	559	
1	a	559	
2	B	527	
2	b	527	
3	D	528	
3	d	528	
4	E	562	
4	e	562	
5	G	534	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	g	534	
6	Q	568	
6	q	568	
7	Z	546	
7	z	546	
8	H	550	
8	h	550	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 61005 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-complex protein 1 subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	a	511	Total	C	N	O	S	0	0
			3851	2411	672	749	19		
1	A	513	Total	C	N	O	S	0	0
			3865	2420	675	751	19		

- Molecule 2 is a protein called T-complex protein 1 subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	b	502	Total	C	N	O	S	0	0
			3811	2385	658	754	14		
2	B	504	Total	C	N	O	S	0	0
			3824	2392	661	757	14		

- Molecule 3 is a protein called T-complex protein 1 subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	d	480	Total	C	N	O	S	0	0
			3655	2284	647	708	16		
3	D	429	Total	C	N	O	S	0	0
			3255	2024	579	638	14		

- Molecule 4 is a protein called T-complex protein 1 subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	e	501	Total	C	N	O	S	0	0
			3839	2411	659	748	21		
4	E	495	Total	C	N	O	S	0	0
			3786	2373	650	742	21		

- Molecule 5 is a protein called T-complex protein 1 subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	g	493	Total	C	N	O	S	0	0
			3769	2368	658	717	26		
5	G	462	Total	C	N	O	S	0	0
			3518	2210	621	664	23		

- Molecule 6 is a protein called T-complex protein 1 subunit theta.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	q	524	Total	C	N	O	S	0	0
			3978	2508	677	767	26		
6	Q	523	Total	C	N	O	S	0	0
			3974	2506	676	766	26		

- Molecule 7 is a protein called T-complex protein 1 subunit zeta.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	z	524	Total	C	N	O	S	0	0
			4030	2535	696	782	17		
7	Z	531	Total	C	N	O	S	0	0
			4086	2566	708	795	17		


- Molecule 8 is a protein called T-complex protein 1 subunit eta.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	h	507	Total	C	N	O	S	0	0
			3882	2447	664	752	19		
8	H	507	Total	C	N	O	S	0	0
			3882	2447	664	752	19		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-complex protein 1 subunit alpha




- Molecule 1: T-complex protein 1 subunit alpha







• Molecule 2: T-complex protein 1 subunit beta






• Molecule 3: T-complex protein 1 subunit delta







• Molecule 3: T-complex protein 1 subunit delta







• Molecule 4: T-complex protein 1 subunit epsilon




• Molecule 4: T-complex protein 1 subunit epsilon




GLY
GLY
PRO
ARG
ALA
PRO
GLN
GLY
PRO
ARG
GLY
ASN
TRP
ASP
GLN
GLU
ASP

• Molecule 7: T-complex protein 1 subunit zeta



• Molecule 7: T-complex protein 1 subunit zeta



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	54639	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	38	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	3.607	Depositor
Minimum map value	-2.255	Depositor
Average map value	0.027	Depositor
Map value standard deviation	0.214	Depositor
Recommended contour level	0.85	Depositor
Map size (\AA)	258.328, 258.328, 258.328	wwPDB
Map dimensions	196, 196, 196	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.318, 1.318, 1.318	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3898	0.41	0/5257
1	a	0.23	0/3884	0.40	0/5239
2	B	0.23	0/3862	0.39	0/5204
2	b	0.23	0/3849	0.39	0/5186
3	D	0.23	0/3279	0.39	0/4417
3	d	0.24	1/3687 (0.0%)	0.39	0/4967
4	E	0.23	0/3830	0.40	0/5158
4	e	0.23	0/3885	0.40	0/5228
5	G	0.23	0/3559	0.39	0/4805
5	g	0.23	0/3813	0.38	0/5150
6	Q	0.24	0/4020	0.41	0/5426
6	q	0.24	0/4024	0.40	0/5431
7	Z	0.23	0/4138	0.39	0/5590
7	z	0.23	0/4081	0.41	0/5513
8	H	0.23	0/3931	0.38	0/5304
8	h	0.24	0/3931	0.39	0/5304
All	All	0.23	1/61671 (0.0%)	0.40	0/83179

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	d	372	ARG	C-N	5.42	1.44	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	509/559 (91%)	481 (94%)	28 (6%)	0	100	100
1	a	507/559 (91%)	483 (95%)	23 (4%)	1 (0%)	47	81
2	B	502/527 (95%)	479 (95%)	23 (5%)	0	100	100
2	b	500/527 (95%)	475 (95%)	25 (5%)	0	100	100
3	D	419/528 (79%)	400 (96%)	18 (4%)	1 (0%)	47	81
3	d	474/528 (90%)	452 (95%)	21 (4%)	1 (0%)	47	81
4	E	491/562 (87%)	467 (95%)	24 (5%)	0	100	100
4	e	497/562 (88%)	473 (95%)	24 (5%)	0	100	100
5	G	456/534 (85%)	440 (96%)	16 (4%)	0	100	100
5	g	489/534 (92%)	469 (96%)	19 (4%)	1 (0%)	47	81
6	Q	521/568 (92%)	500 (96%)	21 (4%)	0	100	100
6	q	522/568 (92%)	498 (95%)	24 (5%)	0	100	100
7	Z	529/546 (97%)	510 (96%)	19 (4%)	0	100	100
7	z	520/546 (95%)	498 (96%)	22 (4%)	0	100	100
8	H	505/550 (92%)	479 (95%)	26 (5%)	0	100	100
8	h	505/550 (92%)	484 (96%)	21 (4%)	0	100	100
All	All	7946/8748 (91%)	7588 (96%)	354 (4%)	4 (0%)	54	85

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	g	195	ASN
3	d	213	VAL
1	a	203	VAL
3	D	213	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	428/471 (91%)	428 (100%)	0	100	100
1	a	427/471 (91%)	427 (100%)	0	100	100
2	B	421/441 (96%)	421 (100%)	0	100	100
2	b	420/441 (95%)	420 (100%)	0	100	100
3	D	365/453 (81%)	365 (100%)	0	100	100
3	d	409/453 (90%)	409 (100%)	0	100	100
4	E	422/483 (87%)	422 (100%)	0	100	100
4	e	428/483 (89%)	428 (100%)	0	100	100
5	G	387/455 (85%)	387 (100%)	0	100	100
5	g	417/455 (92%)	417 (100%)	0	100	100
6	Q	439/473 (93%)	439 (100%)	0	100	100
6	q	439/473 (93%)	439 (100%)	0	100	100
7	Z	448/463 (97%)	448 (100%)	0	100	100
7	z	442/463 (96%)	442 (100%)	0	100	100
8	H	422/454 (93%)	422 (100%)	0	100	100
8	h	422/454 (93%)	422 (100%)	0	100	100
All	All	6736/7386 (91%)	6736 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	a	88	GLN
1	a	89	GLN
1	a	90	GLN
1	a	250	ASN
2	b	142	ASN
2	b	222	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	b	286	ASN
2	b	393	GLN
2	B	84	GLN
3	d	240	GLN
3	d	289	GLN
3	D	284	ASN
3	D	289	GLN
4	e	349	GLN
4	e	397	GLN
4	e	432	ASN
4	e	509	ASN
4	e	536	GLN
4	E	241	ASN
4	E	536	GLN
5	G	470	GLN
6	Q	27	GLN
6	Q	66	ASN
6	Q	91	GLN
6	Q	92	GLN
6	Q	129	GLN
7	z	22	ASN
7	z	385	HIS
7	Z	185	GLN
7	Z	356	GLN
7	Z	430	ASN
8	h	25	GLN
8	h	172	HIS
8	H	231	GLN
8	H	435	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-0756. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 98



Y Index: 98




Z Index: 98

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 58



Y Index: 66




Z Index: 84


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.85. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 304 nm³; this corresponds to an approximate mass of 275 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.228\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-0756 and PDB model 6KRD. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.85 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.85).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 49% of all backbone atoms, 34% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.85) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.3350	 0.1420
A	 0.3070	 0.1350
B	 0.2350	 0.0910
D	 0.3000	 0.1420
E	 0.3330	 0.1350
G	 0.3610	 0.1620
H	 0.2830	 0.1170
Q	 0.3750	 0.1660
Z	 0.4370	 0.1960
a	 0.3420	 0.1280
b	 0.2410	 0.1080
d	 0.3130	 0.1420
e	 0.3330	 0.1220
g	 0.3390	 0.1380
h	 0.3800	 0.1530
q	 0.3490	 0.1460
z	 0.4190	 0.1880

