



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 28, 2023 – 02:17 AM EDT

PDB ID : 3KS8  
Title : Crystal structure of Reston ebolavirus VP35 RNA binding domain in complex with 18bp dsRNA  
Authors : Kimberlin, C.R.; Bornholdt, Z.A.; Li, S.; Woods, V.L.; Macrae, I.J.; Sapphire, E.O.  
Deposited on : 2009-11-20  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)
RNA backbone	3102	1174 (2.80-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	18	 50% 22% 22% 6%
2	F	18	 61% 33% 6%
3	A	184	 56% 9% 34%
3	B	184	 52% 14% 33%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	184	 <p>A horizontal bar chart showing the quality of chain C. The bar is divided into three segments: green (55%), yellow (10%), and grey (34%). A small red square is located at the beginning of the bar.</p>
3	D	184	 <p>A horizontal bar chart showing the quality of chain D. The bar is divided into three segments: green (53%), yellow (13%), and grey (33%). A small red square is located at the beginning of the bar.</p>

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4757 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(\*AP\*GP\*AP\*AP\*GP\*GP\*AP\*GP\*GP\*GP\*AP\*GP\*GP\*GP\*AP\*GP\*GP\*GP\*AP\*GP\*GP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	E	18	404	180	90	117	17	0	0	0

- Molecule 2 is a RNA chain called 5'-R(\*UP\*CP\*CP\*UP\*CP\*CP\*CP\*UP\*CP\*CP\*CP\*UP\*CP\*CP\*UP\*UP\*CP\*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	F	18	357	162	47	131	17	0	0	0

- Molecule 3 is a protein called Polymerase cofactor VP35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	122	952	609	169	168	6	29	0	0
3	B	124	971	623	171	171	6	30	0	0
3	C	122	952	609	169	168	6	28	0	0
3	D	124	971	623	171	171	6	29	0	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	MET	-	expression tag	UNP Q8JPY0
A	147	ALA	-	expression tag	UNP Q8JPY0
A	148	HIS	-	expression tag	UNP Q8JPY0
A	149	HIS	-	expression tag	UNP Q8JPY0
A	150	HIS	-	expression tag	UNP Q8JPY0
A	151	HIS	-	expression tag	UNP Q8JPY0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	152	HIS	-	expression tag	UNP Q8JPY0
A	153	HIS	-	expression tag	UNP Q8JPY0
A	154	VAL	-	expression tag	UNP Q8JPY0
A	155	ASP	-	expression tag	UNP Q8JPY0
A	156	ASP	-	expression tag	UNP Q8JPY0
A	157	ASP	-	expression tag	UNP Q8JPY0
A	158	ASP	-	expression tag	UNP Q8JPY0
A	159	LYS	-	expression tag	UNP Q8JPY0
B	146	MET	-	expression tag	UNP Q8JPY0
B	147	ALA	-	expression tag	UNP Q8JPY0
B	148	HIS	-	expression tag	UNP Q8JPY0
B	149	HIS	-	expression tag	UNP Q8JPY0
B	150	HIS	-	expression tag	UNP Q8JPY0
B	151	HIS	-	expression tag	UNP Q8JPY0
B	152	HIS	-	expression tag	UNP Q8JPY0
B	153	HIS	-	expression tag	UNP Q8JPY0
B	154	VAL	-	expression tag	UNP Q8JPY0
B	155	ASP	-	expression tag	UNP Q8JPY0
B	156	ASP	-	expression tag	UNP Q8JPY0
B	157	ASP	-	expression tag	UNP Q8JPY0
B	158	ASP	-	expression tag	UNP Q8JPY0
B	159	LYS	-	expression tag	UNP Q8JPY0
C	146	MET	-	expression tag	UNP Q8JPY0
C	147	ALA	-	expression tag	UNP Q8JPY0
C	148	HIS	-	expression tag	UNP Q8JPY0
C	149	HIS	-	expression tag	UNP Q8JPY0
C	150	HIS	-	expression tag	UNP Q8JPY0
C	151	HIS	-	expression tag	UNP Q8JPY0
C	152	HIS	-	expression tag	UNP Q8JPY0
C	153	HIS	-	expression tag	UNP Q8JPY0
C	154	VAL	-	expression tag	UNP Q8JPY0
C	155	ASP	-	expression tag	UNP Q8JPY0
C	156	ASP	-	expression tag	UNP Q8JPY0
C	157	ASP	-	expression tag	UNP Q8JPY0
C	158	ASP	-	expression tag	UNP Q8JPY0
C	159	LYS	-	expression tag	UNP Q8JPY0
D	146	MET	-	expression tag	UNP Q8JPY0
D	147	ALA	-	expression tag	UNP Q8JPY0
D	148	HIS	-	expression tag	UNP Q8JPY0
D	149	HIS	-	expression tag	UNP Q8JPY0
D	150	HIS	-	expression tag	UNP Q8JPY0
D	151	HIS	-	expression tag	UNP Q8JPY0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	152	HIS	-	expression tag	UNP Q8JPY0
D	153	HIS	-	expression tag	UNP Q8JPY0
D	154	VAL	-	expression tag	UNP Q8JPY0
D	155	ASP	-	expression tag	UNP Q8JPY0
D	156	ASP	-	expression tag	UNP Q8JPY0
D	157	ASP	-	expression tag	UNP Q8JPY0
D	158	ASP	-	expression tag	UNP Q8JPY0
D	159	LYS	-	expression tag	UNP Q8JPY0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	13	Total O 13 13	0	0
4	F	7	Total O 7 7	0	0
4	A	32	Total O 32 32	0	0
4	B	30	Total O 30 30	0	0
4	C	33	Total O 33 33	0	0
4	D	35	Total O 35 35	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-R(\*AP\*GP\*AP\*AP\*GP\*GP\*AP\*GP\*GP\*GP\*AP\*GP\*GP\*GP\*AP\*GP\*GP\*A)-3'

Chain E: 




- Molecule 2: 5'-R(\*UP\*CP\*CP\*UP\*CP\*CP\*CP\*UP\*CP\*CP\*CP\*UP\*CP\*CP\*UP\*UP\*CP\*U)-3'

Chain F: 



- Molecule 3: Polymerase cofactor VP35

Chain A: 



- Molecule 3: Polymerase cofactor VP35

Chain B: 



- Molecule 3: Polymerase cofactor VP35

Chain C: 





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.69Å 85.69Å 108.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.12 – 2.40 37.10 – 2.29	Depositor EDS
% Data completeness (in resolution range)	84.9 (35.12-2.40) 75.0 (37.10-2.29)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.27 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_222)	Depositor
R, $R_{free}$	0.198 , 0.239 0.196 , 0.236	Depositor DCC
$R_{free}$ test set	1532 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for -h,-k,l 0.467 for h,-h-k,-l 0.022 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4757	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	E	0.89	0/457	1.66	6/716 (0.8%)
2	F	0.77	0/392	1.60	1/604 (0.2%)
3	A	0.38	0/974	0.52	0/1317
3	B	0.36	0/995	0.53	0/1346
3	C	0.38	0/974	0.54	0/1317
3	D	0.36	0/995	0.53	0/1346
All	All	0.49	0/4787	0.87	7/6646 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	14	C	C4'-C3'-C2'	-6.59	96.01	102.60
1	E	16	G	N3-C4-N9	-5.54	122.67	126.00
1	E	9	G	C4'-C3'-C2'	-5.32	97.28	102.60
1	E	13	G	N3-C4-N9	-5.32	122.81	126.00
1	E	12	G	C4'-C3'-C2'	-5.13	97.47	102.60
1	E	16	G	N3-C4-C5	5.07	131.14	128.60
1	E	9	G	C1'-O4'-C4'	-5.04	105.87	109.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	404	0	200	6	0
2	F	357	0	193	5	0
3	A	952	0	984	13	1
3	B	971	0	1001	23	0
3	C	952	0	984	11	1
3	D	971	0	1001	22	0
4	A	32	0	0	0	0
4	B	30	0	0	1	0
4	C	33	0	0	0	0
4	D	35	0	0	0	0
4	E	13	0	0	3	0
4	F	7	0	0	0	0
All	All	4757	0	4363	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:268:GLN:HE21	3:B:268:GLN:HA	1.46	0.79
3:D:268:GLN:HE21	3:D:268:GLN:HA	1.50	0.75
1:E:12:G:N7	4:E:44:HOH:O	2.22	0.73
3:A:287:ARG:CG	3:A:287:ARG:HH11	2.01	0.73
3:C:311:ARG:O	3:C:328:LYS:HE3	1.94	0.68
3:A:311:ARG:O	3:A:328:LYS:HE3	1.94	0.67
3:A:287:ARG:HH11	3:A:287:ARG:HG3	1.59	0.65
1:E:15:A:N7	4:E:113:HOH:O	2.30	0.64
2:F:8:U:H6	2:F:8:U:H5''	1.65	0.62
3:D:263:GLN:NE2	3:D:263:GLN:H	1.98	0.62
3:B:263:GLN:H	3:B:263:GLN:NE2	1.98	0.61
3:B:210:ALA:HB2	3:B:240:LYS:CG	2.32	0.60
3:D:210:ALA:HB2	3:D:240:LYS:CG	2.32	0.60
3:D:283:ILE:HD12	3:D:283:ILE:N	2.17	0.59
3:B:283:ILE:N	3:B:283:ILE:HD12	2.18	0.59
3:B:218:TYR:CE1	3:B:229:HIS:HD2	2.21	0.58
3:A:287:ARG:HB3	3:A:291:ASP:OD2	2.04	0.58
3:D:218:TYR:CE1	3:D:229:HIS:HD2	2.22	0.58
3:C:287:ARG:HB3	3:C:291:ASP:OD2	2.04	0.57
3:A:214:LYS:HE3	3:A:218:TYR:CZ	2.43	0.53
3:A:289:ARG:CA	3:A:319:MET:HE2	2.39	0.53
1:E:14:G:N7	4:E:78:HOH:O	2.34	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:214:LYS:HE3	3:C:218:TYR:CZ	2.44	0.53
3:B:268:GLN:HA	3:B:268:GLN:NE2	2.21	0.52
2:F:8:U:H5''	2:F:8:U:C6	2.45	0.50
3:D:213:LEU:HD23	3:D:245:LEU:HD22	1.92	0.50
3:C:289:ARG:CA	3:C:319:MET:HE2	2.42	0.50
3:D:268:GLN:HA	3:D:268:GLN:NE2	2.24	0.50
3:A:321:ASP:OD1	3:A:323:LYS:HB2	2.12	0.49
3:B:321:ASP:OD2	3:B:323:LYS:HB2	2.12	0.48
3:A:287:ARG:CG	3:A:287:ARG:NH1	2.70	0.48
3:A:228:PHE:O	3:A:232:VAL:HG23	2.13	0.48
3:D:264:CYS:O	3:D:268:GLN:HG2	2.13	0.48
3:B:218:TYR:CD1	3:B:229:HIS:HD2	2.31	0.48
3:B:213:LEU:HD23	3:B:245:LEU:HD22	1.95	0.47
3:D:218:TYR:CD1	3:D:229:HIS:HD2	2.31	0.47
3:D:321:ASP:OD2	3:D:323:LYS:HB2	2.13	0.47
3:C:321:ASP:OD1	3:C:323:LYS:HB2	2.15	0.47
3:B:210:ALA:CB	3:B:240:LYS:HD2	2.45	0.47
3:D:210:ALA:CB	3:D:240:LYS:HD2	2.45	0.47
3:B:264:CYS:O	3:B:268:GLN:HG2	2.13	0.47
3:A:209:SER:HB3	3:A:212:ASP:OD2	2.15	0.46
3:A:289:ARG:HA	3:A:319:MET:HE2	1.96	0.46
3:C:209:SER:HB3	3:C:212:ASP:OD2	2.15	0.46
3:D:283:ILE:N	3:D:283:ILE:CD1	2.78	0.46
1:E:13:G:H1	2:F:6:C:H42	1.61	0.46
3:B:283:ILE:N	3:B:283:ILE:CD1	2.79	0.46
3:C:228:PHE:O	3:C:232:VAL:HG23	2.15	0.46
3:B:210:ALA:HB2	3:B:240:LYS:HG3	1.99	0.45
3:C:221:LEU:HA	3:C:222:PRO:HD3	1.87	0.44
1:E:7:A:C2	2:F:13:C:C2	3.06	0.44
3:A:221:LEU:HA	3:A:222:PRO:HD3	1.87	0.44
2:F:1:U:H1'	3:D:263:GLN:HB2	1.99	0.43
3:D:210:ALA:HB2	3:D:240:LYS:HG3	2.00	0.43
3:B:214:LYS:O	3:B:218:TYR:HB2	2.18	0.43
3:D:214:LYS:HG3	3:D:218:TYR:CE2	2.54	0.43
3:C:289:ARG:HA	3:C:319:MET:HE2	2.00	0.42
3:A:289:ARG:HA	3:A:319:MET:CE	2.50	0.42
3:D:212:ASP:O	3:D:215:GLU:HB3	2.20	0.42
3:B:285:HIS:CE1	3:B:318:LYS:HD3	2.55	0.42
3:D:214:LYS:O	3:D:218:TYR:HB2	2.19	0.42
3:D:285:HIS:CE1	3:D:318:LYS:HD3	2.55	0.42
3:B:218:TYR:CE1	3:B:229:HIS:CD2	3.05	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:273:VAL:HA	3:D:274:PRO:HD3	1.81	0.42
3:B:223:GLY:CA	4:B:105:HOH:O	2.67	0.41
3:B:210:ALA:HB2	3:B:240:LYS:HD2	2.02	0.41
3:B:273:VAL:HA	3:B:274:PRO:HD3	1.81	0.41
3:D:210:ALA:HB2	3:D:240:LYS:HD2	2.02	0.41
3:B:212:ASP:O	3:B:215:GLU:HB3	2.21	0.41
3:D:218:TYR:CE1	3:D:229:HIS:CD2	3.06	0.41
1:E:17:G:O2'	1:E:18:A:H5'	2.20	0.41
3:C:208:ILE:HD12	3:C:209:SER:H	1.86	0.41
3:D:286:ILE:O	3:D:319:MET:HA	2.21	0.41
3:C:289:ARG:HA	3:C:319:MET:CE	2.51	0.41
3:B:222:PRO:O	3:B:229:HIS:HE1	2.04	0.40
3:B:214:LYS:HG3	3:B:218:TYR:CE2	2.56	0.40
3:B:228:PHE:O	3:B:232:VAL:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:278:ASP:OD2	3:C:278:ASP:OD2[3_454]	1.84	0.36

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	120/184 (65%)	117 (98%)	3 (2%)	0	100	100
3	B	122/184 (66%)	117 (96%)	5 (4%)	0	100	100
3	C	120/184 (65%)	117 (98%)	3 (2%)	0	100	100
3	D	122/184 (66%)	116 (95%)	6 (5%)	0	100	100
All	All	484/736 (66%)	467 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	106/160 (66%)	101 (95%)	5 (5%)	26	42
3	B	108/160 (68%)	100 (93%)	8 (7%)	13	22
3	C	106/160 (66%)	102 (96%)	4 (4%)	33	51
3	D	108/160 (68%)	101 (94%)	7 (6%)	17	27
All	All	428/640 (67%)	404 (94%)	24 (6%)	21	34

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	255	SER
3	A	277	GLN
3	A	287	ARG
3	A	319	MET
3	A	321	ASP
3	B	211	LYS
3	B	237	LYS
3	B	263	GLN
3	B	268	GLN
3	B	277	GLN
3	B	283	ILE
3	B	301	ARG
3	B	311	ARG
3	C	255	SER
3	C	277	GLN
3	C	287	ARG
3	C	321	ASP
3	D	211	LYS
3	D	237	LYS
3	D	263	GLN
3	D	268	GLN
3	D	277	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	D	301	ARG
3	D	311	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	253	GLN
3	B	263	GLN
3	B	268	GLN
3	B	285	HIS
3	C	253	GLN
3	D	263	GLN
3	D	268	GLN
3	D	285	HIS

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	E	17/18 (94%)	4 (23%)	1 (5%)
2	F	17/18 (94%)	3 (17%)	0
All	All	34/36 (94%)	7 (20%)	1 (2%)

All (7) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	E	9	G
1	E	13	G
1	E	14	G
1	E	15	A
2	F	9	C
2	F	14	C
2	F	15	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	E	9	G

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	18/18 (100%)	-1.02	0 <a href="#">100</a> <a href="#">100</a>	30, 50, 77, 80	0
2	F	18/18 (100%)	-0.90	0 <a href="#">100</a> <a href="#">100</a>	27, 51, 86, 88	0
3	A	122/184 (66%)	-0.59	1 (0%) <a href="#">86</a> <a href="#">84</a>	21, 35, 67, 92	7 (5%)
3	B	124/184 (67%)	-0.46	0 <a href="#">100</a> <a href="#">100</a>	21, 38, 61, 97	7 (5%)
3	C	122/184 (66%)	-0.59	0 <a href="#">100</a> <a href="#">100</a>	21, 35, 67, 82	7 (5%)
3	D	124/184 (67%)	-0.34	1 (0%) <a href="#">86</a> <a href="#">84</a>	21, 38, 62, 106	7 (5%)
All	All	528/772 (68%)	-0.53	2 (0%) <a href="#">92</a> <a href="#">91</a>	21, 38, 68, 106	28 (5%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	322	GLY	2.1
3	D	207	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.