



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 04:55 AM EDT

PDB ID : 4KWD
Title : Crystal structure of *Aspergillus terreus* aristolochene synthase complexed with (1R,8R,9aS)-1,9a-dimethyl-8-(prop-1-en-2-yl)decahydroquinolizin-5-ium
Authors : Chen, M.; Al-lami, N.; Janvier, M.; D'Antonio, E.L.; Faraldos, J.A.; Cane, D.E.; Allemann, R.K.; Christianson, D.W.
Deposited on : 2013-05-23
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	 89% 8% ..
1	B	314	 91% 6% .
1	C	314	 % 86% 11% .
1	D	314	 14% 80% 14% ...

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aristolochene synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	304	2459	1570	415	458	16	0	1	0
1	B	304	2459	1570	415	458	16	0	1	0
1	C	304	2459	1570	415	458	16	0	1	0
1	D	304	2454	1566	415	458	15	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q9UR08
A	2	HIS	-	expression tag	UNP Q9UR08
A	3	HIS	-	expression tag	UNP Q9UR08
A	4	HIS	-	expression tag	UNP Q9UR08
A	5	HIS	-	expression tag	UNP Q9UR08
A	6	HIS	-	expression tag	UNP Q9UR08
A	7	HIS	-	expression tag	UNP Q9UR08
B	1	MET	-	expression tag	UNP Q9UR08
B	2	HIS	-	expression tag	UNP Q9UR08
B	3	HIS	-	expression tag	UNP Q9UR08
B	4	HIS	-	expression tag	UNP Q9UR08
B	5	HIS	-	expression tag	UNP Q9UR08
B	6	HIS	-	expression tag	UNP Q9UR08
B	7	HIS	-	expression tag	UNP Q9UR08
C	1	MET	-	expression tag	UNP Q9UR08
C	2	HIS	-	expression tag	UNP Q9UR08
C	3	HIS	-	expression tag	UNP Q9UR08
C	4	HIS	-	expression tag	UNP Q9UR08
C	5	HIS	-	expression tag	UNP Q9UR08
C	6	HIS	-	expression tag	UNP Q9UR08
C	7	HIS	-	expression tag	UNP Q9UR08

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MET	-	expression tag	UNP Q9UR08
D	2	HIS	-	expression tag	UNP Q9UR08
D	3	HIS	-	expression tag	UNP Q9UR08
D	4	HIS	-	expression tag	UNP Q9UR08
D	5	HIS	-	expression tag	UNP Q9UR08
D	6	HIS	-	expression tag	UNP Q9UR08
D	7	HIS	-	expression tag	UNP Q9UR08

- Molecule 2 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 9 7 2	0	0
2	B	1	Total O P 9 7 2	0	0
2	C	1	Total O P 9 7 2	0	0
2	D	1	Total O P 9 7 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Mg 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Mg	0	0
			3	3		
3	C	3	Total	Mg	0	0
			3	3		
3	D	3	Total	Mg	0	0
			3	3		

- Molecule 4 is (1R,5R,8R,9aS)-1,9a-dimethyl-8-(prop-1-en-2-yl)octahydro-2H-quinolizinium (three-letter code: JF2) (formula: C₁₄H₂₆N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			15	14	1		
4	B	1	Total	C	N	0	0
			15	14	1		
4	C	1	Total	C	N	0	0
			15	14	1		
4	D	1	Total	C	N	0	0
			15	14	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	336	Total O 336 336	0	0
6	B	353	Total O 353 353	0	0
6	C	229	Total O 229 229	0	0
6	D	131	Total O 131 131	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aristolochene synthase

Chain A:  89% 8% ..





- Molecule 1: Aristolochene synthase

Chain B:  91% 6% .





- Molecule 1: Aristolochene synthase


Chain C:  86% 11% .



- Molecule 1: Aristolochene synthase

Chain D:  14% 80% 14% ...





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.86Å 123.86Å 202.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.37 – 1.86 47.37 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.37-1.86) 99.8 (47.37-1.86)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 1.86Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1370)	Depositor
R, R_{free}	0.193 , 0.226 0.194 , 0.228	Depositor DCC
R_{free} test set	7601 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtrriage
Anisotropy	0.252	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.012 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11000	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, POP, JF2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/2517	0.80	0/3406
1	B	0.81	0/2517	0.83	3/3406 (0.1%)
1	C	0.68	0/2517	0.74	1/3406 (0.0%)
1	D	0.58	0/2509	0.68	3/3396 (0.1%)
All	All	0.72	0/10060	0.76	7/13614 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	169	ARG	NE-CZ-NH2	-10.01	115.30	120.30
1	C	107	ARG	NE-CZ-NH1	-8.70	115.95	120.30
1	B	183	ARG	NE-CZ-NH2	-8.68	115.96	120.30
1	D	169	ARG	NE-CZ-NH1	8.04	124.32	120.30
1	D	57	VAL	CB-CA-C	-6.42	99.20	111.40
1	B	183	ARG	NE-CZ-NH1	6.33	123.47	120.30
1	B	56	ARG	NE-CZ-NH2	-5.88	117.36	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2459	0	2434	9	0
1	B	2459	0	2434	7	0
1	C	2459	0	2434	13	0
1	D	2454	0	2425	34	0
2	A	9	0	0	0	0
2	B	9	0	0	0	0
2	C	9	0	0	1	0
2	D	9	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
4	A	15	0	26	0	0
4	B	15	0	26	1	0
4	C	15	0	26	1	0
4	D	15	0	26	0	0
5	B	6	0	8	0	0
5	C	6	0	8	1	0
6	A	336	0	0	3	0
6	B	353	0	0	5	0
6	C	229	0	0	2	0
6	D	131	0	0	5	0
All	All	11000	0	9847	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:15:PHE:HE2	1:D:296:MET:HE2	1.50	0.76
1:D:135:GLU:HG2	1:D:189:LYS:HE2	1.72	0.72
1:D:165:TYR:CZ	1:D:169:ARG:HG3	2.31	0.66
1:D:36:GLN:HG2	1:D:37:HIS:CD2	2.31	0.65
4:B:705:JF2:H6	6:B:1153:HOH:O	1.96	0.64
1:D:201:ILE:O	6:D:570:HOH:O	2.14	0.64
1:D:15:PHE:CE2	1:D:296:MET:HE2	2.33	0.64
1:A:197:ARG:HD3	1:A:280:LEU:HB3	1.82	0.62
1:A:35:LEU:O	1:A:46:ARG:NH2	2.32	0.61
1:B:154:ARG:HD2	6:B:949:HOH:O	2.01	0.59
1:C:18:LEU:HD22	1:C:63:PRO:HB2	1.85	0.57
1:B:236:CYS:O	6:B:1126:HOH:O	2.17	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:64:LYS:O	1:D:186:MET:HG2	2.06	0.55
1:C:35:LEU:O	1:C:46:ARG:NH2	2.39	0.54
1:A:222:LEU:HD22	1:A:226:LYS:HE3	1.90	0.54
1:D:57:VAL:HG22	1:D:302:TRP:CE2	2.43	0.54
1:D:69:ARG:NH2	1:D:133:ASP:OD1	2.33	0.53
1:D:112:PRO:HD3	1:D:123:TYR:CD1	2.44	0.53
1:D:160:MET:HE1	1:D:165:TYR:HD2	1.74	0.52
1:B:183:ARG:NH2	6:B:1116:HOH:O	2.41	0.52
1:C:196:GLN:HA	1:C:199:ARG:HD3	1.91	0.52
1:B:200:GLU:OE1	1:B:267:ARG:NH2	2.43	0.51
1:C:183:ARG:NH1	6:C:568:HOH:O	2.44	0.51
1:D:57:VAL:HG13	1:D:302:TRP:CG	2.46	0.51
1:A:267:ARG:NH1	6:A:670:HOH:O	2.42	0.50
1:A:25:GLU:HB2	6:A:614:HOH:O	2.11	0.50
1:A:219:GLU:HG2	6:A:653:HOH:O	2.12	0.50
1:D:307:LEU:O	1:D:311:VAL:HG22	2.12	0.49
1:D:75:ARG:HB2	1:D:125:LEU:HD21	1.95	0.48
1:A:160:MET:HE1	1:A:236:CYS:SG	2.54	0.47
1:A:57:VAL:HB	1:A:302:TRP:CE2	2.50	0.47
1:B:15:PHE:HB3	1:B:297:SER:HB3	1.97	0.47
1:D:46:ARG:HD2	6:D:541:HOH:O	2.15	0.46
1:D:66:LEU:HB2	1:D:69:ARG:HD3	1.98	0.46
1:C:213:ASN:O	1:C:217:SER:OG	2.34	0.46
1:D:38:TRP:HB3	1:D:40:PHE:CE1	2.51	0.45
1:D:93:GLU:HG3	1:D:154:ARG:HH22	1.81	0.45
1:D:43:GLU:HG3	1:D:47:LYS:HE2	1.99	0.45
1:D:175:LYS:HG3	6:D:551:HOH:O	2.17	0.45
1:D:15:PHE:HE2	1:D:296:MET:CE	2.27	0.45
1:D:62:PHE:CE1	1:D:181:LEU:HD13	2.52	0.45
1:C:15:PHE:HB3	1:C:297:SER:HB3	1.98	0.45
1:D:222:LEU:HD22	1:D:226:LYS:HE3	2.00	0.44
1:D:266:LEU:HD13	6:D:510:HOH:O	2.17	0.44
1:A:57:VAL:HB	1:A:302:TRP:CD2	2.53	0.43
1:C:149:ARG:NH1	5:C:406:GOL:H11	2.33	0.43
1:C:160:MET:HE1	1:C:165:TYR:HD2	1.83	0.43
1:C:28:LYS:HA	1:C:28:LYS:HD3	1.81	0.43
1:C:46:ARG:HD2	6:C:619:HOH:O	2.18	0.43
1:D:165:TYR:CE2	1:D:169:ARG:HG3	2.53	0.43
1:D:160:MET:HE3	1:D:165:TYR:HB2	2.01	0.43
1:C:169:ARG:HG2	1:C:210:SER:HB3	2.00	0.43
2:C:401:POP:O	4:C:405:JF2:H17	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:CYS:HA	6:B:1023:HOH:O	2.18	0.42
1:C:40:PHE:HA	1:C:41:PRO:HD3	1.76	0.42
1:D:106:SER:HA	1:D:126:TRP:CD1	2.55	0.42
1:B:57:VAL:HB	1:B:302:TRP:CD2	2.54	0.42
1:D:145:PHE:O	1:D:149:ARG:HD2	2.20	0.41
1:C:261:CYS:O	1:C:265:GLU:HG3	2.21	0.41
1:D:57:VAL:HG22	1:D:302:TRP:NE1	2.36	0.41
1:D:165:TYR:OH	1:D:169:ARG:HD2	2.21	0.40
1:D:296:MET:HB3	1:D:296:MET:HE3	1.69	0.40
1:D:303:SER:OG	6:D:555:HOH:O	2.22	0.40
1:D:160:MET:CE	1:D:165:TYR:HD2	2.33	0.40
1:D:194:GLU:OE1	1:D:282:THR:OG1	2.27	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/314 (96%)	300 (99%)	3 (1%)	0	100	100
1	B	303/314 (96%)	300 (99%)	3 (1%)	0	100	100
1	C	303/314 (96%)	299 (99%)	4 (1%)	0	100	100
1	D	302/314 (96%)	293 (97%)	8 (3%)	1 (0%)	41	26
All	All	1211/1256 (96%)	1192 (98%)	18 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	133	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	267/276 (97%)	252 (94%)	15 (6%)	21 7
1	B	267/276 (97%)	258 (97%)	9 (3%)	37 19
1	C	267/276 (97%)	254 (95%)	13 (5%)	25 9
1	D	266/276 (96%)	250 (94%)	16 (6%)	19 5
All	All	1067/1104 (97%)	1014 (95%)	53 (5%)	24 9

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LEU
1	A	25	GLU
1	A	125	LEU
1	A	134	ARG
1	A	173	VAL
1	A	177	LEU
1	A	181	LEU
1	A	196	GLN
1	A	222	LEU
1	A	237	THR
1	A	258	PHE
1	A	266	LEU
1	A	285	LEU
1	A	301	LEU
1	A	307	LEU
1	B	181	LEU
1	B	222	LEU
1	B	237	THR
1	B	240	GLN
1	B	258	PHE
1	B	266	LEU
1	B	285	LEU
1	B	301	LEU
1	B	311	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	28	LYS
1	C	111	LEU
1	C	125	LEU
1	C	181	LEU
1	C	222	LEU
1	C	237	THR
1	C	240	GLN
1	C	247	ASP
1	C	258	PHE
1	C	266	LEU
1	C	267	ARG
1	C	285	LEU
1	C	301	LEU
1	D	36	GLN
1	D	44	LYS
1	D	57	VAL
1	D	125	LEU
1	D	134	ARG
1	D	162	LEU
1	D	167	GLU
1	D	169	ARG
1	D	177	LEU
1	D	181	LEU
1	D	185	SER
1	D	222	LEU
1	D	258	PHE
1	D	266	LEU
1	D	275	LEU
1	D	301	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	196	GLN
1	C	39	ASN
1	D	132	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 12 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	POP	A	401	3	6,8,8	0.85	0	13,13,13	1.08	1 (7%)
4	JF2	D	405	-	14,16,16	0.73	1 (7%)	20,24,24	1.24	1 (5%)
4	JF2	B	705	-	14,16,16	0.61	0	20,24,24	2.06	6 (30%)
4	JF2	C	405	-	14,16,16	0.77	1 (7%)	20,24,24	1.49	4 (20%)
2	POP	B	704	3	6,8,8	0.98	0	13,13,13	1.20	1 (7%)
5	GOL	C	406	-	5,5,5	0.21	0	5,5,5	0.71	0
5	GOL	B	706	-	5,5,5	0.25	0	5,5,5	0.74	0
4	JF2	A	405	-	14,16,16	0.85	1 (7%)	20,24,24	2.04	6 (30%)
2	POP	C	401	3	6,8,8	0.91	0	13,13,13	1.50	2 (15%)
2	POP	D	401	3	6,8,8	0.49	0	13,13,13	1.54	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	POP	A	401	3	-	0/6/6/6	-
4	JF2	D	405	-	-	0/4/32/32	0/2/2/2
4	JF2	B	705	-	-	0/4/32/32	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	JF2	C	405	-	-	0/4/32/32	0/2/2/2
2	POP	B	704	3	-	0/6/6/6	-
5	GOL	C	406	-	-	2/4/4/4	-
5	GOL	B	706	-	-	0/4/4/4	-
4	JF2	A	405	-	-	0/4/32/32	0/2/2/2
2	POP	C	401	3	-	0/6/6/6	-
2	POP	D	401	3	-	0/6/6/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	405	JF2	CAO-CAM	-2.23	1.52	1.55
4	A	405	JF2	CAH-NAN	2.18	1.50	1.47
4	C	405	JF2	CAO-CAM	-2.12	1.52	1.55

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	705	JF2	CAI-NAN-CAH	-5.25	104.67	110.66
4	A	405	JF2	CAE-CAF-CAM	-4.25	106.11	112.71
4	A	405	JF2	CAF-CAE-CAH	4.08	116.57	110.85
4	B	705	JF2	CAD-CAO-CAJ	-3.46	105.15	110.67
2	C	401	POP	P2-O-P1	-3.44	121.02	132.83
2	D	401	POP	O6-P2-O5	3.20	119.85	107.64
2	B	704	POP	P2-O-P1	-3.14	122.06	132.83
4	A	405	JF2	CAO-CAJ-CAL	-3.14	108.43	114.42
4	C	405	JF2	CAD-CAO-CAM	-2.95	107.34	112.04
2	A	401	POP	P2-O-P1	-2.80	123.21	132.83
4	B	705	JF2	CAO-CAJ-CAL	-2.80	109.07	114.42
4	C	405	JF2	CAO-CAJ-CAL	-2.78	109.11	114.42
4	A	405	JF2	CAJ-CAL-CAK	-2.75	108.06	112.52
4	D	405	JF2	CAI-NAN-CAH	-2.70	107.58	110.66
4	B	705	JF2	CAJ-CAL-CAG	-2.66	106.57	109.72
4	B	705	JF2	CAG-CAL-CAK	-2.60	106.99	113.39
4	A	405	JF2	CAI-NAN-CAH	-2.52	107.79	110.66
4	C	405	JF2	CAE-CAF-CAM	-2.52	108.81	112.71
4	C	405	JF2	CAI-NAN-CAH	-2.49	107.82	110.66
2	D	401	POP	O3-P1-O2	2.30	116.42	107.64
2	C	401	POP	O6-P2-O4	2.11	118.93	110.68
4	A	405	JF2	CAJ-CAO-NAN	-2.06	105.34	108.65
4	B	705	JF2	CAE-CAF-CAM	-2.04	109.55	112.71

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	406	GOL	C1-C2-C3-O3
5	C	406	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	705	JF2	1	0
4	C	405	JF2	1	0
5	C	406	GOL	1	0
2	C	401	POP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/314 (96%)	-0.58	0 100 100	10, 17, 32, 50	0
1	B	304/314 (96%)	-0.59	0 100 100	10, 15, 27, 41	0
1	C	304/314 (96%)	-0.30	2 (0%) 87 88	13, 25, 44, 58	0
1	D	304/314 (96%)	0.67	43 (14%) 2 3	21, 38, 66, 96	0
All	All	1216/1256 (96%)	-0.20	45 (3%) 41 39	10, 23, 49, 96	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	280	LEU	5.9
1	D	277	ALA	4.8
1	D	195	LEU	4.6
1	D	279	GLY	4.3
1	D	18	LEU	3.6
1	D	275	LEU	3.6
1	D	188	LEU	3.4
1	D	126	TRP	3.4
1	D	19	CYS	3.4
1	D	72	PHE	3.2
1	D	285	LEU	3.1
1	D	287	ALA	3.1
1	D	197	ARG	3.1
1	D	196	GLN	2.8
1	D	193	SER	2.8
1	D	70	ILE	2.8
1	D	272	VAL	2.8
1	D	191	SER	2.7
1	D	23	VAL	2.7
1	C	311	VAL	2.6
1	D	61	TYR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	190	LEU	2.6
1	D	199	ARG	2.6
1	D	65	ALA	2.5
1	D	22	LEU	2.4
1	D	136	MET	2.4
1	D	56	ARG	2.4
1	D	184	PHE	2.4
1	D	129	MET	2.4
1	D	286	ALA	2.3
1	D	111	LEU	2.2
1	C	16	GLN	2.2
1	D	189	LYS	2.2
1	D	270	LEU	2.2
1	D	14	THR	2.2
1	D	288	TYR	2.2
1	D	16	GLN	2.2
1	D	200	GLU	2.1
1	D	125	LEU	2.1
1	D	284	GLY	2.1
1	D	283	PRO	2.1
1	D	21	PRO	2.1
1	D	294	TYR	2.1
1	D	179	ALA	2.0
1	D	276	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	JF2	D	405	15/15	0.85	0.19	36,45,53,54	0
3	MG	D	404	1/1	0.88	0.11	24,24,24,24	0
5	GOL	C	406	6/6	0.88	0.24	24,40,46,49	0
4	JF2	A	405	15/15	0.91	0.12	18,21,26,28	0
3	MG	D	403	1/1	0.93	0.08	24,24,24,24	0
4	JF2	B	705	15/15	0.93	0.11	18,22,25,27	0
4	JF2	C	405	15/15	0.94	0.11	25,31,35,37	0
3	MG	C	403	1/1	0.95	0.07	22,22,22,22	0
2	POP	D	401	9/9	0.95	0.11	23,28,28,29	0
3	MG	D	402	1/1	0.96	0.06	23,23,23,23	0
5	GOL	B	706	6/6	0.97	0.07	25,26,29,33	0
3	MG	C	404	1/1	0.97	0.07	22,22,22,22	0
3	MG	C	402	1/1	0.98	0.05	19,19,19,19	0
2	POP	C	401	9/9	0.98	0.07	18,19,22,24	0
3	MG	A	402	1/1	0.98	0.06	14,14,14,14	0
3	MG	A	403	1/1	0.98	0.07	13,13,13,13	0
2	POP	A	401	9/9	0.99	0.08	10,11,12,13	0
2	POP	B	704	9/9	0.99	0.09	8,10,10,13	0
3	MG	A	404	1/1	0.99	0.05	11,11,11,11	0
3	MG	B	701	1/1	1.00	0.10	12,12,12,12	0
3	MG	B	702	1/1	1.00	0.10	10,10,10,10	0
3	MG	B	703	1/1	1.00	0.08	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.