

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2023 – 12:25 PM EDT

PDB ID : 7KZE

Title : Substrate-dependent divergence of leukotriene A4 hydrolase aminopeptidase

activity

Authors: Lee, K.H.; Shim, Y.; Paige, M.; Noble, S.M.

Deposited on : 2020-12-10

Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	608	91%	9%
1	В	608	91%	9%
1	С	608	87%	13%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 15254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Leukotriene A-4 hydrolase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	600	Total	С	N	О	S	0	1	0
1	A	608	4865	3125	809	910	21	0	1	0
1	В	608	Total	С	N	О	S	1	1	0
1	Б	000	4865	3125	809	910	21	1		
1	С	607	Total	С	N	О	S	2	2	0
1		007	4860	3123	808	908	21	9		

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0


• Molecule 3 is 1-benzyl-4-methoxybenzene (three-letter code: N0Y) (formula: $C_{14}H_{14}O$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Λ	1	Total C O	0	0
$\begin{array}{c c} 3 & A \end{array}$	А	1	15 14 1	0	U
2	C	1	Total C O	0	0
J		1	15 14 1	0	U

 \bullet Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $\mathrm{C_6H_{14}O_4}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total C O 10 6 4	0	0

 \bullet Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	257	Total O 257 257	0	0
5	В	172	Total O 172 172	0	0
5	С	192	Total O 192 192	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Leukotriene A-4 hydrolase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32	Depositor
Cell constants	139.89Å 139.89Å 84.34Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.84 - 2.90	Depositor
Resolution (A)	53.84 - 2.90	EDS
% Data completeness	98.0 (53.84-2.90)	Depositor
(in resolution range)	94.9 (53.84-2.90)	EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.34 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
P. P.	0.195 , 0.209	Depositor
R, R_{free}	0.194 , 0.209	DCC
R_{free} test set	1957 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.32 \; , 20.4$	EDS
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.32$	Xtriage
	0.025 for -h,-k,l	
Estimated twinning fraction	0.035 for h,-h-k,-l	Xtriage
	0.021 for -k,-h,-l	
F_o, F_c correlation	0.90	EDS
Total number of atoms	15254	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, N0Y, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.23	0/4992	0.41	0/6786
1	В	0.21	0/4992	0.39	0/6786
1	С	0.22	0/4990	0.40	0/6783
All	All	0.22	0/14974	0.40	0/20355

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4865	0	4832	33	0
1	В	4865	0	4832	33	0
1	С	4860	0	4833	44	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
3	A	15	0	0	0	0
3	С	15	0	0	1	0
4	В	10	0	14	2	0
5	A	257	0	0	2	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	В	172	0	0	1	0
5	С	192	0	0	1	0
All	All	15254	0	14511	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$-$ overlap (\mathring{A})
1:A:212:ARG:HD2	5:A:889:HOH:O	1.35	1.23
3:C:702:N0Y:C15	5:C:968:HOH:O	2.27	0.82
1:A:118:LEU:HB2	1:A:130:TYR:HB3	1.86	0.58
1:C:114:ALA:HB2	1:C:139:HIS:HB3	1.86	0.57
1:C:118:LEU:HB2	1:C:130:TYR:HB3	1.90	0.54
1:B:137:ALA:HB2	4:B:702:PGE:H32	1.90	0.53
1:B:210:GLU:HB2	1:B:227:VAL:HG21	1.91	0.53
1:B:398:GLY:HA3	1:B:402:ILE:HD12	1.90	0.53
1:C:159:GLU:HG2	1:C:190:LYS:HG2	1.91	0.52
1:A:22:HIS:HB3	1:A:39:ALA:HB3	1.91	0.52
1:B:588:TYR:CZ	1:B:592:LYS:HD3	2.43	0.52
1:C:210:GLU:HB2	1:C:227:VAL:HG21	1.91	0.52
1:C:270:MET:HB3	1:C:277:PHE:HB2	1.92	0.52
1:A:306:VAL:O	1:A:306:VAL:HG13	2.10	0.51
1:B:270:MET:HB3	1:B:277:PHE:HB2	1.93	0.51
1:B:114:ALA:HB2	1:B:139:HIS:HB3	1.91	0.51
1:C:272:ASN:HB2	1:C:275:LEU:O	2.11	0.50
1:C:489:THR:HG22	1:C:492:LYS:HE3	1.94	0.50
1:C:398:GLY:HA3	1:C:402:ILE:HD12	1.94	0.50
1:A:487:ASN:ND2	1:A:488:ALA:H	2.10	0.50
1:A:15:VAL:HG13	1:A:46:GLU:OE2	2.12	0.49
1:A:365:LEU:HD13	1:A:385:LYS:HG2	1.94	0.49
1:B:3:ILE:HG13	1:B:420:THR:HG21	1.94	0.49
1:C:489:THR:HA	1:C:492:LYS:HG3	1.95	0.48
1:B:177:GLU:OE1	1:B:187:LYS:CE	2.62	0.48
1:B:309:LYS:HA	1:B:418:SER:HB3	1.94	0.48
1:C:135:CYS:HA	1:C:139:HIS:HB2	1.96	0.48
1:A:232:TYR:O	1:A:287:LYS:NZ	2.46	0.47
1:A:251:TYR:CZ	1:A:253:TRP:HB2	2.48	0.47
1:B:212:ARG:O	1:B:220:VAL:HG22	2.14	0.47
1:A:159:GLU:HG2	1:A:190:LYS:HG2	1.96	0.47



 $Continued\ from\ previous\ page...$

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:261:LEU:HD22	1:A:265:PHE:CD1	2.50	0.46
1:A:94:LEU:HD22	1:A:98:GLN:HG2	1.97	0.46
1:C:17:ARG:NH2	1:C:43:GLN:NE2	2.63	0.46
1:B:299:HIS:CD2	1:B:321:THR:HG21	2.50	0.46
1:A:258:LEU:HD23	1:A:276:THR:HB	1.98	0.46
1:B:301:TRP:O	1:B:306:VAL:HG12	2.16	0.46
1:C:22:HIS:HB3	1:C:39:ALA:HB3	1.96	0.46
1:C:564:MET:HG2	1:C:599:THR:HG23	1.97	0.46
1:A:135:CYS:HA	1:A:139:HIS:HB2	1.98	0.46
1:B:148:ASP:HA	1:B:199:CYS:SG	2.56	0.46
1:A:487:ASN:CG	1:A:488:ALA:N	2.70	0.46
1:B:251:TYR:CZ	1:B:253:TRP:HB2	2.51	0.46
1:C:258:LEU:HD23	1:C:276:THR:HB	1.97	0.45
1:A:205:VAL:HG11	1:A:259:LEU:HD22	1.99	0.45
1:C:154:LEU:HD12	1:C:154:LEU:O	2.15	0.45
1:C:205:VAL:HG11	1:C:259:LEU:HD22	1.98	0.45
1:C:511:PRO:HB3	1:C:544:GLN:O	2.15	0.45
1:B:22:HIS:HB3	1:B:39:ALA:HB3	1.98	0.45
1:B:177:GLU:OE1	1:B:187:LYS:NZ	2.47	0.45
1:C:19:LYS:HE2	1:C:99:GLU:OE2	2.16	0.45
1:B:232:TYR:O	1:B:287:LYS:NZ	2.46	0.45
1:C:263:PRO:O	1:C:562:GLY:N	2.40	0.45
1:B:118:LEU:HB2	1:B:130:TYR:HB3	1.98	0.45
1:B:255:GLN:O	1:B:274:CYS:HA	2.17	0.45
1:C:153:LYS:NZ	1:C:307:THR:OG1	2.34	0.45
1:B:137:ALA:HB1	1:B:311:TRP:CE2	2.52	0.44
1:B:374:PRO:HB3	4:B:702:PGE:H4	1.98	0.44
1:A:411:VAL:O	1:A:415:SER:OG	2.27	0.44
1:C:243:ILE:HG23	1:C:331:ARG:HD2	1.99	0.44
1:C:514:LEU:HG	1:C:518:LYS:HE3	1.98	0.44
1:A:511:PRO:HB3	1:A:544:GLN:O	2.17	0.44
1:B:265:PHE:HA	1:B:266:PRO:HD3	1.79	0.44
1:B:487:ASN:HB3	5:B:829:HOH:O	2.17	0.44
1:B:568:ARG:HB2	1:B:569:PRO:HD3	1.99	0.44
1:B:585:VAL:O	1:B:589:GLN:HG2	2.18	0.44
1:C:309:LYS:HA	1:C:418:SER:HB3	1.99	0.44
1:C:198:PRO:HD2	1:C:201:LEU:HD12	2.00	0.44
1:B:551:ILE:HD13	1:B:580:SER:HB3	2.00	0.44
1:C:568:ARG:HB2	1:C:569:PRO:HD3	1.99	0.44
1:C:17:ARG:NH2	1:C:43:GLN:HE22	2.15	0.43
1:C:365:LEU:HD23	1:C:448:LEU:HD23	1.98	0.43



 $Continued\ from\ previous\ page...$

Continued from prec		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:365:LEU:HD13	1:A:385:LYS:CG	2.49	0.43
1:B:3:ILE:CD1	1:B:420:THR:HG21	2.48	0.43
1:C:299:HIS:CD2	1:C:321:THR:HG21	2.52	0.43
1:A:271:GLU:OE1	5:A:801:HOH:O	2.20	0.43
1:A:578:ASP:OD1	1:A:579:LYS:N	2.51	0.43
1:B:135:CYS:HA	1:B:139:HIS:HB2	2.00	0.43
1:C:265:PHE:HA	1:C:266:PRO:HD3	1.81	0.43
1:C:457:LYS:HA	1:C:458:PRO:HD3	1.90	0.43
1:A:309:LYS:HA	1:A:418:SER:HB3	2.00	0.43
1:A:299:HIS:CD2	1:A:321:THR:HG21	2.53	0.42
1:C:578:ASP:OD1	1:C:579:LYS:N	2.52	0.42
1:C:120:PRO:O	1:C:126:LYS:HA	2.19	0.42
1:A:210:GLU:HB2	1:A:227:VAL:HG21	2.01	0.42
1:C:292:VAL:O	1:C:296:GLU:HG2	2.20	0.42
1:A:136:GLN:HA	1:A:137:ALA:HA	1.71	0.42
1:C:591:HIS:O	1:C:595:MET:HG3	2.20	0.42
1:A:270:MET:HB3	1:A:277:PHE:HB2	2.01	0.42
1:C:137:ALA:HB1	1:C:311:TRP:CE2	2.54	0.42
1:A:253:TRP:CZ2	1:A:304:ASN:HB3	2.55	0.42
1:B:136:GLN:HA	1:B:137:ALA:HA	1.72	0.42
1:C:17:ARG:HE	1:C:17:ARG:HB2	1.69	0.41
1:C:279:THR:HG23	1:C:282:LEU:HG	2.02	0.41
1:A:487:ASN:CG	1:A:488:ALA:H	2.23	0.41
1:C:154:LEU:HD12	1:C:154:LEU:C	2.41	0.41
1:A:263:PRO:O	1:A:562:GLY:N	2.40	0.41
1:A:457:LYS:HA	1:A:458:PRO:HD3	1.91	0.41
1:B:43:GLN:NE2	1:B:45:GLN:OE1	2.47	0.41
1:C:27:VAL:HG21	1:C:206:VAL:HG21	2.02	0.41
1:C:180:ASP:HB3	1:C:183:ASP:O	2.19	0.41
1:C:323:TYR:OH	1:C:394:GLU:OE1	2.30	0.41
1:C:519:ARG:O	1:C:523:VAL:HG22	2.21	0.41
1:B:588:TYR:CE2	1:B:592:LYS:HD3	2.56	0.41
1:C:148:ASP:HA	1:C:199:CYS:SG	2.60	0.41
1:C:223:GLU:O	1:C:227:VAL:HG23	2.20	0.41
1:B:205:VAL:HG11	1:B:259:LEU:HD22	2.02	0.41
1:A:265:PHE:HA	1:A:266:PRO:HD3	1.81	0.41
1:A:111:LYS:HD2	1:A:111:LYS:HA	1.92	0.40
1:B:365:LEU:HD23	1:B:448:LEU:HD23	2.04	0.40
1:A:195:VAL:HA	1:A:196:PRO:HD3	1.92	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	607/608 (100%)	589 (97%)	18 (3%)	0	100	100
1	В	607/608 (100%)	591 (97%)	16 (3%)	0	100	100
1	С	607/608 (100%)	589 (97%)	18 (3%)	0	100	100
All	All	1821/1824 (100%)	1769 (97%)	52 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	541/540 (100%)	541 (100%)	0	100	100
1	В	541/540 (100%)	540 (100%)	1 (0%)	93	98
1	С	541/540 (100%)	541 (100%)	0	100	100
All	All	1623/1620 (100%)	1622 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	154	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	589	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	cles
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	PGE	В	702	-	9,9,9	0.44	0	8,8,8	0.29	0	
3	N0Y	С	702	-	16,16,16	1.24	1 (6%)	20,20,20	0.76	1 (5%)	
3	N0Y	A	702	-	16,16,16	1.24	1 (6%)	20,20,20	0.78	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGE	В	702	-	-	4/7/7/7	-
3	N0Y	С	702	-	-	2/6/6/6	0/2/2/2



Continued from previous page...

Mo	l Type	Chain	Res	Link	Chirals	Torsions	Rings
3	N0Y	A	702	-	-	2/6/6/6	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
3	С	702	N0Y	O14-C11	3.75	1.45	1.37
3	A	702	N0Y	O14-C11	3.74	1.45	1.37

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	702	N0Y	C15-O14-C11	-2.88	111.27	117.51
3	С	702	N0Y	C15-O14-C11	-2.83	111.36	117.51

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	N0Y	C10-C11-O14-C15
3	С	702	N0Y	C10-C11-O14-C15
3	A	702	N0Y	C12-C11-O14-C15
3	С	702	N0Y	C12-C11-O14-C15
4	В	702	PGE	O2-C3-C4-O3
4	В	702	PGE	O1-C1-C2-O2
4	В	702	PGE	C1-C2-O2-C3
4	В	702	PGE	C3-C4-O3-C5

There are no ring outliers.


2 monomers are involved in 3 short contacts:


\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
4	В	702	PGE	2	0
3	С	702	N0Y	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring



in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	608/608 (100%)	-0.37	0 100 100	12, 20, 33, 59	0
1	В	608/608 (100%)	-0.11	2 (0%) 94 94	15, 28, 47, 75	3 (0%)
1	С	607/608 (99%)	-0.26	1 (0%) 95 95	16, 25, 39, 65	3 (0%)
All	All	1823/1824 (99%)	-0.24	3 (0%) 95 95	12, 24, 41, 75	6 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	610	ASP	3.5
1	В	547	TRP	2.1
1	С	441	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mo	l Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q<0.9
4	PGE	В	702	10/10	0.86	0.25	26,29,30,35	0







Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	N0Y	С	702	15/15	0.93	0.32	13,14,16,18	0
3	N0Y	A	702	15/15	0.96	0.21	13,14,16,18	0
2	ZN	В	701	1/1	0.97	0.06	27,27,27,27	0
2	ZN	A	701	1/1	0.97	0.08	17,17,17,17	0
2	ZN	С	701	1/1	0.98	0.06	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

