

# Full wwPDB X-ray Structure Validation Report (i)

Nov 16, 2023 – 07:31 AM JST

PDB ID : 6L86

Title : The structure of SfaA

Authors: Chen, T.Y.; Chen, J.; Zhou, J.; Chang, W.

Deposited on : 2019-11-05

Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	315	86%	6%	7%
1	В	315	85%	6%	9%
1	С	315	87%	5%	8%
1	D	315	85%	7%	8%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Taurine catabolism dioxygenase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	292	Total	С	N	О	S	0	0	0
1	A	292	2359	1522	390	441	6	0	U	
1	В	287	Total	С	N	О	S	0	0	0
1	Б	201	2327	1504	384	433	6	U	U	
1	С	290	Total	С	N	О	S	0	0	0
1		290	2347	1515	387	439	6		0	
1	D	280	Total	С	N	О	S	0	0	0
1		289	2336	1509	386	435	6		U	

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP A0A2H4T920
A	-18	GLY	-	expression tag	UNP A0A2H4T920
A	-17	SER	-	expression tag	UNP A0A2H4T920
A	-16	SER	-	expression tag	UNP A0A2H4T920
A	-15	HIS	-	expression tag	UNP A0A2H4T920
A	-14	HIS	_	expression tag	UNP A0A2H4T920
A	-13	HIS	-	expression tag	UNP A0A2H4T920
A	-12	HIS	-	expression tag	UNP A0A2H4T920
A	-11	HIS	-	expression tag	UNP A0A2H4T920
A	-10	HIS	-	expression tag	UNP A0A2H4T920
A	-9	SER	-	expression tag	UNP A0A2H4T920
A	-8	SER	_	expression tag	UNP A0A2H4T920
A	-7	GLY	-	expression tag	UNP A0A2H4T920
A	-6	LEU	-	expression tag	UNP A0A2H4T920
A	-5	VAL	-	expression tag	UNP A0A2H4T920
A	-4	PRO	-	expression tag	UNP A0A2H4T920
A	-3	ARG		expression tag	UNP A0A2H4T920
A	-2	GLY	-	expression tag	UNP A0A2H4T920
A	-1	SER	-	expression tag	UNP A0A2H4T920
A	0	HIS		expression tag	UNP A0A2H4T920
В	-19	MET	_	expression tag	UNP A0A2H4T920



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
В	-18	GLY	-	expression tag	UNP A0A2H4T920
В	-17	SER	_	expression tag	UNP A0A2H4T920
В	-16	SER	_	expression tag	UNP A0A2H4T920
В	-15	HIS	-	expression tag	UNP A0A2H4T920
В	-14	HIS	-	expression tag	UNP A0A2H4T920
В	-13	HIS	-	expression tag	UNP A0A2H4T920
В	-12	HIS	-	expression tag	UNP A0A2H4T920
В	-11	HIS	-	expression tag	UNP A0A2H4T920
В	-10	HIS	-	expression tag	UNP A0A2H4T920
В	-9	SER	-	expression tag	UNP A0A2H4T920
В	-8	SER	-	expression tag	UNP A0A2H4T920
В	-7	GLY	-	expression tag	UNP A0A2H4T920
В	-6	LEU	-	expression tag	UNP A0A2H4T920
В	-5	VAL	-	expression tag	UNP A0A2H4T920
В	-4	PRO	-	expression tag	UNP A0A2H4T920
В	-3	ARG	-	expression tag	UNP A0A2H4T920
В	-2	GLY	-	expression tag	UNP A0A2H4T920
В	-1	SER	-	expression tag	UNP A0A2H4T920
В	0	HIS	-	expression tag	UNP A0A2H4T920
С	-19	MET	-	expression tag	UNP A0A2H4T920
С	-18	GLY	-	expression tag	UNP A0A2H4T920
С	-17	SER	-	expression tag	UNP A0A2H4T920
С	-16	SER	-	expression tag	UNP A0A2H4T920
С	-15	HIS	-	expression tag	UNP A0A2H4T920
С	-14	HIS	-	expression tag	UNP A0A2H4T920
С	-13	HIS	-	expression tag	UNP A0A2H4T920
С	-12	HIS	-	expression tag	UNP A0A2H4T920
С	-11	HIS	-	expression tag	UNP A0A2H4T920
С	-10	HIS	-	expression tag	UNP A0A2H4T920
С	-9	SER	-	expression tag	UNP A0A2H4T920
С	-8	SER	-	expression tag	UNP A0A2H4T920
С	-7	GLY	-	expression tag	UNP A0A2H4T920
С	-6	LEU	-	expression tag	UNP A0A2H4T920
С	-5	VAL	-	expression tag	UNP A0A2H4T920
С	-4	PRO	-	expression tag	UNP A0A2H4T920
С	-3	ARG		expression tag	UNP A0A2H4T920
С	-2	GLY	_	expression tag	UNP A0A2H4T920
С	-1	SER	-	expression tag	UNP A0A2H4T920
С	0	HIS		expression tag	UNP A0A2H4T920
D	-19	MET	-	expression tag	UNP A0A2H4T920
D	-18	GLY	-	expression tag	UNP A0A2H4T920
D	-17	SER	-	expression tag	UNP A0A2H4T920



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP A0A2H4T920
D	-15	HIS	-	expression tag	UNP A0A2H4T920
D	-14	HIS	-	expression tag	UNP A0A2H4T920
D	-13	HIS	-	expression tag	UNP A0A2H4T920
D	-12	HIS	-	expression tag	UNP A0A2H4T920
D	-11	HIS	-	expression tag	UNP A0A2H4T920
D	-10	HIS	-	expression tag	UNP A0A2H4T920
D	-9	SER	-	expression tag	UNP A0A2H4T920
D	-8	SER	-	expression tag	UNP A0A2H4T920
D	-7	GLY	-	expression tag	UNP A0A2H4T920
D	-6	LEU	-	expression tag	UNP A0A2H4T920
D	-5	VAL	-	expression tag	UNP A0A2H4T920
D	-4	PRO	-	expression tag	UNP A0A2H4T920
D	-3	ARG	-	expression tag	UNP A0A2H4T920
D	-2	GLY	-	expression tag	UNP A0A2H4T920
D	-1	SER	-	expression tag	UNP A0A2H4T920
D	0	HIS	_	expression tag	UNP A0A2H4T920

• Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe 1 1	0	0
2	В	1	Total Fe 1 1	0	0
2	С	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0

• Molecule 3 is (2S)-2-hydroxybutanedioic acid (three-letter code: LMR) (formula:  $C_4H_6O_5$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Δ	1	Total C O	0	0
	Λ	1	9 4 5	U	0
3	В	1	Total C O	0	0
3	Ъ	1	9 4 5	U	
3	В	1	Total C O	0	0
3	Ъ	1	9 4 5	U	
3	С	1	Total C O	0	0
3		1	9 4 5	U	
2	3 D	D 1	Total C O	0	0
3			9 4 5	0	

• Molecule 4 is D-MALATE (three-letter code: MLT) (formula:  $C_4H_6O_5$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	A	1	Total 9	C 4	O 5	0	0

#### • Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	166	Total O 166 166	0	0
5	В	146	Total O 146 146	0	0
5	С	131	Total O 131 131	0	0
5	D	140	Total O 140 140	0	0




#### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Taurine catabolism dioxygenase




• Molecule 1: Taurine catabolism dioxygenase



• Molecule 1: Taurine catabolism dioxygenase



• Molecule 1: Taurine catabolism dioxygenase









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.17Å 150.17Å 387.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.10 - 2.23 $49.83 - 2.23$	Depositor EDS
% Data completeness	99.1 (43.10-2.23)	Depositor
(in resolution range)	99.1 (49.83-2.23)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.05 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
D D	0.190 , 0.228	Depositor
$R, R_{free}$	0.190 , $0.228$	DCC
$R_{free}$ test set	3999 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.35, 41.4	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	$\begin{array}{c} 0.006 \text{ for } -1/3*\text{h} + 1/3*\text{k} + 1/3*\text{l}, -\text{k}, 8/3*\text{h} + 4/\\ 3*\text{k} + 1/3*\text{l} \\ 0.014 \text{ for } -2/3*\text{h} - 1/3*\text{k} - 1/3*\text{l}, -1/3*\text{h} - 2/3*\text{k} + \\ 1/3*\text{l}, -4/3*\text{h} + 4/3*\text{k} + 1/3*\text{l} \\ 0.009 \text{ for } -\text{h}, 1/3*\text{h} - 1/3*\text{k} - 1/3*\text{l}, -4/3*\text{h} - 8/3*\text{k} \\ +1/3*\text{l} \end{array}$	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10010	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LMR, MLT, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	$\mathbf{angles}$
IVIOI	Chain	RMSZ $ $ $# Z  >$		RMSZ	# Z  > 5
1	A	0.25	0/2427	0.45	0/3296
1	В	0.25	0/2393	0.45	0/3248
1	С	0.24	0/2414	0.45	0/3279
1	D	0.25	0/2402	0.46	0/3261
All	All	0.25	0/9636	0.45	0/13084

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2359	0	2309	13	0
1	В	2327	0	2280	9	0
1	С	2347	0	2294	10	0
1	D	2336	0	2285	11	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	9	0	4	2	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
3	В	18	0	8	1	0
3	С	9	0	4	0	0
3	D	9	0	4	0	0
4	A	9	0	4	0	0
5	A	166	0	0	0	0
5	В	146	0	0	0	0
5	С	131	0	0	0	0
5	D	140	0	0	0	0
All	All	10010	0	9192	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

distance           HB3         1.72           NE2         2.22           HG12         1.80           D11         1.82           HB3         1.84           CD2         2.22           HD23         1.87           DE1         2.38	0.72 0.72 0.63 0.61 0.61 0.59 0.58 0.56
NE2 2.22 HG12 1.80 D11 1.82 HB3 1.82 H3A 1.84 CD2 2.22 HD23 1.87 OE1 2.38	0.72 0.63 0.61 0.61 0.59 0.58 0.56
HG12 1.80 D11 1.82 HB3 1.82 H3A 1.84 CD2 2.22 HD23 1.87 OE1 2.38	0.63 0.61 0.61 0.59 0.58 0.56
D11     1.82       HB3     1.82       H3A     1.84       CD2     2.22       HD23     1.87       OE1     2.38	0.61 0.61 0.59 0.58 0.56
HB3 1.82 H3A 1.84 CD2 2.22 HD23 1.87 OE1 2.38	0.61 0.59 0.58 0.56
H3A 1.84 CD2 2.22 HD23 1.87 OE1 2.38	0.59 0.58 0.56
CD2 2.22 HD23 1.87 OE1 2.38	0.58 0.56
HD23 1.87 DE1 2.38	0.56
OE1 2.38	
	0.54
TO10	0.04
$4G12 \mid 1.90$	0.53
:HB 2.38	0.53
N:N 2.28	0.53
HG22 1.92	0.51
CG 2.45	0.51
CD2 $2.46$	0.51
H3A 1.93	0.50
:CD 2.16	0.48
CD2 2.50	0.46
HB3 1.95	0.46
:NZ 2.46	0.46
HB3 1.98	0.46
2.28	0.45
HD2 1.69	0.45
HB3 1.98	0.45
:HA 1.98	0.45
	HG12 1.90 :HB 2.38 N:N 2.28 HG22 1.92 CG 2.45 CD2 2.46 :H3A 1.93 :CD 2.16 :CD2 2.50 :HB3 1.95 :NZ 2.46 :HB3 1.98 :C1 2.28 HD2 1.69 HB3 1.98



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
7100111 1	1100111 2	$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
1:D:198:THR:OG1	1:D:200:GLU:HG3	2.17	0.44
1:B:172:PRO:HG2	1:B:175:GLU:HG3	1.99	0.44
1:D:74:PRO:HD3	1:D:92:ILE:CD1	2.48	0.44
1:B:123:PRO:HD2	1:B:127:ARG:HG2	1.99	0.43
1:D:107:PHE:CD2	1:D:158:VAL:HB	2.53	0.43
1:B:145:LEU:HD12	1:B:145:LEU:HA	1.81	0.42
1:D:74:PRO:HD3	1:D:92:ILE:HD11	2.00	0.42
1:C:107:PHE:CD2	1:C:158:VAL:HB	2.55	0.42
1:D:106:GLN:HG3	1:D:204:TYR:CE2	2.55	0.42
1:A:36:TYR:CE2	1:A:197:LEU:HD22	2.55	0.41
1:C:75:GLU:HG3	1:C:76:PHE:CD2	2.55	0.41
1:A:100:PHE:O	1:A:102:HIS:ND1	2.48	0.41
1:B:28:ILE:HG22	1:B:32:LYS:HD2	2.03	0.41
1:A:185:PRO:HG3	1:D:249:HIS:CE1	2.56	0.40
1:D:215:ASP:HB2	1:D:219:LYS:H	1.86	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	288/315 (91%)	281 (98%)	7 (2%)	0	100	100
1	В	281/315~(89%)	273 (97%)	7 (2%)	1 (0%)	34	35
1	С	284/315 (90%)	278 (98%)	5 (2%)	1 (0%)	34	35
1	D	281/315 (89%)	277 (99%)	4 (1%)	0	100	100
All	All	1134/1260 (90%)	1109 (98%)	23 (2%)	2 (0%)	47	53

All (2) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	С	273	PRO
1	В	285	ASP

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outlier		Percentiles		
1	A	258/277~(93%)	255 (99%)	3 (1%)	71 78		
1	В	254/277~(92%)	250 (98%)	4 (2%)	62 70		
1	$\mathbf{C}$	257/277~(93%)	255 (99%)	2 (1%)	81 87		
1	D	255/277~(92%)	253 (99%)	2 (1%)	81 87		
All	All	$1024/1108 \; (92\%)$	1013 (99%)	11 (1%)	73 80		

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	TYR
1	A	197	LEU
1	A	266	ARG
1	В	2	GLU
1	В	105	TYR
1	В	124	GLN
1	В	197	LEU
1	С	105	TYR
1	С	286	GLU
1	D	105	TYR
1	D	197	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	167	HIS



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain H		Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	les		
IVIOI	Wioi Type	Chain	Chain	Chain	rtes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	LMR	В	303	2	8,8,8	1.04	0	10,10,10	1.35	2 (20%)		
4	MLT	A	303	-	8,8,8	1.04	0	10,10,10	1.34	1 (10%)		
3	LMR	С	302	-	8,8,8	1.02	0	10,10,10	1.29	2 (20%)		
3	LMR	D	302	2	8,8,8	1.03	0	10,10,10	1.37	2 (20%)		
3	LMR	A	302	2	8,8,8	1.03	0	10,10,10	1.21	1 (10%)		
3	LMR	В	302	-	8,8,8	0.99	0	10,10,10	1.21	1 (10%)		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LMR	В	303	2	-	0/8/8/8	-
4	MLT	A	303	-	-	2/8/8/8	-
3	LMR	С	302	-	-	0/8/8/8	-



 $Continued\ from\ previous\ page...$ 

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LMR	D	302	2	-	5/8/8/8	-
3	LMR	A	302	2	-	7/8/8/8	-
3	LMR	В	302	-	-	6/8/8/8	-

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	D	302	LMR	O1B-C1-C2	2.85	118.98	112.72
3	С	302	LMR	O1B-C1-C2	2.76	118.79	112.72
4	A	303	MLT	O2-C1-C2	2.76	118.79	112.72
3	A	302	LMR	O1B-C1-C2	2.50	118.20	112.72
3	В	302	LMR	O1B-C1-C2	2.33	117.84	112.72
3	В	303	LMR	O1B-C1-C2	2.33	117.84	112.72
3	D	302	LMR	O1A-C1-C2	-2.17	118.29	122.54
3	С	302	LMR	O1A-C1-C2	-2.17	118.30	122.54
3	В	303	LMR	O1A-C1-C2	-2.06	118.52	122.54

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	LMR	O1A-C1-C2-O2
3	A	302	LMR	O1A-C1-C2-C3
3	A	302	LMR	O1B-C1-C2-O2
3	В	302	LMR	O1A-C1-C2-O2
3	В	302	LMR	O1B-C1-C2-O2
3	A	302	LMR	O1B-C1-C2-C3
3	В	302	LMR	O1B-C1-C2-C3
3	A	302	LMR	C2-C3-C4-O4A
3	В	302	LMR	O1A-C1-C2-C3
3	A	302	LMR	C2-C3-C4-O4B
3	В	302	LMR	C2-C3-C4-O4A
3	D	302	LMR	C2-C3-C4-O4A
3	D	302	LMR	C2-C3-C4-O4B
4	A	303	MLT	C2-C3-C4-O4
4	A	303	MLT	C2-C3-C4-O5
3	A	302	LMR	O2-C2-C3-C4
3	D	302	LMR	O2-C2-C3-C4
3	В	302	LMR	C2-C3-C4-O4B



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	302	LMR	O1A-C1-C2-C3
3	D	302	LMR	O1B-C1-C2-C3


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	303	LMR	1	0
3	A	302	LMR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	292/315~(92%)	-0.43	4 (1%) 75 76	18, 28, 52, 85	0
1	В	287/315 (91%)	-0.42	0 100 100	18, 31, 57, 76	0
1	С	290/315 (92%)	-0.46	1 (0%) 94 94	20, 32, 55, 96	0
1	D	289/315 (91%)	-0.46	2 (0%) 87 87	20, 32, 55, 81	0
All	All	1158/1260 (91%)	-0.44	7 (0%) 89 89	18, 31, 56, 96	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	GLY	4.5
1	A	7	SER	4.1
1	A	6	GLN	3.8
1	D	86	PRO	3.1
1	A	9	SER	2.4
1	С	295	ALA	2.1
1	D	222	GLU	2.0

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	FE2	D	301	1/1	0.69	0.18	51,51,51,51	1
3	LMR	A	302	9/9	0.84	0.21	32,39,51,59	0
3	LMR	D	302	9/9	0.87	0.22	33,46,57,58	0
2	FE2	С	301	1/1	0.91	0.12	63,63,63,63	1
3	LMR	В	303	9/9	0.93	0.14	28,35,43,45	0
3	LMR	С	302	9/9	0.93	0.17	37,43,53,57	0
2	FE2	В	301	1/1	0.93	0.07	45,45,45,45	0
3	LMR	В	302	9/9	0.94	0.13	22,28,38,44	0
2	FE2	A	301	1/1	0.95	0.10	46,46,46,46	0
4	MLT	A	303	9/9	0.95	0.13	20,25,32,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













# Electron density around LMR D 302: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)














# Electron density around LMR C 302: $2 {\rm mF}_o\text{-DF}_c \ ({\rm at}\ 0.7\ {\rm rmsd})\ {\rm in\ gray}$ ${\rm mF}_o\text{-DF}_c \ ({\rm at}\ 3\ {\rm rmsd})\ {\rm in\ purple}\ ({\rm negative})$


and green (positive)
















#### Electron density around LMR B 302:


 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
















# 6.5 Other polymers (i)

There are no such residues in this entry.

