



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 02:23 AM BST

PDB ID : 1LEG  
Title : Crystal Structure of H-2Kb bound to the dEV8 peptide  
Authors : Luz, J.G.; Huang, M.; Garcia, K.C.; Rudolph, M.G.; Apostolopoulos, V.;  
Teyton, L.; Wilson, I.A.  
Deposited on : 2002-04-09  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
2	B	99	
3	P	8	
4	C	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	C	2	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN, K-B ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2233	1408	393	423	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	CSO	CYS	modified residue	UNP P01901


- Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	821	524	138	152	7	0	0	0

- Molecule 3 is a protein called NADH-ubiquinone oxidoreductase MLRQ subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	P	8	76	51	10	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	C	3	38	22	2	14	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
6	A	1	5	4	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	281	Total	O	0	0
			281	281		
7	B	92	Total	O	0	0
			92	92		
7	P	7	Total	O	0	0
			7	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN, K-B ALPHA CHAIN



- Molecule 2: BETA-2-MICROGLOBULIN



- Molecule 3: NADH-ubiquinone oxidoreductase MLRQ subunit



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.69Å 87.69Å 45.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.75 17.82 – 1.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.75) 86.4 (17.82-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 1.65Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.217 , 0.241 0.210 , 0.229	Depositor DCC
$R_{free}$ test set	2294 reflections (4.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 63.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3572	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, PO4, CSO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/2286	0.70	2/3102 (0.1%)
2	B	0.49	0/847	0.76	0/1148
3	P	3.50	2/78 (2.6%)	1.30	1/102 (1.0%)
All	All	0.73	2/3211 (0.1%)	0.73	3/4352 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	8	VAL	C-OXT	26.37	1.73	1.23
3	P	8	VAL	C-O	-14.69	0.95	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	8	VAL	CA-C-O	9.40	139.84	120.10
1	A	272	LEU	CA-CB-CG	6.70	130.71	115.30
1	A	28	VAL	N-CA-C	-5.58	95.93	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2233	0	2119	69	0
2	B	821	0	796	35	0
3	P	76	0	70	12	0
4	C	38	0	34	5	0
5	A	14	0	13	0	0
6	A	5	0	0	0	0
6	B	5	0	0	1	0
7	A	281	0	0	6	0
7	B	92	0	0	3	0
7	P	7	0	0	0	0
All	All	3572	0	3032	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:8:VAL:C	3:P:8:VAL:OXT	1.73	1.24
3:P:2:GLN:HE21	3:P:3:TYR:H	1.08	0.98
1:A:195:PRO:O	1:A:196:GLU:HG3	1.73	0.88
1:A:173:LYS:NZ	4:C:3:FUC:H63	1.88	0.88
2:B:73:THR:HB	2:B:76:ASP:OD2	1.74	0.88
1:A:50:ARG:HH11	1:A:50:ARG:HG3	1.41	0.85
7:A:1092:HOH:O	4:C:3:FUC:H4	1.76	0.85
3:P:8:VAL:OXT	3:P:8:VAL:CA	2.25	0.85
2:B:48:LYS:HD3	2:B:68:THR:OG1	1.76	0.84
1:A:223:GLU:HB3	1:A:225:ILE:HD13	1.59	0.84
1:A:263:HIS:CD2	1:A:265:GLY:H	1.95	0.84
1:A:263:HIS:HD2	1:A:265:GLY:H	1.25	0.83
1:A:128:GLU:HG3	7:A:1116:HOH:O	1.81	0.80
1:A:50:ARG:HH11	1:A:50:ARG:CG	1.95	0.79
2:B:2:GLN:HE21	2:B:86:SER:HA	1.47	0.79
1:A:62:ARG:NH1	3:P:1:GLU:OE2	2.20	0.75
1:A:258:THR:HG22	1:A:273:ARG:HG3	1.69	0.74
1:A:50:ARG:NH1	1:A:50:ARG:HG3	2.02	0.71
1:A:225:ILE:O	1:A:226:GLN:HB2	1.91	0.71
2:B:38:GLN:HB2	2:B:45:LYS:HD3	1.71	0.71
1:A:6:ARG:HH11	1:A:6:ARG:HG3	1.55	0.70
1:A:173:LYS:HZ1	4:C:3:FUC:H63	1.54	0.70
1:A:173:LYS:HZ2	4:C:3:FUC:H63	1.56	0.68
1:A:6:ARG:NH1	1:A:6:ARG:HG3	2.10	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:38:GLN:HB2	2:B:45:LYS:CD	2.24	0.66
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.32	0.64
1:A:12:VAL:HG22	1:A:94:THR:HG23	1.81	0.63
1:A:266:LEU:HD13	1:A:270:LEU:HG	1.80	0.62
1:A:57:PRO:HD2	1:A:58:GLU:OE2	1.99	0.62
1:A:216:THR:HB	1:A:225:ILE:HD11	1.83	0.61
1:A:23:MET:HE1	2:B:54:MET:HG3	1.83	0.60
2:B:73:THR:HG22	2:B:75:THR:H	1.66	0.60
1:A:219:LEU:HB3	1:A:224:LEU:HD21	1.83	0.59
1:A:9:VAL:HG22	1:A:24:GLU:HG2	1.84	0.58
3:P:2:GLN:HE21	3:P:3:TYR:N	1.92	0.58
1:A:58:GLU:CD	1:A:58:GLU:H	2.08	0.56
1:A:101:CYS:HB2	7:A:1272:HOH:O	2.04	0.56
2:B:2:GLN:NE2	2:B:86:SER:HA	2.20	0.56
1:A:194:ARG:HD3	1:A:198:LYS:HB3	1.86	0.56
2:B:48:LYS:CE	2:B:69:GLU:H	2.18	0.56
3:P:8:VAL:OXT	3:P:8:VAL:HA	2.05	0.55
2:B:19:LYS:HD3	7:B:1074:HOH:O	2.06	0.55
2:B:83:LYS:HE3	2:B:90:PRO:CG	2.37	0.55
1:A:44:ARG:HH22	1:A:61:GLU:HA	1.72	0.55
1:A:23:MET:CE	2:B:54:MET:HG3	2.37	0.55
2:B:38:GLN:HB2	2:B:45:LYS:CE	2.38	0.54
2:B:91:LYS:HE2	7:B:1078:HOH:O	2.07	0.54
2:B:48:LYS:HE3	2:B:69:GLU:H	1.72	0.54
3:P:2:GLN:NE2	3:P:3:TYR:H	1.91	0.54
1:A:222:GLU:HG2	1:A:222:GLU:O	2.08	0.54
2:B:39:MET:H	2:B:45:LYS:NZ	2.06	0.54
1:A:258:THR:HG23	7:A:1215:HOH:O	2.09	0.52
2:B:83:LYS:HG3	2:B:90:PRO:HB3	1.91	0.52
1:A:50:ARG:CG	1:A:50:ARG:NH1	2.63	0.51
1:A:177:ALA:HA	4:C:1:NAG:H82	1.92	0.51
1:A:14:ARG:NH1	1:A:21:ARG:HB2	2.25	0.51
1:A:217:TRP:HD1	1:A:228:MET:HE2	1.76	0.50
1:A:66:LYS:HG3	3:P:4:LYS:HG2	1.91	0.50
1:A:255:GLN:HE21	1:A:273:ARG:NH1	2.10	0.50
2:B:1:ILE:CD1	2:B:1:ILE:N	2.75	0.49
2:B:1:ILE:HD12	2:B:1:ILE:N	2.26	0.49
1:A:42:ASN:O	1:A:42:ASN:CG	2.50	0.49
1:A:6:ARG:HH11	1:A:6:ARG:CG	2.25	0.49
1:A:217:TRP:HD1	1:A:228:MET:CE	2.25	0.49
1:A:191:HIS:HB2	1:A:274:TRP:NE1	2.28	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:36:GLU:CD	2:B:83:LYS:HD2	2.33	0.48
1:A:210:PRO:O	1:A:263:HIS:HE1	1.97	0.48
1:A:190:THR:OG1	1:A:202:ARG:HB3	2.14	0.48
1:A:263:HIS:HD2	1:A:265:GLY:N	2.02	0.48
1:A:133:TRP:CH2	1:A:156:LEU:HD23	2.50	0.47
1:A:117:ALA:HB2	2:B:60:TRP:CD2	2.49	0.47
1:A:258:THR:HG22	1:A:273:ARG:CG	2.42	0.47
1:A:225:ILE:HD12	1:A:225:ILE:H	1.79	0.47
1:A:28:VAL:HG11	1:A:179:LEU:HD13	1.95	0.46
1:A:225:ILE:CD1	1:A:225:ILE:H	2.29	0.46
2:B:7:ILE:HD12	2:B:91:LYS:HD3	1.96	0.46
1:A:256:TYR:N	1:A:256:TYR:CD1	2.81	0.46
1:A:263:HIS:CD2	1:A:265:GLY:N	2.76	0.46
1:A:167:TRP:CE2	3:P:1:GLU:HG3	2.52	0.45
1:A:195:PRO:C	1:A:196:GLU:HG3	2.33	0.45
2:B:38:GLN:OE1	2:B:45:LYS:HD3	2.17	0.44
1:A:167:TRP:CD2	3:P:1:GLU:HG3	2.52	0.44
2:B:48:LYS:HE3	2:B:69:GLU:HB2	1.99	0.44
2:B:38:GLN:HB2	2:B:45:LYS:HE2	2.00	0.44
2:B:34:HIS:HA	7:B:1020:HOH:O	2.16	0.44
1:A:144:LYS:NZ	1:A:148:GLU:OE2	2.37	0.43
1:A:225:ILE:HD12	1:A:225:ILE:N	2.33	0.43
2:B:83:LYS:HE3	2:B:90:PRO:HG3	2.00	0.43
2:B:1:ILE:HD12	2:B:1:ILE:H1	1.82	0.43
1:A:115:GLN:HG2	7:A:1100:HOH:O	2.19	0.43
1:A:228:MET:HB2	1:A:246:SER:O	2.19	0.43
1:A:128:GLU:HG3	1:A:128:GLU:H	1.55	0.42
1:A:133:TRP:O	1:A:144:LYS:HE2	2.19	0.42
2:B:83:LYS:HG3	2:B:90:PRO:HG3	2.01	0.42
2:B:13:HIS:O	2:B:14:PRO:C	2.58	0.42
3:P:4:LYS:HE3	3:P:4:LYS:HB3	1.71	0.42
2:B:39:MET:H	2:B:45:LYS:HZ3	1.67	0.42
2:B:36:GLU:OE2	2:B:83:LYS:HD2	2.20	0.42
2:B:83:LYS:CG	2:B:90:PRO:HG3	2.48	0.42
2:B:33:PRO:O	6:B:990:PO4:O3	2.37	0.42
1:A:255:GLN:HB2	7:A:1175:HOH:O	2.19	0.41
1:A:233:THR:OG1	1:A:243:LYS:HD2	2.20	0.41
3:P:2:GLN:HG3	3:P:3:TYR:N	2.35	0.41
1:A:230:LEU:C	1:A:230:LEU:CD1	2.89	0.41
1:A:108:ARG:HH11	1:A:108:ARG:CG	2.34	0.41
1:A:217:TRP:HB2	1:A:228:MET:CE	2.51	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.89	0.40
1:A:37:ASP:OD1	1:A:39:ASP:HB2	2.21	0.40
1:A:41:GLU:O	1:A:42:ASN:HB3	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/274 (99%)	264 (97%)	5 (2%)	2 (1%)	22	8
2	B	97/99 (98%)	96 (99%)	0	1 (1%)	15	4
3	P	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	374/381 (98%)	365 (98%)	6 (2%)	3 (1%)	19	6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	ILE
1	A	226	GLN
2	B	49	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	206 (89%)	25 (11%)	6	1
2	B	94/94 (100%)	90 (96%)	4 (4%)	29	9
3	P	8/8 (100%)	7 (88%)	1 (12%)	4	0
All	All	333/333 (100%)	303 (91%)	30 (9%)	9	1

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	39	ASP
1	A	41	GLU
1	A	42	ASN
1	A	45	TYR
1	A	50	ARG
1	A	58	GLU
1	A	88	SER
1	A	108	ARG
1	A	128	GLU
1	A	134	THR
1	A	149	GLN
1	A	154	GLU
1	A	194	ARG
1	A	196	GLU
1	A	197	ASP
1	A	220	ASN
1	A	222	GLU
1	A	227	ASP
1	A	228	MET
1	A	229	GLU
1	A	230	LEU
1	A	255	GLN
1	A	268	GLU
1	A	273	ARG
2	B	1	ILE
2	B	56	PHE
2	B	58	LYS
2	B	73	THR
3	P	8	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	114	GLN
1	A	255	GLN
1	A	263	HIS
2	B	2	GLN
3	P	2	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSO	A	121	1	3,6,7	0.75	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	121	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates i

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	C	1	1,4	14,14,15	0.67	0	17,19,21	0.68	1 (5%)
4	NAG	C	2	4	14,14,15	0.72	1 (7%)	17,19,21	0.82	0
4	FUC	C	3	4	10,10,11	0.50	0	14,14,16	1.09	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1
4	FUC	C	3	4	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2	NAG	C1-C2	2.11	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	3	FUC	C1-C2-C3	3.09	113.47	109.67
4	C	1	NAG	C2-N2-C7	-2.04	119.99	122.90

There are no chirality outliers.

All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	C	1	NAG	C4-C5-C6-O6
4	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	3	FUC	4	0
4	C	1	NAG	1	0

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	PO4	B	990	-	4,4,4	0.88	0	6,6,6	0.73	0
5	NAG	A	900	1	14,14,15	0.56	0	17,19,21	0.67	0
6	PO4	A	991	-	4,4,4	0.81	0	6,6,6	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	900	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	990	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	273/274 (99%)	0.14	13 (4%) 30 36	11, 18, 46, 65	0
2	B	99/99 (100%)	0.29	4 (4%) 38 45	11, 21, 39, 59	0
3	P	8/8 (100%)	0.60	1 (12%) 3 5	18, 26, 33, 35	0
All	All	380/381 (99%)	0.19	18 (4%) 31 37	11, 19, 44, 65	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	10.5
1	A	225	ILE	7.3
1	A	226	GLN	6.3
1	A	274	TRP	5.8
1	A	41	GLU	5.6
1	A	220	ASN	5.2
1	A	42	ASN	4.6
1	A	256	TYR	3.8
2	B	48	LYS	3.6
1	A	196	GLU	3.5
1	A	255	GLN	3.4
1	A	89	LYS	3.1
1	A	227	ASP	3.0
1	A	222	GLU	2.5
1	A	40	ALA	2.5
2	B	49	VAL	2.2
3	P	8	VAL	2.1
2	B	73	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	121	7/8	0.86	0.10	17,17,22,25	0

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	C	2	14/15	0.71	0.40	59,63,69,72	0
4	FUC	C	3	10/11	0.77	0.24	25,52,63,70	0
4	NAG	C	1	14/15	0.88	0.16	25,35,52,60	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	900	14/15	0.60	0.35	51,61,65,69	0
6	PO4	A	991	5/5	0.67	0.33	69,72,79,100	0
6	PO4	B	990	5/5	0.91	0.16	44,45,54,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.