



Full wwPDB X-ray Structure Validation Report

Feb 10, 2024 – 02:37 PM EST

PDB ID : 2LH2
Title : X-RAY STRUCTURAL INVESTIGATION OF LEGHEMOGLOBIN. VI. STRUCTURE OF ACETATE-FERRILEGHEMOGLOBIN AT A RESOLUTION OF 2.0 ANGSTROMS (RUSSIAN)
Authors : Vainshtein, B.K.; Harutyunyan, E.H.; Kuranova, I.P.; Borisov, V.V.; Sosfenov, N.I.; Pavlovsky, A.G.; Grebenko, A.I.; Konareva, N.V.
Deposited on : 1982-04-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	153	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LEGHEMOGLOBIN (AQUO MET).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	153	1180	761	193	225	1	36	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	GLU	GLN	conflict	UNP P02240
A	150	ASP	ASN	conflict	UNP P02240

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	3	0

- Molecule 3 is OXYGEN ATOM (three-letter code: O) (formula: O).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	67	Total O 67 67	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LEGHEMOGLOBIN (AQUO MET)

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	B 1 1 2	Depositor
Cell constants a, b, c, α , β , γ	93.22Å 38.26Å 52.01Å 90.00° 90.00° 98.80°	Depositor
Resolution (Å)	(Not available) – 2.00 9.93 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 96.4 (9.93-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available) 0.500 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtrriage
Anisotropy	0.156	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	1.23 , 279.2	EDS
L-test for twinning ¹	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.51	EDS
Total number of atoms	1291	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	3.25	138/1214 (11.4%)	2.13	39/1648 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8

All (138) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	95	SER	CB-OG	10.60	1.56	1.42
1	A	138	TYR	CB-CG	9.88	1.66	1.51
1	A	15	TRP	CD2-CE2	9.48	1.52	1.41
1	A	138	TYR	CZ-OH	9.47	1.53	1.37
1	A	112	GLU	CG-CD	9.30	1.65	1.51
1	A	132[A]	SER	CA-CB	9.09	1.66	1.52
1	A	132[B]	SER	CA-CB	9.09	1.66	1.52
1	A	132[C]	SER	CA-CB	9.09	1.66	1.52
1	A	120	GLU	CG-CD	8.72	1.65	1.51
1	A	106	HIS	CB-CG	8.58	1.65	1.50
1	A	72	TYR	CE1-CZ	8.49	1.49	1.38
1	A	123	GLY	CA-C	8.48	1.65	1.51
1	A	15	TRP	CB-CG	8.37	1.65	1.50
1	A	5	GLU	CD-OE2	8.26	1.34	1.25
1	A	94	GLY	CA-C	8.13	1.64	1.51
1	A	18	PHE	CB-CG	8.06	1.65	1.51
1	A	65	GLY	C-O	8.02	1.36	1.23
1	A	13	SER	CA-CB	7.94	1.64	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	35	GLU	CD-OE2	7.69	1.34	1.25
1	A	72	TYR	CG-CD2	7.61	1.49	1.39
1	A	44	PHE	CB-CG	7.49	1.64	1.51
1	A	128	GLU	CD-OE2	-7.40	1.17	1.25
1	A	46	PHE	CB-CG	7.30	1.63	1.51
1	A	138	TYR	CD2-CE2	7.18	1.50	1.39
1	A	140	GLU	CB-CG	7.15	1.65	1.52
1	A	129	GLU	CD-OE1	7.07	1.33	1.25
1	A	15	TRP	CZ3-CH2	7.06	1.51	1.40
1	A	144	VAL	CB-CG2	7.06	1.67	1.52
1	A	110	VAL	CB-CG2	7.05	1.67	1.52
1	A	11	VAL	CB-CG1	7.03	1.67	1.52
1	A	121	VAL	CB-CG2	7.03	1.67	1.52
1	A	35	GLU	CD-OE1	-6.99	1.18	1.25
1	A	134	TRP	N-CA	6.96	1.60	1.46
1	A	72	TYR	C-O	6.94	1.36	1.23
1	A	101	GLY	CA-C	6.90	1.62	1.51
1	A	111	LYS	N-CA	6.88	1.60	1.46
1	A	85	VAL	CB-CG2	6.87	1.67	1.52
1	A	113	ALA	CA-CB	6.85	1.66	1.52
1	A	102	VAL	CB-CG1	6.84	1.67	1.52
1	A	88	ALA	N-CA	6.83	1.60	1.46
1	A	104	ASP	N-CA	6.79	1.59	1.46
1	A	13	SER	CB-OG	-6.76	1.33	1.42
1	A	40	ALA	CA-CB	6.75	1.66	1.52
1	A	68	PHE	C-O	6.74	1.36	1.23
1	A	27	HIS	CE1-NE2	6.71	1.48	1.32
1	A	152	ALA	C-O	6.64	1.35	1.23
1	A	79	GLU	CB-CG	6.62	1.64	1.52
1	A	138	TYR	CD1-CE1	6.59	1.49	1.39
1	A	124	ALA	N-CA	6.57	1.59	1.46
1	A	109	VAL	CA-CB	6.48	1.68	1.54
1	A	124	ALA	C-O	6.44	1.35	1.23
1	A	95	SER	N-CA	6.41	1.59	1.46
1	A	6	SER	CA-CB	6.40	1.62	1.52
1	A	11	VAL	N-CA	6.38	1.59	1.46
1	A	111	LYS	CD-CE	6.37	1.67	1.51
1	A	14	SER	CB-OG	-6.36	1.33	1.42
1	A	61	GLN	C-O	6.36	1.35	1.23
1	A	86	THR	N-CA	6.33	1.59	1.46
1	A	145	ILE	N-CA	6.33	1.58	1.46
1	A	15	TRP	NE1-CE2	-6.31	1.29	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	45	SER	CB-OG	6.31	1.50	1.42
1	A	67	VAL	CB-CG2	6.31	1.66	1.52
1	A	28	ARG	CZ-NH1	6.29	1.41	1.33
1	A	99	SER	C-O	6.28	1.35	1.23
1	A	149	MET	C-O	6.26	1.35	1.23
1	A	68	PHE	CG-CD2	6.22	1.48	1.38
1	A	73	GLU	CG-CD	6.20	1.61	1.51
1	A	15	TRP	C-O	6.18	1.35	1.23
1	A	15	TRP	CD1-NE1	6.18	1.48	1.38
1	A	16	GLU	CG-CD	6.17	1.61	1.51
1	A	148	GLU	CB-CG	6.16	1.63	1.52
1	A	148	GLU	CD-OE2	6.09	1.32	1.25
1	A	79	GLU	C-O	6.07	1.34	1.23
1	A	97	HIS	CA-CB	6.07	1.67	1.53
1	A	102	VAL	N-CA	6.05	1.58	1.46
1	A	122	VAL	N-CA	6.05	1.58	1.46
1	A	68	PHE	CE1-CZ	6.03	1.48	1.37
1	A	55	GLN	C-O	6.03	1.34	1.23
1	A	2	ALA	CA-CB	6.01	1.65	1.52
1	A	106	HIS	ND1-CE1	5.99	1.49	1.34
1	A	9	ALA	CA-CB	5.96	1.65	1.52
1	A	97	HIS	CG-CD2	-5.90	1.25	1.35
1	A	63	HIS	CE1-NE2	5.90	1.46	1.32
1	A	98	VAL	CB-CG2	5.88	1.65	1.52
1	A	99	SER	N-CA	5.86	1.58	1.46
1	A	27	HIS	CG-ND1	5.83	1.51	1.38
1	A	76	ILE	C-O	5.82	1.34	1.23
1	A	75	ALA	N-CA	5.80	1.57	1.46
1	A	79	GLU	CD-OE2	5.80	1.32	1.25
1	A	53	VAL	CA-CB	5.79	1.66	1.54
1	A	75	ALA	C-O	5.78	1.34	1.23
1	A	90	LEU	CA-CB	5.77	1.67	1.53
1	A	77	GLN	CG-CD	5.76	1.64	1.51
1	A	8	ALA	C-O	5.72	1.34	1.23
1	A	18	PHE	N-CA	5.71	1.57	1.46
1	A	152	ALA	N-CA	5.69	1.57	1.46
1	A	111	LYS	CB-CG	5.66	1.67	1.52
1	A	58	PRO	N-CD	5.66	1.55	1.47
1	A	59	GLU	CD-OE2	5.64	1.31	1.25
1	A	4	THR	C-O	5.64	1.34	1.23
1	A	52	GLU	CG-CD	-5.62	1.43	1.51
1	A	84	VAL	CB-CG1	5.61	1.64	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	120	GLU	CA-CB	5.60	1.66	1.53
1	A	96	VAL	CB-CG1	5.60	1.64	1.52
1	A	71	VAL	CB-CG1	5.51	1.64	1.52
1	A	125	LYS	CD-CE	5.50	1.65	1.51
1	A	68	PHE	N-CA	5.48	1.57	1.46
1	A	69	LYS	C-O	5.48	1.33	1.23
1	A	92	ASN	CB-CG	5.43	1.63	1.51
1	A	140	GLU	CD-OE2	5.42	1.31	1.25
1	A	153	ALA	C-OXT	5.41	1.33	1.23
1	A	92	ASN	C-O	5.39	1.33	1.23
1	A	30	PHE	CG-CD2	-5.38	1.30	1.38
1	A	127	SER	C-O	5.38	1.33	1.23
1	A	141	LEU	N-CA	5.37	1.57	1.46
1	A	82	GLY	CA-C	5.35	1.60	1.51
1	A	45	SER	N-CA	5.34	1.57	1.46
1	A	130	LEU	N-CA	5.33	1.57	1.46
1	A	52	GLU	N-CA	5.31	1.56	1.46
1	A	143	ILE	CA-CB	5.29	1.67	1.54
1	A	3	LEU	CA-CB	5.24	1.65	1.53
1	A	115	LEU	N-CA	5.20	1.56	1.46
1	A	58	PRO	CA-C	-5.20	1.42	1.52
1	A	148	GLU	CG-CD	-5.19	1.44	1.51
1	A	15	TRP	CG-CD1	5.17	1.44	1.36
1	A	133	ALA	CA-C	5.16	1.66	1.52
1	A	15	TRP	N-CA	5.14	1.56	1.46
1	A	134	TRP	CD2-CE2	-5.14	1.35	1.41
1	A	81	THR	CA-CB	5.12	1.66	1.53
1	A	79	GLU	CD-OE1	5.12	1.31	1.25
1	A	64	ALA	N-CA	5.11	1.56	1.46
1	A	57	ASN	C-O	5.09	1.33	1.23
1	A	138	TYR	N-CA	5.08	1.56	1.46
1	A	144	VAL	CA-C	5.08	1.66	1.52
1	A	38	PRO	N-CA	5.06	1.55	1.47
1	A	43	LEU	CA-CB	5.06	1.65	1.53
1	A	17	GLU	CB-CG	5.05	1.61	1.52
1	A	20	ALA	CA-CB	5.01	1.62	1.52

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	TYR	CB-CG-CD1	-9.86	115.09	121.00
1	A	112	GLU	OE1-CD-OE2	-9.81	111.52	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	GLU	OE1-CD-OE2	-8.52	113.07	123.30
1	A	16	GLU	OE1-CD-OE2	-8.04	113.66	123.30
1	A	28	ARG	NE-CZ-NH2	-7.90	116.35	120.30
1	A	15	TRP	CG-CD2-CE3	-7.64	127.03	133.90
1	A	138	TYR	CB-CG-CD2	7.04	125.22	121.00
1	A	46	PHE	CB-CG-CD2	6.99	125.69	120.80
1	A	87	ASP	CB-CG-OD2	-6.77	112.20	118.30
1	A	108	PRO	N-CA-CB	6.77	111.43	103.30
1	A	17	GLU	OE1-CD-OE2	-6.72	115.24	123.30
1	A	29	PHE	CB-CG-CD1	-6.70	116.11	120.80
1	A	128	GLU	OE1-CD-OE2	-6.52	115.47	123.30
1	A	120	GLU	OE1-CD-OE2	-6.52	115.47	123.30
1	A	54	PRO	N-CA-CB	6.46	111.05	103.30
1	A	126	TRP	CE2-CD2-CG	-6.41	102.17	107.30
1	A	29	PHE	CD1-CG-CD2	6.38	126.60	118.30
1	A	30	PHE	CB-CG-CD2	-6.28	116.41	120.80
1	A	72	TYR	CG-CD1-CE1	-6.24	116.31	121.30
1	A	134	TRP	CG-CD1-NE1	-6.17	103.93	110.10
1	A	23	PRO	N-CA-CB	6.14	110.67	103.30
1	A	48	LYS	CB-CA-C	-5.87	98.66	110.40
1	A	68	PHE	CB-CG-CD1	-5.84	116.71	120.80
1	A	126	TRP	NE1-CE2-CD2	5.79	113.09	107.30
1	A	138	TYR	CD1-CE1-CZ	5.79	125.01	119.80
1	A	15	TRP	CH2-CZ2-CE2	-5.79	111.61	117.40
1	A	15	TRP	CD2-CE3-CZ3	-5.53	111.61	118.80
1	A	15	TRP	CD1-NE1-CE2	-5.53	104.03	109.00
1	A	131	ASN	O-C-N	5.50	131.51	122.70
1	A	142	ALA	O-C-N	5.44	131.40	122.70
1	A	8	ALA	O-C-N	5.35	131.26	122.70
1	A	124	ALA	CB-CA-C	-5.30	102.14	110.10
1	A	15	TRP	NE1-CE2-CZ2	-5.18	124.70	130.40
1	A	76	ILE	O-C-N	5.17	130.98	122.70
1	A	149	MET	O-C-N	5.14	130.92	122.70
1	A	12	LYS	O-C-N	5.13	130.91	122.70
1	A	138	TYR	O-C-N	5.08	130.83	122.70
1	A	3	LEU	N-CA-CB	5.06	120.52	110.40
1	A	29	PHE	CB-CG-CD2	-5.02	117.29	120.80

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	151	ASP	Sidechain
1	A	17	GLU	Sidechain
1	A	19	ASN	Sidechain
1	A	42	ASP	Sidechain
1	A	57	ASN	Sidechain
1	A	7	GLN	Sidechain
1	A	87	ASP	Sidechain
1	A	92	ASN	Sidechain

CLOSE-CONTACTS INFOmissingINFO

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	153/153 (100%)	148 (97%)	3 (2%)	2 (1%)	12 6

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	ALA
1	A	3	LEU

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/125 (102%)	123 (97%)	4 (3%)	40 40

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	34	LEU
1	A	96	VAL
1	A	151	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	25	HIS
1	A	61	GLN
1	A	63	HIS
1	A	77	GLN

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.5 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	154	1,3	41,50,50	4.04	29 (70%)	45,82,82	2.31	20 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	154	1,3	-	1/12/54/54	-

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	154	HEM	C1D-C2D	7.53	1.59	1.44
2	A	154	HEM	FE-NB	7.21	2.32	1.96
2	A	154	HEM	C1B-C2B	7.11	1.58	1.44
2	A	154	HEM	C4D-C3D	6.87	1.56	1.45
2	A	154	HEM	CAB-C3B	6.41	1.64	1.47
2	A	154	HEM	C3B-C4B	6.36	1.57	1.44
2	A	154	HEM	CBD-CGD	6.31	1.65	1.50
2	A	154	HEM	CAA-C2A	6.06	1.60	1.52
2	A	154	HEM	C1A-NA	5.58	1.47	1.36
2	A	154	HEM	CAD-C3D	5.18	1.64	1.51
2	A	154	HEM	C3C-C2C	4.67	1.46	1.40
2	A	154	HEM	C3C-CAC	4.65	1.57	1.47
2	A	154	HEM	CMD-C2D	4.60	1.60	1.50
2	A	154	HEM	CMB-C2B	4.53	1.60	1.50
2	A	154	HEM	C4A-NA	3.98	1.44	1.36
2	A	154	HEM	FE-ND	3.95	2.16	1.96
2	A	154	HEM	CMC-C2C	3.78	1.60	1.51
2	A	154	HEM	CBA-CGA	3.69	1.59	1.50
2	A	154	HEM	C4A-CHB	3.64	1.51	1.41
2	A	154	HEM	O2A-CGA	-3.60	1.18	1.30
2	A	154	HEM	C2A-C3A	3.28	1.47	1.37
2	A	154	HEM	C1A-CHA	3.03	1.49	1.41
2	A	154	HEM	CMA-C3A	2.75	1.57	1.51
2	A	154	HEM	O1D-CGD	2.54	1.30	1.22
2	A	154	HEM	O2D-CGD	-2.45	1.22	1.30
2	A	154	HEM	CHB-C1B	2.43	1.41	1.35
2	A	154	HEM	CBD-CAD	-2.42	1.44	1.52
2	A	154	HEM	C4D-ND	2.42	1.44	1.40
2	A	154	HEM	C4B-NB	2.19	1.43	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	154	HEM	CMA-C3A-C4A	-5.13	120.58	128.46
2	A	154	HEM	C4C-CHD-C1D	4.44	128.41	122.56
2	A	154	HEM	C4A-C3A-C2A	4.28	109.97	107.00
2	A	154	HEM	C2C-C3C-C4C	3.78	109.54	106.90
2	A	154	HEM	C4B-C3B-C2B	-3.58	104.27	107.11
2	A	154	HEM	O2A-CGA-O1A	-3.46	114.68	123.30
2	A	154	HEM	C3C-C4C-NC	-3.29	104.74	110.94
2	A	154	HEM	O2D-CGD-O1D	-3.28	115.13	123.30
2	A	154	HEM	CHD-C1D-ND	-3.12	121.04	124.43
2	A	154	HEM	CMC-C2C-C3C	2.99	130.28	124.68
2	A	154	HEM	C4B-CHC-C1C	2.92	126.41	122.56
2	A	154	HEM	CHC-C4B-NB	-2.91	121.27	124.43
2	A	154	HEM	CAD-C3D-C4D	2.80	129.55	124.66
2	A	154	HEM	CHB-C1B-NB	-2.66	121.09	124.38
2	A	154	HEM	C3D-C4D-ND	2.53	112.98	110.17
2	A	154	HEM	CBA-CAA-C2A	-2.53	108.31	112.62
2	A	154	HEM	CMA-C3A-C2A	2.39	129.44	124.94
2	A	154	HEM	CHA-C4D-ND	-2.15	121.72	124.38
2	A	154	HEM	C2B-C1B-NB	2.07	112.29	109.84
2	A	154	HEM	C1B-NB-C4B	-2.07	102.94	105.07

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	154	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	154	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.6 Other polymers [i](#)

There are no such residues in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.