

Full wwPDB NMR Structure Validation Report (i)

Jun 4, 2023 – 10:03 AM EDT

PDB ID : 2LHW BMRB ID : 17869

Title : Tri-O-GalNAc glycosylated Mucin sequence based on MUC2 Mucin glycopro-

tein tandem repeat

Authors: Borgert, A.; Heimburg-Molinaro, J.; Lasanajak, Y.; Ju, T.; Liu, M.; Thomp-

son, P.; Ragupathi, G.; Barany, G.; Cummings, R.; Smith, D.; Live, D.

Deposited on : 2011-08-18

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

wwPDB-ShiftChecker : v1.2 BMRB Restraints Analysis : v1.2

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	m NMR archive $(# m Entries)$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	9	44%	44%	11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mal	Chain	Compound	Dec	Total mo	dels with violations
WIOI	Chain	Compound	res	Chirality	Geometry
2	A	A2G	11	2	-



2 Ensemble composition and analysis (i)

This entry contains 39 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms (4) was below the domain threshold value (8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust




3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 201 atoms, of which 103 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called MUC2 Mucin Domain Peptide.

Mol	Chain	Residues		At	oms			Trace
1	Λ	0	Total	С	Н	N	О	1
1	A	9	117	36	61	9	11	1

• Molecule 2 is 2-acetamido-2-deoxy-alpha-D-galactopyranose (three-letter code: A2G) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues		\mathbf{At}	oms		
2	Λ	1	Total	С	Н	N	О
2	A	1	28	8	14	1	5
2	Λ	1	Total	С	Н	N	О
	Λ	1	28	8	14	1	5
2	Λ	1	Total	С	Н	N	О
	Α	1	28	8	14	1	5



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 44% 44% 11%

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 44%




4.2.2 Score per residue for model 2

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 78% 22%







Chain A: 22% 44% 33%



4.2.4 Score per residue for model 4

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 22% 33% 33% 11%



4.2.5 Score per residue for model 5

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%



4.2.6 Score per residue for model 6

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%




4.2.7 Score per residue for model 7

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 22% 11%







Chain A: 44% 11% 44%



4.2.9 Score per residue for model 9

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 78% 22%



4.2.10 Score per residue for model 10

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 78% 11% 11%



4.2.11 Score per residue for model 11

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%




4.2.12 Score per residue for model 12


• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 44% 22% 33%











4.2.14 Score per residue for model 14

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 44% 22% 22% 11%



4.2.15 Score per residue for model 15

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 11% 22% 11%



4.2.16 Score per residue for model 16

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 33% 44% 22%




4.2.17 Score per residue for model 17

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%







Chain A: 67% 22% 11%



4.2.19 Score per residue for model 19

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 44% 44% 11%



4.2.20 Score per residue for model 20

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 22% 11%



4.2.21 Score per residue for model 21

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 33% 33% 33%




4.2.22 Score per residue for model 22

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%







Chain A: 78% 22%



4.2.24 Score per residue for model 24

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 22% 67% 11%



4.2.25 Score per residue for model 25

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 78% 22%



4.2.26 Score per residue for model 26

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 22% 11%




4.2.27 Score per residue for model 27

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 78% 22%







Chain A: 67% 22% 11%



4.2.29 Score per residue for model 29

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 89% 11%



4.2.30 Score per residue for model 30

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 22% 22%



4.2.31 Score per residue for model 31

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 33%




4.2.32 Score per residue for model 32

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 11% 22%







Chain A: 33% 56% 11%



4.2.34 Score per residue for model 34

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 67% 33%



4.2.35 Score per residue for model 35

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 33% 33% 33%



4.2.36 Score per residue for model 36

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 33% 56% 11%



4.2.37 Score per residue for model 37

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 33% 56% 11%





4.2.38 Score per residue for model 38

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 33% 11%



4.2.39 Score per residue for model 39

• Molecule 1: MUC2 Mucin Domain Peptide

Chain A: 56% 22% 22%





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: torsion space simulated annealing.

Of the 100 calculated structures, 39 were deposited, based on the following criterion: structures with the least restraint violations.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	142
Number of shifts mapped to atoms	142
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, NH2, ACE

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	56	61	59	3±2
2	A	42	42	36	1±1
All	All	3822	4017	3705	131

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:7:LEU:H	1:A:7:LEU:HD23	0.81	1.36	36	4
1:A:5:THR:O	1:A:7:LEU:N	0.80	2.14	31	10
1:A:7:LEU:HD23	1:A:7:LEU:N	0.71	2.00	36	7
1:A:7:LEU:HD12	1:A:7:LEU:H	0.66	1.50	20	3
1:A:7:LEU:HD12	1:A:7:LEU:N	0.64	2.08	27	3
1:A:6:PRO:O	1:A:7:LEU:C	0.63	2.36	39	9
1:A:7:LEU:H	1:A:7:LEU:CD2	0.63	2.06	21	4
1:A:5:THR:O	1:A:6:PRO:O	0.61	2.19	12	2
1:A:7:LEU:HD23	2:A:12:A2G:H8	0.60	1.74	31	1
1:A:6:PRO:O	1:A:8:LYS:N	0.56	2.39	4	10
1:A:6:PRO:C	1:A:8:LYS:H	0.56	2.04	37	8

Continued on next page...



Continued from previous page...

A + 1		Clasta (Å)	Distance (%)	Models		
Atom-1	Atom-2	$\operatorname{Clash}(\mathrm{\AA})$	$\operatorname{Distance}(\check{\mathbf{A}})$	Worst	Total	
1:A:7:LEU:N	1:A:7:LEU:CD2	0.56	2.69	13	8	
1:A:7:LEU:H	1:A:7:LEU:CD1	0.56	2.11	20	1	
1:A:7:LEU:N	1:A:7:LEU:HD22	0.55	2.16	2	2	
1:A:6:PRO:O	1:A:9:NH2:N	0.55	2.40	24	1	
1:A:5:THR:HG23	2:A:10:A2G:O7	0.55	2.02	15	8	
1:A:6:PRO:O	2:A:12:A2G:H8A	0.52	2.04	12	1	
1:A:7:LEU:HD21	2:A:12:A2G:O7	0.52	2.04	33	1	
1:A:7:LEU:O	1:A:9:NH2:N	0.51	2.43	35	3	
1:A:6:PRO:O	1:A:7:LEU:O	0.51	2.29	11	5	
1:A:7:LEU:N	1:A:7:LEU:CD1	0.49	2.75	27	3	
2:A:11:A2G:HO4	2:A:11:A2G:H15	0.49	1.50	5	1	
1:A:5:THR:CG2	2:A:10:A2G:O7	0.47	2.62	2	2	
1:A:7:LEU:CD2	2:A:12:A2G:O7	0.47	2.63	17	2	
1:A:7:LEU:HD21	2:A:12:A2G:C7	0.46	2.39	33	1	
1:A:7:LEU:CD1	2:A:12:A2G:O7	0.46	2.64	25	1	
1:A:6:PRO:C	1:A:8:LYS:N	0.45	2.70	6	8	
1:A:7:LEU:CD2	1:A:7:LEU:H	0.45	2.24	13	2	
1:A:7:LEU:C	1:A:9:NH2:N	0.45	2.69	35	1	
1:A:7:LEU:O	1:A:8:LYS:CB	0.44	2.66	12	5	
1:A:7:LEU:O	1:A:8:LYS:C	0.43	2.56	24	2	
2:A:11:A2G:O6	2:A:11:A2G:O4	0.43	2.35	23	1	
1:A:7:LEU:CD1	1:A:7:LEU:H	0.43	2.27	27	3	
1:A:7:LEU:CD2	2:A:12:A2G:H8	0.42	2.43	31	1	
1:A:1:ACE:O	1:A:2:PRO:O	0.42	2.38	13	5	
1:A:1:ACE:O	1:A:3:THR:HG23	0.42	2.14	4	1	
1:A:7:LEU:O	1:A:8:LYS:O	0.42	2.37	8	1	

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	7/9 (78%)	4±1 (59±14%)	1±1 (18±12%)	2±1 (23±14%)	0 1
All	All	273/351 (78%)	160 (59%)	50 (18%)	63 (23%)	0 1



All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	7	LEU	22
1	A	2	PRO	18
1	A	6	PRO	14
1	A	8	LYS	9

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric		Outliers	Percentiles	
1	A	7/7 (100%)	5±1 (77±15%)	2±1 (23±15%)	3	29
All	All	273/273 (100%)	211 (77%)	62 (23%)	3	29

All 5 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	5	THR	19
1	A	3	THR	16
1	A	7	LEU	12
1	A	8	LYS	9
1	A	4	THR	6

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.



6.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Pos	Link		Bond leng	ths
WIOI	туре	Chain	rtes	Lilik	Counts	RMSZ	#Z>2
2	A2G	A	11	1	14,14,15	0.64 ± 0.01	0±0 (0±0%)
2	A2G	A	10	1	14,14,15	0.64 ± 0.01	0±0 (0±0%)
2	A2G	A	12	1	14,14,15	0.65 ± 0.01	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

7	/[a]	Tuno	Chain	Dec	Tiple		Bond ang	gles
1	101	туре	Chain	nes	Lilik	Counts	RMSZ	#Z>2
	2	A2G	A	11	1	17,19,21	0.83 ± 0.01	0±0 (0±0%)
	2	A2G	A	10	1	17,19,21	0.83 ± 0.01	0±0 (0±0%)
	2	A2G	A	12	1	17,19,21	0.83 ± 0.01	0±0 (0±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A2G	A	12	1	-	$0\pm0,6,23,26$	$0\pm0,1,1,1$
2	A2G	A	11	1	-	$0\pm0,6,23,26$	$0\pm0,1,1,1$
2	A2G	A	10	1	-	$0\pm0,6,23,26$	$0\pm0,1,1,1$

There are no bond-length outliers.

All unique angle outliers are listed below.



Mol	$oxed{Chain} oxed{Res} oxed{Type} oxed{Atoms} oxed{Z} oxed{Observed}^{\hspace{-0.5em} o}$		Ideal(°)	Mod					
IVIOI	Cham	II ICS	Tees Type	11001115		Observed()	racar()	Worst	Total
2	A	12	A2G	O5-C1-C2	2.12	107.94	111.29	35	2

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	A	11	A2G	C1	2

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 91% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	142
Number of shifts mapped to atoms	142
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 85 atoms were assigned a chemical shift out of a possible 93. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	24/31 (77%)	12/12 (100%)	7/14 (50%)	5/5 (100%)
Sidechain	61/62 (98%)	41/41 (100%)	20/20 (100%)	0/1 (0%)
Overall	85/93 (91%)	53/53 (100%)	27/34 (79%)	5/6 (83%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 91%, i.e. 85 atoms were assigned a chemical shift out of a possible 93. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.



	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	24/31 (77%)	12/12 (100%)	7/14 (50%)	5/5 (100%)
Sidechain	61/62 (98%)	41/41 (100%)	20/20 (100%)	0/1 (0%)
Overall	85/93 (91%)	53/53 (100%)	27/34 (79%)	5/6 (83%)

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	4	THR	СВ	80.03	61.12 - 78.27	6.0
1	A	3	THR	СВ	79.75	61.12 - 78.27	5.9

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	98
Intra-residue ($ i-j =0$)	52
Sequential (i-j =1)	14
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range (i-j ≥5)	32
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	6
Number of unmapped restraints	0
Number of restraints per residue	11.6
Number of long range restraints per residue ¹	3.6

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.1	0.17
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

$\mathbf{Bins}\;(^{\circ})$	Average number of violations per model	\mathbf{Max} (°)
1.0-10.0 (Small)	0.7	2.7
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dogtosinta tuna	Count	$\%^{1}$	Vic	\mathbf{lated}^{5}	3	Consis	tentl	${f y}$ Violated 4
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	52	53.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	5	5.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	19	19.4	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	28	28.6	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	14	14.3	2	14.3	2.0	0	0.0	0.0
Backbone-Backbone	7	7.1	1	14.3	1.0	0	0.0	0.0
Backbone-Sidechain	5	5.1	1	20.0	1.0	0	0.0	0.0
Sidechain-Sidechain	2	2.0	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range ($ i-j \ge 5$)	32	32.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	14	14.3	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	18	18.4	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	98	100.0	2	2.0	2.0	0	0.0	0.0
Backbone-Backbone	12	12.2	1	8.3	1.0	0	0.0	0.0
Backbone-Sidechain	38	38.8	1	2.6	1.0	0	0.0	0.0
Sidechain-Sidechain	48	49.0	0	0.0	0.0	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	5	M (8)	M (Å)	\mathbf{SD}^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	$SD^*(A)$	Median (Å)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0
11	0	1	0	0	0	1	0.11	0.11	0.0	0.11

Continued on next page...




Continued from previous page...

M - 1-1 ID			nber o		ations	3	Ν .σ (&)	М (%)	CD6 (Å)	N/ - 1 (&)
Model ID	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (Å)	Max (Å)	${ m SD}^6 \ (m \AA)$	Median (Å)
12	0	0	0	0	0	0	0.0	0.0	0.0	0.0
13	0	1	0	0	0	1	0.13	0.13	0.0	0.13
14	0	0	0	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0	0	0	0.0	0.0	0.0	0.0
21	0	0	0	0	0	0	0.0	0.0	0.0	0.0
22	0	0	0	0	0	0	0.0	0.0	0.0	0.0
23	0	0	0	0	0	0	0.0	0.0	0.0	0.0
24	0	0	0	0	0	0	0.0	0.0	0.0	0.0
25	0	0	0	0	0	0	0.0	0.0	0.0	0.0
26	0	0	0	0	0	0	0.0	0.0	0.0	0.0
27	0	0	0	0	0	0	0.0	0.0	0.0	0.0
28	0	0	0	0	0	0	0.0	0.0	0.0	0.0
29	0	0	0	0	0	0	0.0	0.0	0.0	0.0
30	0	0	0	0	0	0	0.0	0.0	0.0	0.0
31	0	2	0	0	0	2	0.15	0.17	0.03	0.15
32	0	0	0	0	0	0	0.0	0.0	0.0	0.0
33	0	0	0	0	0	0	0.0	0.0	0.0	0.0
34	0	0	0	0	0	0	0.0	0.0	0.0	0.0
35	0	0	0	0	0	0	0.0	0.0	0.0	0.0
36	0	0	0	0	0	0	0.0	0.0	0.0	0.0
37	0	0	0	0	0	0	0.0	0.0	0.0	0.0
38	0	0	0	0	0	0	0.0	0.0	0.0	0.0
39	0	0	0	0	0	0	0.0	0.0	0.0	0.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation



9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 96(IR:52, SQ:12, MR:0, LR:32, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restr	aints	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Count ⁶	%		
0	1	0	0	0	1	1	2.6		
0	0	0	0	0	0	2	5.1		
0	1	0	0	0	1	3	7.7		
0	0	0	0	0	0	4	10.3		
0	0	0	0	0	0	5	12.8		
0	0	0	0	0	0	6	15.4		

Continued on next page...




Continued from previous page...

		$\frac{\textit{jrom pr}}{\mathbf{of vio}}$				Fraction	n of the ensemble
IR^1	SQ^2	MR^3		IC^5	Total	Count ⁶	%
0	0	0	0	0	0	7	17.9
0	0	0	0	0	0	8	20.5
0	0	0	0	0	0	9	23.1
0	0	0	0	0	0	10	25.6
0	0	0	0	0	0	11	28.2
0	0	0	0	0	0	12	30.8
0	0	0	0	0	0	13	33.3
0	0	0	0	0	0	14	35.9
0	0	0	0	0	0	15	38.5
0	0	0	0	0	0	16	41.0
0	0	0	0	0	0	17	43.6
0	0	0	0	0	0	18	46.2
0	0	0	0	0	0	19	48.7
0	0	0	0	0	0	20	51.3
0	0	0	0	0	0	21	53.8
0	0	0	0	0	0	22	56.4
0	0	0	0	0	0	23	59.0
0	0	0	0	0	0	24	61.5
0	0	0	0	0	0	25	64.1
0	0	0	0	0	0	26	66.7
0	0	0	0	0	0	27	69.2
0	0	0	0	0	0	28	71.8
0	0	0	0	0	0	29	74.4
0	0	0	0	0	0	30	76.9
0	0	0	0	0	0	31	79.5
0	0	0	0	0	0	32	82.1
0	0	0	0	0	0	33	84.6
0	0	0	0	0	0	34	87.2
0	0	0	0	0	0	35	89.7
0	0	0	0	0	0	36	92.3
0	0	0	0	0	0	37	94.9
0	0	0	0	0	0	38	97.4
0	0	0	0	0	0	39	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	${f Models}^1$	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,28)	1:A:8:LYS:H	1:A:7:LEU:HA	3	0.14	0.02	0.13

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram



9.5.2 Table: All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,28)	1:A:8:LYS:H	1:A:7:LEU:HA	31	0.17
(1,28)	1:A:8:LYS:H	1:A:7:LEU:HA	13	0.13
(1,98)	1:A:8:LYS:H	1:A:7:LEU:HB3	31	0.12
(1,98)	1:A:8:LYS:H	1:A:7:LEU:HG	31	0.12
(1,28)	1:A:8:LYS:H	1:A:7:LEU:HA	11	0.11



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

A1 - 4	Carret	$\%^{1}$	Vie	olated	3	Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\frac{1}{\%}$	Count	$\%^2$	$\%^1$
•	6	100.0	3	50.0	50.0	0	0.0	0.0
Total	6	100.0	3	50.0	50.0	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories



10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

M. J.LTD	N	umber of violations	N (0)	N /((0)	CD (0)	N. 1: (0)
Model ID		Total	$\mathbf{Mean} (^{\circ})$	$\mathbf{Max} \ (^{\circ})$	\mathbf{SD} (°)	$ \ \mathbf{Median} (^{\circ}) $
1	0	0	0.0	0.0	0.0	0.0
2	0	0	0.0	0.0	0.0	0.0
3	1	1	1.4	1.4	0.0	1.4
4	0	0	0.0	0.0	0.0	0.0
5	0	0	0.0	0.0	0.0	0.0
6	1	1	1.7	1.7	0.0	1.7
7	2	2	1.15	1.2	0.05	1.15
8	1	1	1.6	1.6	0.0	1.6
9	0	0	0.0	0.0	0.0	0.0
10	1	1	1.4	1.4	0.0	1.4
11	1	1	1.5	1.5	0.0	1.5
12	1	1	1.4	1.4	0.0	1.4
13	0	0	0.0	0.0	0.0	0.0
14	1	1	2.3	2.3	0.0	2.3
15	0	0	0.0	0.0	0.0	0.0
16	2	2	1.65	1.9	0.25	1.65
17	1	1	1.4	1.4	0.0	1.4
18	1	1	1.1	1.1	0.0	1.1
19	0	0	0.0	0.0	0.0	0.0
20	2	2	1.7	1.7	0.0	1.7
21	0	0	0.0	0.0	0.0	0.0
22	1	1	1.3	1.3	0.0	1.3
23	1	1	2.1	2.1	0.0	2.1
24	1	1	1.1	1.1	0.0	1.1
25	1	1	1.2	1.2	0.0	1.2
26	0	0	0.0	0.0	0.0	0.0
27	0	0	0.0	0.0	0.0	0.0
28	0	0	0.0	0.0	0.0	0.0
29	1	1	2.7	2.7	0.0	2.7
30	0	0	0.0	0.0	0.0	0.0
31	0	0	0.0	0.0	0.0	0.0
32	0	0	0.0	0.0	0.0	0.0
33	2	2	1.6	2.1	0.5	1.6
34	2	2	1.25	1.3	0.05	1.25
35	0	0	0.0	0.0	0.0	0.0
36	1	1	1.4	1.4	0.0	1.4
37	1	1	1.2	1.2	0.0	1.2
38	0	0	0.0	0.0	0.0	0.0


Continued on next page...



Continued from previous page...

Model ID		umber of violations Total	Mean (°)	Max (°)	SD (°)	Median (°)
39	0	0	0.0	0.0	0.0	0.0

10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

N	umber of violated restraints	Fraction of the ensemble			
	Total	Count ¹	%		
1	1	1	2.6		
0	0	2	5.1		
0	0	3	7.7		
0	0	4	10.3		
0	0	5	12.8		

Continued on next page...




 $Continued\ from\ previous\ page...$

	umber of violated restraints	Fraction of the ensemble			
	Total	Count ¹	%		
0	0	6	15.4		
0	0	7	17.9		
0	0	8	20.5		
0	0	9	23.1		
1	1	10	25.6		
0	0	11	28.2		
0	0	12	30.8		
0	0	13	33.3		
0	0	14	35.9		
1	1	15	38.5		
0	0	16	41.0		
0	0	17	43.6		
0	0	18	46.2		
0	0	19	48.7		
0	0	20	51.3		
0	0	21	53.8		
0	0	22	56.4		
0	0	23	59.0		
0	0	24	61.5		
0	0	25	64.1		
0	0	26	66.7		
0	0	27	69.2		
0	0	28	71.8		
0	0	29	74.4		
0	0	30	76.9		
0	0	31	79.5		
0	0	32	82.1		
0	0	33	84.6		
0	0	34	87.2		
0	0	35	89.7		
0	0	36	92.3		
0	0	37	94.9		
0	0	38	97.4		
0	0	39	100.0		

 $^{^{\}rm 1}$ Number of models with violations



10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	15	1.56	0.38	1.4
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	10	1.34	0.17	1.35


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,3)	1:A:5:THR:HA	1:A:5:THR:CA	1:A:5:THR:CB	1:A:5:THR:HB	29	2.7
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	14	2.3
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	23	2.1
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	33	2.1
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	16	1.9
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	20	1.7
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	6	1.7
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	20	1.7
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	8	1.6
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	11	1.5
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	12	1.4
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	16	1.4
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	17	1.4
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	3	1.4
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	10	1.4
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	36	1.4
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	22	1.3
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	34	1.3
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	25	1.2
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	7	1.2
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	34	1.2

Continued on next page...



$Continued\ from\ previous\ page...$

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	37	1.2
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	7	1.1
(1,2)	1:A:4:THR:HA	1:A:4:THR:CA	1:A:4:THR:CB	1:A:4:THR:HB	33	1.1
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	18	1.1
(1,1)	1:A:3:THR:HA	1:A:3:THR:CA	1:A:3:THR:CB	1:A:3:THR:HB	24	1.1

