

Full wwPDB X-ray Structure Validation Report (i)

Oct 9, 2023 – 10:43 PM EDT

PDB ID : 7LPF

Title: The internal aldimine form of the wild-type Salmonella typhimurium Trypto-

phan Synthase in complex with inhibitor N-(4'-trifluoromethoxybenzenesulf onyl)-2-amino-1-ethylphosphate (F9F) at the enzyme alpha-site and sodium

ion at the metal coordination site at 1.10 Angstrom resolution

Authors: Hilario, E.; Dunn, M.F.; Mueller, L.J.

Deposited on : 2021-02-11

Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

 $Xtriage\ (Phenix) \quad : \quad 1.13$

EDS : 2.35.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

\mathbf{M}	ol	Chain	Length	Quality of chain				
-		A	240	20%				
	-	A	268	87%	13%			
		ъ	20-	14%				
2	2	В	397	89%	10% •			

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	F9F	A	301	-	-	X	-
4	PEG	A	302	-	-	X	-



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 6031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Tryptophan synthase alpha chain.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	Δ	268	Total	С	N	О	S	0	19	0
1	11	200	2086	1316	366	395	9		12	

• Molecule 2 is a protein called Tryptophan synthase beta chain.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	393	Total 3107	C 1947	N 546	O 592	S 22	0	16	0


• Molecule 3 is 2-({[4-(TRIFLUOROMETHOXY)PHENYL]SULFONYL}AMINO)ETHYL DIHYDROGEN PHOSPHATE (three-letter code: F9F) (formula: C₉H₁₁F₃NO₇PS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	F	N	О	Р	S	0	0
)	A	1	22	9	3	1	7	1	1	U	0




• Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
4	A	1	Total 7	C 4	O 3	0	0


 \bullet Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O S 4 2 1 1	0	0
5	В	1	Total C O S 4 2 1 1	0	0




• Molecule 6 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).



Mo	Chain	Residues	Atoms			ZeroOcc	AltConf		
6	В	1	Total	С	N	Ō	P	0	0
			15	8	1	Б	1		

 \bullet Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	В	1	Total C O 4 2 2	0	0



• Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	Total Na 1 1	0	0

• Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	296	Total O 301 301	0	4
9	В	469	Total O 480 480	0	11



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan synthase alpha chain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	182.23Å 59.18Å 67.33Å	Depositor
a, b, c, α , β , γ	90.00° 94.70° 90.00°	Depositor
Resolution (Å)	28.52 - 1.10	Depositor
resolution (A)	28.50 - 1.10	EDS
% Data completeness	96.3 (28.52-1.10)	Depositor
(in resolution range)	96.3 (28.50-1.10)	EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) > 1$	1.44 (at 1.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
P.P.	0.172 , 0.199	Depositor
R, R_{free}	0.180 , 0.208	DCC
R_{free} test set	13730 reflections $(4.94%)$	wwPDB-VP
Wilson B-factor (Å ²)	7.2	Xtriage
Anisotropy	1.776	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.46, 85.1	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6031	wwPDB-VP
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.45% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG, NA, F9F, PLP, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.63	0/2127	0.73	0/2891
2	В	0.65	0/3166	0.76	0/4272
All	All	0.64	0/5293	0.75	0/7163

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2086	0	2049	45	0
2	В	3107	0	3054	38	0
3	A	22	0	9	16	0
4	A	7	0	10	12	0
5	A	4	0	6	0	0
5	В	4	0	6	1	0
6	В	15	0	6	0	0
7	В	4	0	6	0	0
8	В	1	0	0	0	0
9	A	301	0	0	10	0
9	В	480	0	0	13	0
All	All	6031	0	5146	87	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 8.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
1 A 224[D] CIV (9 A 901 DOD D17	distance (Å)	overlap (Å)
1:A:234[B]:GLY:O	3:A:301:F9F:P17	2.03	1.17
1:A:218:GLU:HG3	9:A:615:HOH:O	1.50	1.07
1:A:234[B]:GLY:O	3:A:301:F9F:O19	1.73	1.05
1:A:234[B]:GLY:O	3:A:301:F9F:O20	1.76	1.03
1:A:178:SER:OG	1:A:211[B]:GLY:O	1.84	0.94
1:A:234[A]:GLY:HA3	9:A:416:HOH:O	1.70	0.91
2:B:217:LEU:HD11	9:B:927:HOH:O	1.76	0.83
1:A:173[A]:TYR:OH	9:A:401:HOH:O	1.76	0.79
1:A:234[B]:GLY:HA3	9:A:416:HOH:O	1.83	0.79
1:A:42[B]:ASP:OD1	9:A:402:HOH:O	2.05	0.74
2:B:181[B]:TYR:CE2	9:B:638:HOH:O	2.40	0.74
1:A:49[B]:GLU:OE1	3:A:301:F9F:S12	2.46	0.72
2:B:217:LEU:HG	9:B:774:HOH:O	1.90	0.71
1:A:64:ILE:HD11	9:A:416:HOH:O	1.90	0.71
2:B:27:GLN:O	2:B:30:GLU:HG3	1.92	0.70
3:A:301:F9F:H151	9:A:416:HOH:O	1.90	0.70
4:A:302:PEG:H21	2:B:16:TYR:OH	1.93	0.69
1:A:153:ILE:HG23	1:A:177:LEU:HG	1.76	0.68
2:B:181[B]:TYR:CZ	9:B:638:HOH:O	2.47	0.67
4:A:302:PEG:H32	2:B:283:LYS:NZ	2.11	0.65
1:A:53:PRO:HA	1:A:68:ASN:HD22	1.62	0.65
1:A:175:TYR:CE2	3:A:301:F9F:H5	2.32	0.64
2:B:44:GLN:HG3	9:B:718:HOH:O	1.98	0.64
1:A:154[B]:CYS:SG	1:A:166:VAL:HG11	2.38	0.64
1:A:153:ILE:HD11	3:A:301:F9F:H6	1.80	0.63
1:A:234[B]:GLY:CA	9:A:416:HOH:O	2.42	0.63
2:B:217:LEU:HD21	9:B:927:HOH:O	1.99	0.62
1:A:49[B]:GLU:OE1	3:A:301:F9F:O22	2.16	0.62
1:A:49[B]:GLU:OE1	3:A:301:F9F:O21	2.19	0.61
1:A:185:ALA:O	1:A:188:ARG:NH1	2.37	0.58
1:A:53:PRO:HA	1:A:68:ASN:ND2	2.18	0.57
1:A:49[B]:GLU:CD	3:A:301:F9F:O21	2.42	0.57
2:B:99:LYS:HZ1	2:B:126:LEU:HD13	1.69	0.57
1:A:153:ILE:CG2	1:A:177:LEU:HG	2.33	0.57
2:B:148:ARG:O	2:B:152[A]:MET:HE2	2.06	0.56
2:B:216[B]:ILE:HG22	2:B:222:ARG:O	2.07	0.53
1:A:138:PRO:HG2	4:A:302:PEG:H11	1.90	0.53



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
2:B:341[A]:ARG:HD2	9:B:541:HOH:O	2.08	0.53
4:A:302:PEG:C3	2:B:283:LYS:HZ1	2.22	0.53
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.45	0.51
2:B:223:LEU:HD23	9:B:774:HOH:O	2.10	0.51
1:A:234[B]:GLY:C	9:A:416:HOH:O	2.48	0.51
1:A:225[B]:ARG:HB2	1:A:225[B]:ARG:CZ	2.40	0.50
4:A:302:PEG:H32	2:B:283:LYS:HZ1	1.75	0.50
2:B:117:VAL:HG13	2:B:152[A]:MET:HE1	1.93	0.50
2:B:99:LYS:NZ	2:B:126:LEU:HD13	2.27	0.49
1:A:100:LEU:C	1:A:100:LEU:HD13	2.34	0.48
2:B:143:SER:OG	2:B:144:PRO:HD3	2.13	0.48
1:A:156:PRO:HA	1:A:196:LEU:HD11	1.96	0.48
1:A:22:PHE:HA	1:A:49[B]:GLU:O	2.13	0.48
2:B:216[A]:ILE:HG23	9:B:774:HOH:O	2.13	0.48
2:B:99:LYS:HE3	2:B:126:LEU:HD22	1.96	0.47
1:A:139:PHE:HE1	4:A:302:PEG:C1	2.28	0.47
2:B:96:LEU:HD23	2:B:99:LYS:HE2	1.97	0.47
2:B:103:LYS:NZ	2:B:181[B]:TYR:O	2.40	0.46
2:B:379[B]:ARG:NH2	9:B:751[B]:HOH:O	2.12	0.46
2:B:159:VAL:HG13	2:B:172[A]:GLU:HG3	1.97	0.46
1:A:178:SER:HB3	1:A:189:GLY:HA3	1.98	0.46
1:A:183:THR:CG2	3:A:301:F9F:H142	2.46	0.45
1:A:153:ILE:HD13	1:A:175:TYR:HB3	1.98	0.45
2:B:300:ILE:HD11	2:B:390:ILE:CD1	2.46	0.45
3:A:301:F9F:H3	9:A:497:HOH:O	2.15	0.45
1:A:111:ILE:CD1	4:A:302:PEG:H31	2.47	0.45
4:A:302:PEG:H22	4:A:302:PEG:H42	1.71	0.45
2:B:159:VAL:CG1	2:B:172[A]:GLU:HG3	2.48	0.44
1:A:211[A]:GLY:O	1:A:212:PHE:HB2	2.18	0.44
2:B:303:GLY:HA3	2:B:350:GLU:OE2	2.17	0.44
2:B:275:ARG:NH2	9:B:510:HOH:O	2.51	0.43
1:A:195:HIS:O	1:A:199:LYS:HG2	2.17	0.43
5:B:403:DMS:H11	9:B:846:HOH:O	2.18	0.43
4:A:302:PEG:C2	2:B:283:LYS:HZ1	2.31	0.43
2:B:379[B]:ARG:HD2	2:B:381:ASP:OD1	2.18	0.43
1:A:175:TYR:HE2	3:A:301:F9F:O22	2.02	0.42
2:B:360[A]:LYS:HG3	9:B:653:HOH:O	2.18	0.42
1:A:233[B]:SER:O	3:A:301:F9F:O20	2.38	0.42
1:A:139:PHE:HE1	4:A:302:PEG:H12	1.83	0.42
1:A:127:LEU:HD11	1:A:153:ILE:HD11	2.01	0.42
1:A:175:TYR:CE2	3:A:301:F9F:O22	2.73	0.42



I 'omtamalod	tmom	mmonia	maaa
Continued	110111	DIEUIUU	DUJUE
00,000,000	.,	p	p = 9 =

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\left(\operatorname{\mathring{A}} ight)$	overlap (Å)
1:A:22:PHE:HA	1:A:49[A]:GLU:O	2.19	0.41
1:A:111:ILE:HD11	4:A:302:PEG:H31	2.01	0.41
2:B:95:ALA:C	2:B:99:LYS:HZ3	2.23	0.41
2:B:193:GLY:HA2	2:B:280:PHE:O	2.21	0.40
1:A:183:THR:HG21	3:A:301:F9F:H142	2.02	0.40
2:B:16:TYR:O	2:B:281:GLY:HA2	2.22	0.40
2:B:86:HIS:CE1	2:B:236:ASN:HB3	2.55	0.40
1:A:54:PHE:H	1:A:68:ASN:HD22	1.69	0.40
4:A:302:PEG:H12	2:B:16:TYR:HE1	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	278/268 (104%)	269 (97%)	7 (2%)	2 (1%)	22	4
2	В	407/397 (102%)	398 (98%)	9 (2%)	0	100	100
All	All	685/665 (103%)	667 (97%)	16 (2%)	2 (0%)	47	15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	235[A]	SER
1	A	235[B]	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	F	erce	entiles	3
1	A	212/208 (102%)	212 (100%)	0		100	100	
2	В	322/311 (104%)	321 (100%)	1 (0%)		92	76	
All	All	534/519 (103%)	533 (100%)	1 (0%)		93	77	

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	В	207	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	ASN
1	A	109	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with $ Z > 2$ is considered an outlier worth inspection.
RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	eles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	F9F	A	301	-	22,22,22	0.39	0	32,33,33	0.55	0
5	DMS	В	403	-	3,3,3	0.22	0	3,3,3	0.21	0
5	DMS	A	303	-	3,3,3	0.21	0	3,3,3	0.10	0
4	PEG	A	302	-	6,6,6	0.29	0	5,5,5	0.46	0
7	EDO	В	402	-	3,3,3	0.47	0	2,2,2	0.46	0
6	PLP	В	401	2	15,15,16	1.08	1 (6%)	20,22,23	1.35	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	F9F	A	301	-	-	7/20/20/20	0/1/1/1
6	PLP	В	401	2	-	0/6/6/8	0/1/1/1
7	EDO	В	402	-	-	0/1/1/1	-
4	PEG	A	302	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
6	В	401	PLP	C4A-C4	-2.69	1.46	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
6	В	401	PLP	O4P-C5A-C5	3.09	115.23	109.35
6	В	401	PLP	C6-C5-C4	-3.02	115.78	118.16
6	В	401	PLP	O3P-P-O4P	-2.01	101.38	106.73

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	PEG	C4-C3-O2-C2
4	A	302	PEG	O2-C3-C4-O4
4	A	302	PEG	O1-C1-C2-O2
3	A	301	F9F	C3-C4-S12-O21



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	301	F9F	C5-C4-S12-O21
3	A	301	F9F	C14-N13-S12-O22
3	A	301	F9F	C3-C4-S12-N13
3	A	301	F9F	C6-C1-O7-C8
3	A	301	F9F	C2-C1-O7-C8
3	A	301	F9F	C5-C4-S12-N13


There are no ring outliers.

3 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	F9F	16	0
5	В	403	DMS	1	0
4	A	302	PEG	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ	Z>:	2	$OWAB(A^2)$	Q < 0.9
1	A	268/268 (100%)	1.39	54 (20%)	1	2	9, 18, 41, 71	0
2	В	393/397~(98%)	1.12	54 (13%)	3	4	8, 12, 26, 62	0
All	All	661/665 (99%)	1.23	108 (16%)	1	3	8, 15, 34, 71	0

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	190	ALA	14.2
1	A	191	LEU	14.0
2	В	393	ALA	10.0
1	A	192	PRO	9.6
2	В	391	LEU	9.4
1	A	193	LEU	8.6
2	В	392	LYS	8.5
1	A	157	ASN	8.3
2	В	390	ILE	6.6
1	A	194	HIS	6.2
2	В	385	PHE	5.8
1	A	185	ALA	5.8
1	A	195	HIS	5.8
1	A	187	ASN	5.1
1	A	158	ALA	4.8
1	A	186	GLU	4.7
2	В	394	ARG	4.6
1	A	182	VAL	4.5
1	A	13	ASP	4.2
1	A	268	ALA	4.1
1	A	203	TYR	3.9
1	A	160	ASP	3.9
2	В	389	ASP	3.9
1	A	161	ASP	3.9



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	A	73	ALA	3.8
2	В	388	HIS	3.7
1	A		159 ASP	
1	A	234[A]	GLY	3.7
1	A	15	ARG	3.6
2	В	2	THR	3.6
	В	160	HIS	3.6
2	A	196	LEU	3.6
1	A	188	ARG	3.5
2	В	140	GLU	3.5
1	A	189	GLY	3.5
2	В	162	GLY	3.5
1	A	74	ALA	3.4
1	A	250	GLN	3.4
2	В	159	VAL	3.4
1	A	246	ALA	3.4
2	В	163	SER	3.3
1	A	253	ALA	3.3
2	В	157	ILE	3.3
1	A	204	HIS	3.2
2	В	374	VAL	3.2
1	A	109	ASN	3.1
1	A	156	PRO	3.1
1	A	180	SER	3.1
2	В	136	ALA	3.0
2	В	139	VAL	2.9
2	В	386	THR	2.9
2	В	137	LYS	2.9
2	В	228	ILE	2.8
2	В	238	ILE	2.8
2	В	143	SER	2.8
2	В	181[A]	TYR	2.8
2	В	237	ALA	2.8
2	В	141	ARG	2.8
2	В	229	ALA	2.7
2	В	253	ILE	2.7
2	В	147	PHE	2.7
2	В	286[A]	MET	2.7
2	В	355	LEU	2.7
1	A	214	ILE	2.7
1	A	75	GLY	2.6
1	A	184	GLY	2.6



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	RSRZ
2	В	372	LEU	2.6
1	A	257	257 SER	
2	В	179	179 GLY	
1	A	164	ARG	2.6
2	В	230	CYS	2.6
1	A	151	ILE	2.5
1	A	247	SER	2.5
1	A	235[A]	SER	2.5
1	A	249	LYS	2.5
2	В	73	LEU	2.5
1	A	12	ASN	2.5
2	В	158	PRO	2.5
1	A	202	GLU	2.4
2	В	75	LEU	2.4
1	A	183	THR	2.4
2	В	255	VAL	2.3
2	В	231	VAL	2.3
2	В	290	ALA	2.3
1	A	224	VAL	2.3
1	A	181	GLY	2.3
1	A	50	LEU	2.2
1	A	267	ARG	2.2
2	В	376	LEU	2.2
1	A	199	LYS	2.2
2	В	164	ALA	2.2
2	В	39	PRO	2.2
2	В	325	VAL	2.2
2	В	58	ALA	2.2
2	В	373	VAL	2.2
1	A	62	PRO	2.2
2	В	198	PRO	2.2
1	A	173[A]	TYR	2.1
1	A	260[A]	SER	2.1
2	В	316	LEU	2.1
2	В	144	PRO	2.1
1	A	9	ALA	2.1
2	В	264	THR	2.1
2	В	197	TYR	2.1
2	В	200	ILE	2.0
2	В	180	SER	2.0
1	A	99	LEU	2.0
2	В	176	ASP	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
7	EDO	В	402	4/4	0.88	0.14	16,25,25,36	0
5	DMS	A	303	4/4	0.93	0.15	37,38,39,42	0
3	F9F	A	301	22/22	0.94	0.14	18,20,26,28	0
4	PEG	A	302	7/7	0.95	0.14	23,28,35,40	0
5	DMS	В	403	4/4	0.97	0.13	20,23,25,28	0
6	PLP	В	401	15/16	0.99	0.10	9,10,12,13	0
8	NA	В	404	1/1	0.99	0.08	13,13,13,13	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around F9F A 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PLP B 401:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

