

# Full wwPDB X-ray Structure Validation Report (i)

#### Feb 3, 2024 – 12:34 PM EST

:	1LUG
:	Full Matrix Error Analysis of Carbonic Anhydrase
:	Merritt, E.A.; Le Trong, I.; Behnke, C.A.
:	2002-05-22
:	0.95  Å(reported)
	: : :

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 0.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
Clashscore	141614	1321 (1.06-0.86)
Ramachandran outliers	138981	1233 (1.06-0.86)
Sidechain outliers	138945	1235 (1.06-0.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	259	80%	16%	•••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

M	l Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SUA	А	1002[B]	-	Х	-	-



#### 1LUG

# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 2516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Carbonic Anhydrase II.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	259	Total 2120	C 1361	N 363	O 393	${ m S} { m 3}$	0	16	0


• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Ator	ns	ZeroOcc	AltConf
2	А	1	Total 1	Zn 1	0	0

• Molecule 3 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Hg 1 1	0	0


• Molecule 4 is MERCURIBENZOIC ACID (three-letter code: MBO) (formula: C<sub>7</sub>H<sub>5</sub>HgO<sub>2</sub>).






Mol	Chain	Residues	L	Ator	ns		ZeroOcc	AltConf
4	А	1	Total 20	C 14	Hg 2	0 4	0	1

• Molecule 5 is (4-SULFAMOYL-PHENYL)-THIOCARBAMIC ACID O-(2-THIOPHEN-3-Y L-ETHYL) ESTER (three-letter code: SUA) (formula:  $C_{13}H_{14}N_2O_3S_3$ ).



Mol	Chain	Residues		Atc	$\mathbf{ms}$			ZeroOcc	AltConf
5	А	1	Total	С	N	0	S	0	1
			18	11	2	3	2		

• Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total         C         O           12         6         6	0	1
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 7 is water.

$\mathbf{N}$	ſol	Chain	Residues	Atoms	ZeroOcc	AltConf
	7	А	337	Total         O           338         338	0	1




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Carbonic Anhydrase II





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	42.06Å 41.31Å 71.87Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $104.23^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	15.00 - 0.95	Depositor
% Data completeness	(Not available) (15.00-0.95)	Depositor
(in resolution range)	(1007 available) (15.00-0.55)	Depositor
$R_{merge}$	0.07	Depositor
R <sub>sym</sub>	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
$R, R_{free}$	0.119 , $0.141$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2516	wwPDB-VP
Average B, all atoms $(Å^2)$	14.0	wwPDB-VP



# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SUA, ZN, HG, MBO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Cl	Chain	Bond lengths		Bond angles	
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	1.39	26/2251~(1.2%)	1.75	70/3045~(2.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	27	ARG	CD-NE	-10.91	1.27	1.46
1	А	14	GLU	CD-OE2	10.06	1.36	1.25
1	А	2	SER	CB-OG	8.99	1.53	1.42
1	А	3	HIS	C-O	8.13	1.38	1.23
1	А	151	SER	CB-OG	-7.61	1.32	1.42
1	А	231	ASN	C-N	6.71	1.45	1.33
1	А	3	HIS	CA-C	6.70	1.70	1.52
1	А	233	GLU	CG-CD	-6.62	1.42	1.51
1	А	2	SER	CA-CB	6.55	1.62	1.52
1	А	158	LYS	CE-NZ	6.38	1.65	1.49
1	А	213	GLU	CD-OE1	-6.22	1.18	1.25
1	А	80	LYS	CB-CG	-6.15	1.35	1.52
1	А	165	SER	CB-OG	-6.09	1.34	1.42
1	А	76	LYS	C-N	-6.07	1.20	1.34
1	А	181	ARG	CZ-NH1	6.04	1.41	1.33
1	А	237	GLU	CD-OE1	5.98	1.32	1.25
1	А	129[A]	ASP	CA-CB	-5.95	1.40	1.53
1	А	129[B]	ASP	CA-CB	-5.95	1.40	1.53

All (26) bond length outliers are listed below:



Continued on next page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	9	LYS	CA-CB	5.46	1.66	1.53
1	А	232	GLY	CA-C	5.35	1.60	1.51
1	А	19	ASP	CG-OD1	-5.30	1.13	1.25
1	А	89	ARG	CZ-NH2	-5.29	1.26	1.33
1	А	220	GLU	CD-OE1	-5.21	1.20	1.25
1	А	3	HIS	CA-CB	5.18	1.65	1.53
1	А	112[A]	LYS	CG-CD	-5.14	1.34	1.52
1	А	112[B]	LYS	CG-CD	-5.14	1.34	1.52

Continued from previous page...

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	27	ARG	NE-CZ-NH1	17.63	129.12	120.30
1	А	2	SER	C-N-CA	15.27	159.88	121.70
1	А	164	ASP	CB-CG-OD1	15.27	132.04	118.30
1	А	58	ARG	NE-CZ-NH1	15.27	127.93	120.30
1	А	27	ARG	NE-CZ-NH2	-13.99	113.31	120.30
1	А	3	HIS	CA-C-O	-13.01	92.78	120.10
1	А	3	HIS	CA-C-N	12.92	145.63	117.20
1	А	164	ASP	CB-CG-OD2	-12.73	106.84	118.30
1	А	181	ARG	NE-CZ-NH2	-11.93	114.33	120.30
1	А	19	ASP	CB-CG-OD1	10.88	128.09	118.30
1	А	3	HIS	N-CA-CB	10.52	129.54	110.60
1	А	27	ARG	CG-CD-NE	10.43	133.71	111.80
1	А	225	PHE	CB-CG-CD1	9.46	127.42	120.80
1	А	251[A]	LYS	CA-CB-CG	9.34	133.95	113.40
1	А	251[B]	LYS	CA-CB-CG	9.34	133.95	113.40
1	А	161	ASP	CB-CG-OD2	-9.30	109.93	118.30
1	А	3	HIS	CB-CA-C	-9.12	92.17	110.40
1	А	27	ARG	CD-NE-CZ	8.96	136.15	123.60
1	А	58	ARG	NE-CZ-NH2	-8.60	116.00	120.30
1	А	14	GLU	OE1-CD-OE2	8.29	133.25	123.30
1	А	14	GLU	CG-CD-OE1	-7.97	102.36	118.30
1	А	75[A]	ASP	N-CA-CB	-7.64	96.85	110.60
1	А	75[B]	ASP	N-CA-CB	-7.64	96.85	110.60
1	А	171	LYS	CD-CE-NZ	7.63	129.25	111.70
1	А	76	LYS	C-N-CA	7.44	140.29	121.70
1	А	3	HIS	CG-ND1-CE1	7.29	118.41	108.20
1	А	181	ARG	NE-CZ-NH1	7.20	123.90	120.30
1	А	89	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	А	129[A]	ASP	CB-CA-C	6.87	124.14	110.40
1	А	129[B]	ASP	CB-CA-C	6.87	124.14	110.40

Continued on next page...



1 T	TIC	7
ΤL	100	J

	Continued from previous page										
Mol	Chain	$\operatorname{Res}$	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$				
1	А	220	GLU	OE1-CD-OE2	-6.83	115.11	123.30				
1	А	3	HIS	ND1-CG-CD2	-6.69	96.63	106.00				
1	А	2	SER	CA-C-N	6.58	131.68	117.20				
1	А	225	PHE	CB-CG-CD2	-6.55	116.22	120.80				
1	А	224	LYS	CB-CG-CD	6.55	128.62	111.60				
1	А	76	LYS	O-C-N	-6.27	112.67	122.70				
1	А	3	HIS	C-N-CA	6.26	137.35	121.70				
1	А	74	GLN	CG-CD-OE1	-6.20	109.20	121.60				
1	А	126	LYS	CD-CE-NZ	-6.12	97.61	111.70				
1	А	53	GLN	CA-CB-CG	-6.07	100.06	113.40				
1	А	55	THR	CA-CB-CG2	-5.96	104.05	112.40				
1	А	64[A]	HIS	CG-ND1-CE1	-5.93	97.99	105.70				
1	А	64[B]	HIS	CG-ND1-CE1	-5.93	97.99	105.70				
1	А	129[A]	ASP	CA-CB-CG	-5.87	100.48	113.40				
1	А	129[B]	ASP	CA-CB-CG	-5.87	100.48	113.40				
1	А	114	TYR	CB-CG-CD1	5.86	124.51	121.00				
1	А	227	LYS	CG-CD-CE	5.85	129.46	111.90				
1	А	52	ASP	CB-CG-OD1	5.85	123.57	118.30				
1	А	245	ARG	NE-CZ-NH1	5.81	123.21	120.30				
1	А	34[A]	ASP	CB-CG-OD2	5.76	123.48	118.30				
1	А	34[B]	ASP	CB-CG-OD2	5.76	123.48	118.30				
1	А	167	LYS	CD-CE-NZ	-5.71	98.57	111.70				
1	А	74	GLN	OE1-CD-NE2	5.65	134.89	121.90				
1	А	2	SER	CA-CB-OG	-5.64	95.96	111.20				
1	А	89	ARG	NH1-CZ-NH2	-5.61	113.23	119.40				
1	А	236	PRO	CA-N-CD	-5.50	103.80	111.50				
1	А	85[A]	ASP	CB-CG-OD2	5.40	123.16	118.30				
1	А	85[B]	ASP	CB-CG-OD2	5.40	123.16	118.30				
1	А	235	GLU	OE1-CD-OE2	-5.30	116.94	123.30				
1	А	51	TYR	CB-CG-CD1	5.30	124.18	121.00				
1	А	85[A]	ASP	N-CA-CB	5.30	120.13	110.60				
1	А	85[B]	ASP	N-CA-CB	5.30	120.13	110.60				
1	А	175	PHE	CB-CG-CD2	-5.14	117.20	120.80				
1	А	232	GLY	O-C-N	5.12	130.90	122.70				
1	А	19	ASP	OD1-CG-OD2	-5.12	113.58	123.30				
1	А	76	LYS	CA-C-N	5.12	128.45	117.20				
1	А	89	ARG	NE-CZ-NH2	5.11	122.86	120.30				
1	А	8	GLY	O-C-N	-5.02	114.67	122.70				
1	А	174	ASP	CB-CG-OD1	5.01	122.81	118.30				
1	А	132	LYS	CA-CB-CG	5.00	124.41	113.40				

Continued from previous page...

There are no chirality outliers.



All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	2	SER	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2120	0	4	8	0
2	А	1	0	0	0	0
3	А	1	0	0	0	0
4	А	20	0	0	1	0
5	А	18	0	6	2	0
6	А	18	0	23	4	0
7	А	338	0	0	5	0
All	All	2516	0	33	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1205[B]:MBO:CE1	4:A:1205[B]:MBO:HG	1.91	0.77
6:A:1004:GOL:H2	7:A:2255:HOH:O	2.04	0.58
1:A:92:GLN:NE2	5:A:1002[A]:SUA:H91	2.23	0.54
1:A:112[B]:LYS:NZ	7:A:2228:HOH:O	2.45	0.49
1:A:80:LYS:NZ	7:A:2111:HOH:O	2.45	0.49
1:A:64[A]:HIS:CE1	7:A:2214:HOH:O	2.65	0.49
1:A:92:GLN:CG	5:A:1002[A]:SUA:H91	2.44	0.47
1:A:14:GLU:OE1	6:A:1004:GOL:H32	2.14	0.47
6:A:1004:GOL:C2	7:A:2255:HOH:O	2.61	0.47
1:A:164:ASP:OD1	1:A:164:ASP:N	2.48	0.46
1:A:94:HIS:CG	6:A:1003[B]:GOL:H2	2.55	0.41

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	271/259~(105%)	259~(96%)	11 (4%)	1 (0%)	34 11

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	3	HIS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/224~(107%)	230~(96%)	9~(4%)	33 5	

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	3	HIS
1	А	52	ASP
1	А	64[A]	HIS
1	А	64[B]	HIS
1	А	74	GLN
1	А	92	GLN
1	А	164	ASP
1	А	260[A]	LYS
1	А	260[B]	LYS



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	gles
WIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
6	GOL	А	1003[B]	-	$5,\!5,\!5$	1.18	0	$5,\!5,\!5$	1.21	1 (20%)
5	SUA	А	1002[A]	3	$17,\!17,\!22$	3.75	3 (17%)	23,23,30	6.05	10 (43%)
5	SUA	А	1002[B]	-	$17,\!12,\!22$	41.79	3 (17%)	23,17,30	24.63	10 (43%)
4	MBO	А	1205[A]	1,7	7,10,10	3.18	4 (57%)	11,13,13	2.15	3 (27%)
6	GOL	А	1004	-	$5,\!5,\!5$	1.30	0	$5,\!5,\!5$	2.06	2 (40%)
6	GOL	А	1003[A]	-	$5,\!5,\!5$	1.17	1 (20%)	$5,\!5,\!5$	2.61	2 (40%)
4	MBO	А	1205[B]	1	7,10,10	2.50	4 (57%)	11,13,13	1.90	4 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	А	1003[B]	-	-	2/4/4/4	-
5	SUA	А	1002[A]	3	-	7/14/14/16	0/1/1/2
5	SUA	А	1002[B]	-	-	6/14/8/16	0/1/1/2
4	MBO	А	1205[A]	1,7	-	0/4/4/4	0/1/1/1
6	GOL	А	1004	-	-	0/4/4/4	-
6	GOL	А	1003[A]	-	-	1/4/4/4	-
4	MBO	А	1205[B]	1	-	0/4/4/4	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	А	1002[B]	SUA	O3-C7	167.58	3.37	1.33
5	А	1002[B]	SUA	C7-S2	39.34	2.36	1.66
5	А	1002[A]	SUA	O3-C7	10.87	1.46	1.33
5	А	1002[A]	SUA	C7-N2	-9.93	1.26	1.34
5	А	1002[B]	SUA	C7-N2	6.70	1.40	1.34
4	А	1205[A]	MBO	CE4-CZ	5.06	1.60	1.49
4	А	1205[A]	MBO	OZ2-CZ	-4.82	1.15	1.30
4	А	1205[B]	MBO	OZ1-CZ	3.90	1.34	1.22
5	А	1002[A]	SUA	C7-S2	-3.68	1.59	1.66
4	А	1205[B]	MBO	CE6-CE5	3.64	1.45	1.38
4	А	1205[A]	MBO	OZ1-CZ	3.04	1.31	1.22
4	А	1205[A]	MBO	CE3-CE2	2.84	1.43	1.38
4	А	1205[B]	MBO	CE3-CE4	2.73	1.44	1.39
4	А	1205[B]	MBO	CE4-CZ	2.38	1.54	1.49
6	А	1003[A]	GOL	01-C1	2.28	1.52	1.42

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	1002[B]	SUA	O3-C7-S2	-109.92	49.98	125.10
5	А	1002[B]	SUA	O3-C7-N2	-33.22	42.46	111.94
5	А	1002[B]	SUA	S2-C7-N2	-24.23	69.48	123.02
5	А	1002[A]	SUA	C8-O3-C7	20.62	135.85	119.11
5	А	1002[A]	SUA	O3-C7-S2	-13.01	116.21	125.10
5	А	1002[A]	SUA	O3-C7-N2	11.85	136.72	111.94
5	А	1002[B]	SUA	C8-O3-C7	11.48	128.43	119.11
5	А	1002[A]	SUA	S2-C7-N2	-7.72	105.97	123.02
6	А	1003[A]	GOL	O2-C2-C1	4.70	129.82	109.12
4	А	1205[A]	MBO	CE6-CE5-CE4	-4.02	116.10	120.78
5	А	1002[A]	SUA	C6-N2-C7	-3.76	123.28	130.00
4	А	1205[A]	MBO	OZ2-CZ-OZ1	3.54	131.21	123.35

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	1205[A]	MBO	CE5-CE6-CE1	3.41	128.20	121.56
5	А	1002[B]	SUA	C6-N2-C7	-3.03	124.59	130.00
6	А	1004	GOL	C3-C2-C1	-3.01	99.99	111.70
6	А	1004	GOL	O2-C2-C3	3.01	122.38	109.12
4	А	1205[B]	MBO	CE2-CE3-CE4	-2.92	117.38	120.78
6	А	1003[A]	GOL	O1-C1-C2	2.91	124.17	110.20
4	А	1205[B]	MBO	CE5-CE6-CE1	2.54	126.52	121.56
4	А	1205[B]	MBO	OZ1-CZ-CE4	2.36	127.75	121.45
5	А	1002[A]	SUA	O3-C8-C9	-2.25	100.16	108.94
5	А	1002[B]	SUA	O3-C8-C9	-2.25	100.16	108.94
5	А	1002[A]	SUA	C1-C2-C3	2.24	121.77	119.45
5	А	1002[B]	SUA	C1-C2-C3	2.24	121.77	119.45
5	А	1002[A]	SUA	O2-S1-N1	2.15	110.56	107.36
5	А	1002[B]	SUA	O2-S1-N1	2.15	110.56	107.36
5	А	1002[A]	SUA	O1-S1-N1	2.06	110.42	107.36
5	А	1002[B]	SUA	O1-S1-N1	2.06	110.42	107.36
4	А	1205[B]	MBO	CE3-CE4-CZ	-2.03	116.40	120.39
6	А	1003[B]	GOL	C3-C2-C1	2.03	119.58	111.70
5	А	1002[A]	SUA	C10-C9-C8	-2.02	100.08	112.32
5	А	1002[B]	SUA	C10-C9-C8	-2.02	100.08	112.32

Continued from previous page...

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	А	1002[A]	SUA	N2-C7-O3-C8
5	А	1002[A]	SUA	S2-C7-O3-C8
5	А	1002[A]	SUA	C9-C8-O3-C7
5	А	1002[B]	SUA	O3-C7-N2-C6
5	А	1002[B]	SUA	S2-C7-O3-C8
6	А	1003[A]	GOL	C1-C2-C3-O3
5	А	1002[B]	SUA	C1-C6-N2-C7
5	А	1002[B]	SUA	C5-C6-N2-C7
6	А	1003[B]	GOL	C1-C2-C3-O3
6	А	1003[B]	GOL	O2-C2-C3-O3
5	А	1002[A]	SUA	O3-C7-N2-C6
5	А	1002[A]	SUA	S2-C7-N2-C6
5	А	1002[A]	SUA	C4-C3-S1-O2
5	А	1002[B]	SUA	C4-C3-S1-O2
5	А	1002[A]	SUA	C2-C3-S1-O2
5	А	1002[B]	SUA	C2-C3-S1-O2




There are no ring outliers.


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	1003[B]	GOL	1	0
5	А	1002[A]	SUA	2	0
6	А	1004	GOL	3	0
4	А	1205[B]	MBO	1	0

4 monomers are involved in 7 short contacts:


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	А	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	76:LYS	С	77:ALA	N	1.20



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

