

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2023 – 02:52 AM EDT

PDB ID : 7LUT

Title: The internal aldimine form of the wild-type Salmonella typhimurium Trypto-

phan Synthase in complex with cesium ion at the metal coordination site and the product L-tryptophan at the enzyme beta-site at 1.60 Angstrom resolution

Authors: Hilario, E.; Dunn, M.F.; Mueller, L.J.

Deposited on : 2021-02-23

Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	<u> </u>					
1	A	268	91%	5%	-				
2	В	397	89%	10%	-				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Chain Res C		Geometry	Clashes	Electron density
8	PEG	В	412	_	_	X	-



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 5819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Tryptophan synthase alpha chain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	256	Total 1939	C 1234	N 335	O 362	S 8	0	2	0

• Molecule 2 is a protein called Tryptophan synthase beta chain.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
2	В	394	Total	С	N	О	S	0	Q	0
	Ъ	394	3037	1905	533	578	21	0	8	

• Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O S 4 2 1 1	0	0
3	A	1	Total C O S 4 2 1 1	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C O S 4 2 1 1	0	0
3	В	1	Total C O S 4 2 1 1	0	0
3	В	1	Total C O S 4 2 1 1	0	0
3	В	1	Total C O S 4 2 1 1	0	0
3	В	1	Total C O S 4 2 1 1	0	0
3	В	1	Total C O S 4 2 1 1	0	0

 \bullet Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	В	1	Total C O 4 2 2	0	0
4	В	1	Total C O 4 2 2	0	0

• Molecule 5 is CESIUM ION (three-letter code: CS) (formula: Cs) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cs 1 1	0	0
5	В	5	Total Cs 6 6	0	1

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	В	4	Total Cl 4 4	0	0


• Molecule 7 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
7	В	1	Total	С	N	О	Р	0	0
'		1	15	8	1	5	1		


• Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	В	1	Total C 7 4	O 3	0	0

• Molecule 9 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf		
9	В	1	Total 15	C 11	N 2	O 2	0	0

• Molecule 10 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	277	Total O 288 288	0	11
10	В	444	Total O 462 462	0	18



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan synthase alpha chain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	182.41Å 61.09Å 67.06Å	Donositor
a, b, c, α , β , γ	90.00° 94.86° 90.00°	Depositor
Resolution (Å)	32.28 - 1.60	Depositor
Resolution (A)	32.26 - 1.60	EDS
% Data completeness	93.8 (32.28-1.60)	Depositor
(in resolution range)	93.8 (32.26-1.60)	EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	2.22 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D	0.187 , 0.217	Depositor
R, R_{free}	0.188 , 0.219	DCC
R_{free} test set	4520 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	13.4	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 52.8	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5819	wwPDB-VP
Average B, all atoms $(Å^2)$	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, CL, PEG, EDO, CS, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.41	0/1979	0.75	0/2691	
2	В	0.53	1/3095 (0.0%)	0.83	1/4180 (0.0%)	
All	All	0.49	1/5074 (0.0%)	0.80	1/6871 (0.0%)	

All (1) bond length outliers are listed below:

N	/Iol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
	2	В	11	GLU	CD-OE1	6.75	1.33	1.25

All (1) bond angle outliers are listed below:

Mo	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	В	342	HIS	CA-CB-CG	-5.15	104.84	113.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1939	0	1924	13	0
2	В	3037	0	2995	42	0
3	A	8	0	12	2	0
3	В	24	0	36	1	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	5	3	0
4	В	8	0	12	1	0
5	A	1	0	0	0	0
5	В	6	0	0	0	0
6	A	1	0	0	0	0
6	В	4	0	0	0	0
7	В	15	0	6	1	0
8	В	7	0	10	4	0
9	В	15	0	9	1	0
10	A	288	0	0	2	0
10	В	462	0	0	10	0
All	All	5819	0	5009	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
2:B:223:LEU:HD23	10:B:657:HOH:O	1.77	0.82
1:A:107:PHE:CE2	4:A:303:EDO:H21	2.19	0.78
1:A:1:MET:HG2	10:A:521[B]:HOH:O	1.88	0.74
1:A:117:ARG:HG2	3:A:302:DMS:C2	2.21	0.71
2:B:217:LEU:HG	10:B:657:HOH:O	1.92	0.69
2:B:216:ILE:HG23	10:B:657:HOH:O	1.95	0.66
2:B:300:ILE:HD11	2:B:390:ILE:CD1	2.27	0.65
2:B:216:ILE:CG2	10:B:657:HOH:O	2.45	0.65
2:B:44[B]:GLN:HG3	10:B:711:HOH:O	1.97	0.64
2:B:87:LYS:HD2	2:B:114:GLN:HG3	1.78	0.64
1:A:107:PHE:CE2	4:A:303:EDO:C2	2.84	0.59
1:A:117:ARG:HG2	3:A:302:DMS:H22	1.85	0.58
2:B:216:ILE:HG21	2:B:224:PRO:HD3	1.85	0.58
2:B:10:GLY:H	8:B:412:PEG:H32	1.70	0.57
1:A:107:PHE:CZ	4:A:303:EDO:H21	2.40	0.57
1:A:64:ILE:CD1	1:A:238:VAL:HG21	2.35	0.56
2:B:79:ASP:HB2	2:B:379:ARG:HB3	1.90	0.54
2:B:300:ILE:HD11	2:B:390:ILE:HD13	1.90	0.53
2:B:134[B]:MET:CE	2:B:139:VAL:HG22	2.39	0.52
3:B:404:DMS:H22	10:B:600:HOH:O	2.10	0.52
2:B:360[B]:LYS:HG3	10:B:679:HOH:O	2.10	0.52
2:B:10:GLY:H	8:B:412:PEG:C3	2.22	0.52

Continued on next page...



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
2:B:337:LYS:NZ	10:B:504:HOH:O	2.43	0.52
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.45	0.51
2:B:134[B]:MET:CE	2:B:156:VAL:CG1	2.90	0.50
2:B:65:ILE:HD11	2:B:338:THR:HG22	1.93	0.50
2:B:109:GLU:O	2:B:115:HIS:HD2	1.95	0.49
2:B:134[B]:MET:HE1	2:B:156:VAL:HG11	1.95	0.49
2:B:134[B]:MET:CE	2:B:156:VAL:HG11	2.44	0.48
1:A:100:LEU:C	1:A:100:LEU:HD13	2.35	0.47
10:A:412[A]:HOH:O	4:B:411:EDO:H22	2.14	0.47
2:B:134[B]:MET:HE1	2:B:156:VAL:CG1	2.45	0.47
2:B:134[B]:MET:HE2	2:B:134[B]:MET:HB3	1.64	0.47
2:B:306:PHE:CD1	2:B:307:PRO:HD2	2.49	0.47
2:B:134[B]:MET:HE3	2:B:139:VAL:HG22	1.97	0.46
2:B:173:ALA:HB1	2:B:186:TYR:CE1	2.50	0.46
2:B:337:LYS:HE3	2:B:391:LEU:HD11	1.98	0.46
2:B:10:GLY:N	8:B:412:PEG:H32	2.31	0.44
2:B:87:LYS:NZ	7:B:401:PLP:O3	2.51	0.44
1:A:64:ILE:HD12	1:A:238:VAL:HG21	2.00	0.43
2:B:300:ILE:CD1	2:B:390:ILE:HD13	2.49	0.43
1:A:36:ILE:O	1:A:40:LEU:HG	2.19	0.43
2:B:341:ARG:NH1	2:B:342:HIS:NE2	2.67	0.42
1:A:104:ASN:HB2	2:B:278:ILE:O	2.19	0.42
2:B:219:LYS:NZ	10:B:513:HOH:O	2.51	0.42
2:B:301:SER:OG	2:B:350:GLU:HG3	2.20	0.42
1:A:57:PRO:HA	1:A:102:TYR:CZ	2.54	0.42
2:B:6:ASN:HD21	8:B:412:PEG:C1	2.33	0.42
2:B:87:LYS:HE2	9:B:420:TRP:OXT	2.20	0.41
2:B:65:ILE:CD1	2:B:338:THR:HG22	2.51	0.41
2:B:129:LYS:HE3	2:B:129:LYS:HB3	1.72	0.41
2:B:132:ILE:HD13	2:B:149:MET:SD	2.61	0.41
2:B:279:TYR:CE2	2:B:280:PHE:HB2	2.56	0.41
2:B:86:HIS:CE1	2:B:236:ASN:HB3	2.56	0.41
2:B:114:GLN:HG2	10:B:548:HOH:O	2.20	0.41
1:A:191:LEU:HA	1:A:192:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Perce	\mathbf{ntiles}
1	A	254/268~(95%)	250 (98%)	4 (2%)	0	100	100
2	В	400/397 (101%)	393 (98%)	7 (2%)	0	100	100
All	All	654/665 (98%)	643 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	199/208 (96%)	199 (100%)	0	100	100	
2	В	314/311 (101%)	314 (100%)	0	100	100	
All	All	513/519 (99%)	513 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 12 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Clasica	Dan	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DMS	A	302	-	3,3,3	0.23	0	3,3,3	0.24	0
3	DMS	В	405	-	3,3,3	0.25	0	3,3,3	0.16	0
8	PEG	В	412	-	6,6,6	0.32	0	5,5,5	0.32	0
3	DMS	В	404	-	3,3,3	0.27	0	3,3,3	0.21	0
3	DMS	В	407	-	3,3,3	0.25	0	3,3,3	0.17	0
9	TRP	В	420	-	14,16,16	0.81	0	16,22,22	1.23	2 (12%)
3	DMS	В	402	-	3,3,3	0.68	0	3,3,3	0.22	0
7	PLP	В	401	2	15,15,16	0.75	0	20,22,23	1.37	3 (15%)
3	DMS	В	406	-	3,3,3	0.28	0	3,3,3	0.04	0
3	DMS	В	403	-	3,3,3	0.22	0	3,3,3	0.18	0
4	EDO	В	410	-	3,3,3	0.13	0	2,2,2	0.28	0
3	DMS	A	301	-	3,3,3	0.25	0	3,3,3	0.17	0
4	EDO	В	411	-	3,3,3	0.11	0	2,2,2	0.11	0
4	EDO	A	303	-	3,3,3	1.20	1 (33%)	2,2,2	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PEG	В	412	-	-	3/4/4/4	-
9	TRP	В	420	-	-	0/7/8/8	0/2/2/2
7	PLP	В	401	2	-	0/6/6/8	0/1/1/1
4	EDO	В	410	-	-	1/1/1/1	-
4	EDO	В	411	-	-	1/1/1/1	-
4	EDO	A	303	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
4	A	303	EDO	O2-C2	-2.02	1.31	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathbf{Ideal}(^o)$
7	В	401	PLP	O4P-C5A-C5	3.63	116.28	109.35
9	В	420	TRP	CH2-CZ2-CE2	-2.84	115.99	120.08
7	В	401	PLP	C4A-C4-C5	2.40	123.41	120.94
9	В	420	TRP	CZ3-CE3-CD2	-2.25	117.77	120.89
7	В	401	PLP	O3P-P-O4P	-2.08	101.21	106.73

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	В	412	PEG	O2-C3-C4-O4
8	В	412	PEG	O1-C1-C2-O2
4	В	410	EDO	O1-C1-C2-O2
4	A	303	EDO	O1-C1-C2-O2
8	В	412	PEG	C4-C3-O2-C2
4	В	411	EDO	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	DMS	2	0
8	В	412	PEG	4	0
3	В	404	DMS	1	0
9	В	420	TRP	1	0
7	В	401	PLP	1	0


Continued on next page...




Continued from previous page...

Mo	Chain	Res	Type	Clashes	Symm-Clashes
4	В	411	EDO	1	0
4	A	303	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ > #RSRZ > 2		$OWAB(A^2)$	Q<0.9
1	A	256/268~(95%)	-0.07	14 (5%) 25 22	10, 18, 32, 42	0
2	В	394/397~(99%)	-0.11	10 (2%) 57 55	7, 13, 27, 57	0
All	All	650/665~(97%)	-0.09	24 (3%) 41 39	7, 15, 30, 57	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	393	ALA	4.1
1	A	194[A]	HIS	3.5
2	В	385	PHE	3.4
1	A	191	LEU	3.3
1	A	268	ALA	3.1
2	В	160	HIS	2.9
2	В	2	THR	2.9
1	A	13	ASP	2.9
1	A	212	PHE	2.8
1	A	250	GLN	2.8
1	A	195	HIS	2.5
2	В	159	VAL	2.5
1	A	192	PRO	2.5
2	В	394	ARG	2.5
1	A	246	ALA	2.5
1	A	247	SER	2.4
2	В	163	SER	2.4
2	В	391	LEU	2.3
2	В	141	ARG	2.3
1	A	193	LEU	2.3
1	A	157	ASN	2.2
1	A	178	SER	2.2
1	A	15	ARG	2.1
2	В	392	LYS	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q<0.9
6	CL	В	417	1/1	0.73	0.07	60,60,60,60	0
3	DMS	В	403	4/4	0.77	0.18	50,55,58,60	0
4	EDO	В	411	4/4	0.78	0.23	41,43,43,45	0
8	PEG	В	412	7/7	0.78	0.24	27,32,33,33	0
3	DMS	В	406	4/4	0.79	0.21	61,64,65,70	0
6	CL	В	416	1/1	0.84	0.07	62,62,62,62	0
3	DMS	В	405	4/4	0.86	0.18	51,54,55,59	0
6	CL	В	419	1/1	0.90	0.14	51,51,51,51	0
4	EDO	В	410	4/4	0.90	0.12	28,29,29,30	0
3	DMS	A	302	4/4	0.92	0.17	30,34,36,39	0
3	DMS	В	404	4/4	0.92	0.21	39,40,40,41	0
3	DMS	В	407	4/4	0.93	0.21	49,51,52,55	0
4	EDO	A	303	4/4	0.93	0.18	7,12,13,24	0
9	TRP	В	420	15/15	0.93	0.10	17,17,19,19	0
3	DMS	В	402	4/4	0.94	0.15	22,22,23,25	0
6	CL	A	305	1/1	0.96	0.12	41,41,41,41	0
3	DMS	A	301	4/4	0.96	0.11	39,39,42,44	0
5	CS	В	413	1/1	0.97	0.08	40,40,40,40	1
5	CS	В	415	1/1	0.98	0.04	21,21,21,21	1
5	CS	A	304	1/1	0.98	0.04	22,22,22,22	1
6	CL	В	418	1/1	0.98	0.05	21,21,21,21	0
5	CS	В	414	1/1	0.99	0.03	15,15,15,15	1
7	PLP	В	401	15/16	0.99	0.08	8,9,10,11	0
5	CS	В	409[B]	1/1	1.00	0.06	26,26,26,26	1
5	CS	В	408	1/1	1.00	0.03	14,14,14,14	0
5	CS	В	409[A]	1/1	1.00	0.06	13,13,13,13	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















Electron density around CS A 304:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around CS B 414:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



















Electron density around CS B 408: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

