



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2023 – 12:49 AM EDT

PDB ID : 7LXS  
Title : Structural and Biochemical Insight into Assembly of Molecular Motors Involved in Viral DNA Packaging  
Authors : Ortega, M.E.  
Deposited on : 2021-03-04  
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	55	64% (green), 29% (yellow), 7% (orange)
1	B	55	67% (green), 27% (yellow), 5% (orange)
1	C	55	62% (green), 33% (yellow), 5% (orange)
1	D	55	65% (green), 27% (yellow), 7% (orange)
1	E	55	56% (green), 38% (yellow), 5% (orange)
1	F	55	64% (green), 31% (yellow), 5% (orange)
1	G	55	67% (green), 27% (yellow), 5% (orange)

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	H	55	 64% 31% 5%

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Terminase, small subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	55	412	259	72	79	2	0	0	0
1	B	55	412	259	72	79	2	0	0	0
1	C	55	412	259	72	79	2	0	0	0
1	D	55	412	259	72	79	2	0	0	0
1	E	55	412	259	72	79	2	0	0	0
1	F	55	412	259	72	79	2	0	0	0
1	G	55	412	259	72	79	2	0	0	0
1	H	55	412	259	72	79	2	0	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Terminase, small subunit

Chain A:  64% 29% 7%




- Molecule 1: Terminase, small subunit

Chain B:  67% 27% 5%



- Molecule 1: Terminase, small subunit

Chain C:  62% 33% 5%




- Molecule 1: Terminase, small subunit

Chain D:  65% 27% 7%




- Molecule 1: Terminase, small subunit

Chain E:  56% 38% 5%




- Molecule 1: Terminase, small subunit


Chain F:  64% 31% 5%



- Molecule 1: Terminase, small subunit



- Molecule 1: Terminase, small subunit



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	24.75Å 50.76Å 68.53Å 90.00° 88.36° 90.11°	Depositor
Resolution (Å)	24.74 – 2.21 24.74 – 2.21	Depositor EDS
% Data completeness (in resolution range)	94.6 (24.74-2.21) 87.2 (24.74-2.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	28.53 (at 2.22Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.140 , 0.142 0.134 , 0.137	Depositor DCC
$R_{free}$ test set	1606 reflections (10.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.9	Xtrriage
Anisotropy	0.405	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 0.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.187 for h,-k,-l 0.479 for -h,k,-l 0.189 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.78	EDS
Total number of atoms	3296	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 59.96 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6342e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/419	0.82	0/566
1	B	0.79	0/419	0.80	0/566
1	C	0.81	0/419	0.87	0/566
1	D	0.85	0/419	0.82	0/566
1	E	0.80	0/419	0.78	0/566
1	F	0.81	0/419	0.77	0/566
1	G	0.82	0/419	0.81	0/566
1	H	0.80	0/419	0.78	0/566
All	All	0.81	0/3352	0.81	0/4528

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	412	0	406	40	0
1	B	412	0	406	21	0
1	C	412	0	406	46	0
1	D	412	0	406	21	0
1	E	412	0	406	38	0
1	F	412	0	406	19	0
1	G	412	0	406	23	0
1	H	412	0	406	28	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3296	0	3248	198	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 198 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ALA:HB3	1:C:55:ALA:CB	1.59	1.32
1:A:55:ALA:CB	1:C:55:ALA:HB3	1.62	1.28
1:G:54:ASP:HB3	1:H:1:MET:HE2	1.26	1.17
1:E:55:ALA:CB	1:H:55:ALA:HB3	1.76	1.15
1:A:55:ALA:CB	1:C:55:ALA:CB	2.24	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	53/55 (96%)	48 (91%)	5 (9%)	0	100	100
1	B	53/55 (96%)	49 (92%)	4 (8%)	0	100	100
1	C	53/55 (96%)	49 (92%)	4 (8%)	0	100	100
1	D	53/55 (96%)	48 (91%)	4 (8%)	1 (2%)	8	4
1	E	53/55 (96%)	49 (92%)	4 (8%)	0	100	100
1	F	53/55 (96%)	48 (91%)	5 (9%)	0	100	100
1	G	53/55 (96%)	48 (91%)	5 (9%)	0	100	100
1	H	53/55 (96%)	48 (91%)	5 (9%)	0	100	100
All	All	424/440 (96%)	387 (91%)	36 (8%)	1 (0%)	47	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	37	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	41/41 (100%)	37 (90%)	4 (10%)	8	6
1	B	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	C	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	D	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	E	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	F	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	G	41/41 (100%)	38 (93%)	3 (7%)	14	14
1	H	41/41 (100%)	38 (93%)	3 (7%)	14	14
All	All	328/328 (100%)	303 (92%)	25 (8%)	13	12

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	10	GLU
1	F	7	LYS
1	H	35	LYS
1	F	6	LYS
1	F	35	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	2	ASN
1	H	2	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.