

# Full wwPDB NMR Structure Validation Report (i)

### Jun 6, 2023 – 01:32 AM EDT

PDB ID : 2M7X BMRB ID : 19216

Title : Structural and Functional Analysis of Transmembrane Segment IV of the Salt

Tolerance Protein Sod2

Authors: Ullah, A.; Kemp, G.; Lee, B.; Alves, C.; Young, H.; Sykes, B.D.; Fliegel, L.

Deposited on : 2013-05-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 50%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mo	l Chain	Length			Quality of chain	
1	A	38	21%	5%	53%	21%



# 2 Ensemble composition and analysis (i)

This entry contains 25 models. Model 16 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core   Residue range (total)   Backbone RMSD (Å)   Medoid m				
1	A:132-A:141 (10)	0.81	16	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	5, 6, 7, 8, 11, 12, 14, 19, 21, 24
2	1, 2, 3, 4, 13, 16, 20, 22, 25
3	9, 10, 23
4	15, 17
Single-model clusters	18



# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 446 atoms, of which 231 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Na(+)/H(+) antiporter.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	20	Total	С	Н	N	О	S	0
	A	30	446	143	231	32	39	1	U

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	GLY	-	expression tag	UNP P36606
A	121	SER	-	expression tag	UNP P36606
A	122	LYS	-	expression tag	UNP P36606
A	123	LYS	-	expression tag	UNP P36606
A	124	LYS	-	expression tag	UNP P36606
A	155	LYS	-	expression tag	UNP P36606
A	156	LYS	-	expression tag	UNP P36606
A	157	LYS	-	expression tag	UNP P36606



# 4 Residue-property plots (i)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%

#### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

• Molecule 1: Na(+)/H(+) antiporter



#### 4.2.2 Score per residue for model 2

• Molecule 1: Na(+)/H(+) antiporter





#### 4.2.3 Score per residue for model 3

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 24% . 53% 21%



#### 4.2.4 Score per residue for model 4

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%

GLY CLYS LYS LYS L126 P126 P126 P127 P128 P128 P129 P131 P142 P142 P142 P142 P142 P143 P145 P146 P150 P150 P160 P170 P17

#### 4.2.5 Score per residue for model 5

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 16% 8% • 53% 21%

#### 4.2.6 Score per residue for model 6

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%

GLY GLY LYS LYS LYS LYS LYS 1129 Q128 Q128 Q128 T142 T142 T142 T142 T144 T142 T144 T142 T144 T148 T

#### 4.2.7 Score per residue for model 7

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%





#### 4.2.8 Score per residue for model 8

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 13% 11% • 53% 21%



### 4.2.9 Score per residue for model 9

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 16% 11% 53% 21%

#### GLY CLYS CLYS CLYS CLYS F126 P129 P129 P131 F131 F

#### 4.2.10 Score per residue for model 10

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 24% • 53% 21%



#### 4.2.11 Score per residue for model 11

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 16% 11% 53% 21%



#### 4.2.12 Score per residue for model 12

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%





#### 4.2.13 Score per residue for model 13

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%

0 0 1 V S S R R 1 V S S R R 1 V S S R R 1 V S S R R 1 V S S R R 1 V S S R R 1 V S R 1

### 4.2.14 Score per residue for model 14

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%

#### 4.2.15 Score per residue for model 15

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%

#### 4.2.16 Score per residue for model 16 (medoid)

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 24% • 53% 21%

#### 4.2.17 Score per residue for model 17

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%



#### 4.2.18 Score per residue for model 18

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 13% 11% • 53% 21%



### 4.2.19 Score per residue for model 19

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 13% 13% 53% 21%



#### 4.2.20 Score per residue for model 20

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%



#### 4.2.21 Score per residue for model 21

• Molecule 1: Na(+)/H(+) antiporter


Chain A: 16% 11% 53% 21%



#### 4.2.22 Score per residue for model 22

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 24% • 53% 21%





### 4.2.23 Score per residue for model 23

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 21% 5% 53% 21%

### 4.2.24 Score per residue for model 24

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%

#### 4.2.25 Score per residue for model 25

• Molecule 1: Na(+)/H(+) antiporter

Chain A: 18% 8% 53% 21%



# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: simulated annealing.

Of the 50 calculated structures, 25 were deposited, based on the following criterion: *structures* with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	210
Number of shifts mapped to atoms	176
Number of unparsed shifts	0
Number of shifts with mapping errors	34
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	50%



# 6 Model quality (i)

### 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	65	76	76	2±2
All	All	1625	1900	1900	42

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom 1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:132:LEU:HD13	1:A:133:GLY:N	0.91	1.80	18	1
1:A:132:LEU:O	1:A:132:LEU:HD22	0.74	1.81	18	1
1:A:138:ALA:O	1:A:141:ILE:HG22	0.71	1.85	20	3
1:A:132:LEU:HD22	1:A:132:LEU:C	0.63	2.15	18	1
1:A:134:SER:O	1:A:137:ILE:HG22	0.62	1.94	18	4
1:A:135:LEU:H	1:A:135:LEU:HD23	0.57	1.58	5	1
1:A:132:LEU:HD13	1:A:132:LEU:C	0.57	2.19	18	1
1:A:137:ILE:HG23	1:A:138:ALA:N	0.56	2.15	9	1
1:A:141:ILE:HD12	1:A:141:ILE:N	0.52	2.20	4	4
1:A:132:LEU:O	1:A:134:SER:N	0.52	2.43	11	5
1:A:140:CYS:SG	1:A:141:ILE:N	0.51	2.83	6	1
1:A:135:LEU:H	1:A:135:LEU:CD2	0.50	2.17	5	1
1:A:134:SER:OG	1:A:135:LEU:N	0.50	2.44	19	1
1:A:132:LEU:C	1:A:134:SER:N	0.46	2.68	5	4

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:141:ILE:N	1:A:141:ILE:CD1	0.46	2.78	4	4
1:A:137:ILE:CG2	1:A:138:ALA:N	0.46	2.79	9	1
1:A:140:CYS:O	1:A:140:CYS:SG	0.45	2.74	2	1
1:A:132:LEU:O	1:A:135:LEU:N	0.45	2.49	18	1
1:A:138:ALA:O	1:A:140:CYS:N	0.43	2.50	13	2
1:A:138:ALA:C	1:A:140:CYS:N	0.43	2.71	13	2
1:A:132:LEU:C	1:A:132:LEU:CD2	0.42	2.84	18	1
1:A:132:LEU:O	1:A:132:LEU:HD23	0.40	2.17	19	1

### 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed Outliers Percent		entiles		
1	A	10/38 (26%)	9±1 (92±9%)	1±1 (6±7%)	0±0 (2±4%)	13	57
All	All	250/950~(26%)	231 (92%)	15 (6%)	4 (2%)	13	57

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	133	GLY	4

### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Perce	ntiles
1	A	7/32 (22%)	6±1 (83±11%)	1±1 (17±11%)	5	41
All	All	175/800~(22%)	146 (83%)	29 (17%)	5	41

All 6 unique residues with a non-rotameric sidechain are listed below. They are sorted by the



frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	140	CYS	10
1	A	134	SER	9
1	A	136	LEU	5
1	A	135	LEU	2
1	A	132	LEU	2
1	A	137	ILE	1

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.6 Ligand geometry (i)

There are no ligands in this entry.

## 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 50% for the well-defined parts and 44% for the entire structure.

### 7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	210
Number of shifts mapped to atoms	176
Number of unparsed shifts	0
Number of shifts with mapping errors	34
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. All 34 occurrences are reported below.

T: / ID	aı ·	Ъ	TD.	<b>A</b> .		Shift Data		
List ID	Chain	Res	Type	$egin{aligned}  ext{Type} &  ext{Atom} \end{aligned}$	Value	Uncertainty	Ambiguity	
1	A	120	GLY	HA2	3.785		2	
1	A	120	GLY	HA3	3.913		2	
1	A	121	SER	N	115.757		1	
1	A	121	SER	Н	8.85		1	
1	A	121	SER	HA	4.337		1	
1	A	122	LYS	N	123.238		1	
1	A	122	LYS	Н	8.753		1	
1	A	122	LYS	HA	4.034		1	
1	A	122	LYS	HB2	1.808	•	2	
1	A	123	LYS	N	117.981	•	1	
1	A	123	LYS	Н	7.91		1	
1	A	123	LYS	HA	3.862		1	
1	A	123	LYS	HB2	1.744		2	
1	A	123	LYS	HG2	1.441		2	

Continued on next page...



Continued from previous page...

	Chain			Atom		ι	
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	124	LYS	N	115.479	•	1
1	A	124	LYS	Н	7.445	•	1
1	A	124	LYS	HA	4.077	•	1
1	A	124	LYS	HB2	1.765	•	2
1	A	124	LYS	HG2	1.413	•	2
1	A	155	LYS	N	118.434	•	1
1	A	155	LYS	Н	7.637	•	1
1	A	155	LYS	HA	4.184	•	1
1	A	155	LYS	HB2	1.823	•	2
1	A	155	LYS	HG2	1.449	•	2
1	A	156	LYS	N	119.263	•	1
1	A	156	LYS	Н	8.03		1
1	A	156	LYS	HA	4.18	•	1
1	A	156	LYS	HB2	1.808	•	2
1	A	156	LYS	HG2	1.443	•	2
1	A	157	LYS	N	118.155	•	1
1	A	157	LYS	Н	7.902	•	1
1	A	157	LYS	HA	4.289	•	1
1	A	157	LYS	HB2	1.747	•	2
1	A	157	LYS	HG2	1.375	•	2

### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	0		None (insufficient data)
$^{13}C_{\beta}$	0		None (insufficient data)
<sup>13</sup> C′	0		None (insufficient data)
$^{15}N$	35	$1.55 \pm 0.55$	Should be applied

### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 50%, i.e. 63 atoms were assigned a chemical shift out of a possible 127. 0 out of 3 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}{ m C}$	$^{15}{ m N}$
Backbone	31/52~(60%)	21/22~(95%)	0/20~(0%)	10/10 (100%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}\mathbf{N}$
Sidechain	32/75 (43%)	32/52~(62%)	$0/23 \ (0\%)$	0/0 (%)
Overall	63/127 (50%)	53/74 (72%)	0/43 (0%)	10/10 (100%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 44%, i.e. 175 atoms were assigned a chemical shift out of a possible 396. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	87/149 (58%)	59/61 (97%)	0/60 (0%)	28/28 (100%)
Sidechain	88/227 (39%)	88/154 (57%)	0/71 (0%)	0/2 (0%)
Aromatic	0/20~(0%)	0/10 (0%)	0/10 (0%)	0/0 (%)
Overall	175/396 (44%)	147/225 (65%)	0/141 (0%)	28/30 (93%)

#### 7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	129	ILE	HG21	2.17	-0.56 - 2.11	5.2
1	A	129	ILE	HG22	2.17	-0.56 - 2.11	5.2
1	A	129	ILE	HG23	2.17	-0.56 - 2.11	5.2

### 7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







# 8 NMR restraints analysis (i)

### 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	0
Intra-residue ( i-j =0)	0
Sequential ( i-j =1)	0
Medium range ( $ i-j >1$ and $ i-j <5$ )	0
Long range ( i-j ≥5)	0
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	41
Number of unmapped restraints	0
Number of restraints per residue	0
Number of long range restraints per residue <sup>1</sup>	0

<sup>&</sup>lt;sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

### 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.


### 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation. There are no distance restraints

### 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^{\circ}$  are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	$\mathbf{Max} (^{\circ})$
1.0-10.0 (Small)	0.0	1.4





Continued from previous page...

Bins (°)	Average number of violations per model	Max (°)
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



# 9 Distance violation analysis (i)

No distance restraints data found



# 10 Dihedral-angle violation analysis (i)


### 10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle tree	Count % <sup>1</sup>		${f Violated}^3$			Consistently Violated <sup>4</sup>		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	<b>%</b> ¹
PHI	21	51.2	0	0.0	0.0	0	0.0	0.0
PSI	20	48.8	1	5.0	2.4	0	0.0	0.0
Total	41	100.0	1	2.4	2.4	0	0.0	0.0

 $<sup>^1</sup>$  percentage calculated with respect to total number of dihedral-angle restraints,  $^2$  percentage calculated with respect to number of restraints in a particular dihedral-angle type,  $^3$  violated in at least one model,  $^4$  violated in all the models

### 10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




# 10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Madalin	Number of violations		<b>N</b> /L (0)	N/ (0)	CD (0)	N ( - 1' (0)	
Model ID	PHI	PSI	Total	$\mathbf{Mean}  (^{\circ})$	$\mathbf{Max} \ (^{\circ})$	$\mathbf{SD}$ (°)	$\mid$ Median (°) $\mid$
1	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0.0	0.0	0.0	0.0
11	0	0	0	0.0	0.0	0.0	0.0
12	0	0	0	0.0	0.0	0.0	0.0
13	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0.0	0.0	0.0	0.0
18	0	1	1	1.4	1.4	0.0	1.4
19	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0.0	0.0	0.0	0.0
21	0	0	0	0.0	0.0	0.0	0.0
22	0	0	0	0.0	0.0	0.0	0.0
23	0	0	0	0.0	0.0	0.0	0.0
24	0	0	0	0.0	0.0	0.0	0.0
25	0	0	0	0.0	0.0	0.0	0.0



### 10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

# 10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Num	iber o	f violated restraints	Fraction of the ensemble			
PHI	PSI	Total	Count <sup>1</sup>	%		
0	1	1	1	4.0		
0	0	0	2	8.0		
0	0	0	3	12.0		
0	0	0	4	16.0		
0	0	0	5	20.0		
0	0	0	6	24.0		
0	0	0	7	28.0		
0	0	0	8	32.0		
0	0	0	9	36.0		
0	0	0	10	40.0		
0	0	0	11	44.0		

Continued on next page...




Continued from previous page...

Num	ıber o	f violated restraints	Fraction of the ensemble			
PHI	PSI	Total	Count <sup>1</sup>	%		
0	0	0	12	48.0		
0	0	0	13	52.0		
0	0	0	14	56.0		
0	0	0	15	60.0		
0	0	0	16	64.0		
0	0	0	17	68.0		
0	0	0	18	72.0		
0	0	0	19	76.0		
0	0	0	20	80.0		
0	0	0	21	84.0		
0	0	0	22	88.0		
0	0	0	23	92.0		
0	0	0	24	96.0		
0	0	0	25	100.0		

 $<sup>^{1}</sup>$  Number of models with violations

### 10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)





### 10.4 Most violated dihedral-angle restraints in the ensemble (i)

No violations found

### 10.5 All violated dihedral-angle restraints (i)

#### 10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

#### 10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,11)	1:A:132:LEU:N	1:A:132:LEU:CA	1:A:132:LEU:C	1:A:133:GLY:N	18	1.4

