Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2023 – 01:12 AM EDT

PDB ID	:	7M8S
Title	:	Crystal Structure of HLA-A*02:01 in complex with KLNDLCFTNV, an 10-mer epitope from SARS-CoV-2 Spike (S386-395)
Authors	:	Gras, S.; Nguyen, A.T.; Szeto, C.
Deposited on	:	2021-03-30
Resolution	:	2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	91% 8% •
1	D	276	8% 92% 7% •
2	B	100	2% 90% 9% •
2	E	100	2% 87% 12% •
3	C	10	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	10	80% 10% 10%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	2253	1408	410	426	9	0	0	0
1	D	274	2248	1404	412	423	9	0	1	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	848	539	145	160	4	0	1	0
2	E	99	829	528	140	158	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	expression tag	UNP P61769
E	0	MET	-	expression tag	UNP P61769

- Molecule 3 is a protein called Spike protein S1 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	10	81	51	13	16	1	0	0	0
3	F	10	81	51	13	16	1	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	113	113	113	0	0
5	B	46	46	46	0	0
5	C	9	9	9	0	0
5	D	84	84	84	0	0
5	E	35	35	35	0	0
5	F	7	7	7	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HLA class I histocompatibility antigen, A alpha chain

Chain A: 91% 8%



• Molecule 1: HLA class I histocompatibility antigen, A alpha chain

Chain D: 8% 92% 7%



• Molecule 2: Beta-2-microglobulin

Chain B: 2% 90% 9%



• Molecule 2: Beta-2-microglobulin

Chain E: 2% 87% 12%



• Molecule 3: Spike protein S1 peptide

Chain C: 100%

There are no outlier residues recorded for this chain.

• Molecule 3: Spike protein S1 peptide

Chain F: 80% 10% 10%


K1
L2
T8
N9
V10

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	64.60Å 86.69Å 163.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.08 – 2.35 46.08 – 2.35	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.08-2.35) 100.0 (46.08-2.35)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.94 (at 2.34Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.197 , 0.247 0.199 , 0.251	Depositor DCC
R_{free} test set	1956 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	41.2	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 33.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6640	wwPDB-VP
Average B, all atoms (Å^2)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.45	0/2319	0.63	0/3149
1	D	0.41	0/2313	0.62	0/3139
2	B	0.43	0/871	0.61	0/1176
2	E	0.40	0/852	0.61	0/1152
3	C	0.52	0/81	0.66	0/107
3	F	0.51	0/81	0.84	0/107
All	All	0.43	0/6517	0.62	0/8830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2253	0	2103	12	0
1	D	2248	0	2102	8	0
2	B	848	0	815	4	0
2	E	829	0	794	4	0
3	C	81	0	83	0	0
3	F	81	0	83	1	0
4	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	113	0	0	0	0
5	B	46	0	0	0	0
5	C	9	0	0	0	0
5	D	84	0	0	0	0
5	E	35	0	0	0	0
5	F	7	0	0	0	0
All	All	6640	0	5988	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:ARG:NH1	1:A:48:ARG:NH1	2.23	0.86
1:D:253:GLN:HB3	1:D:256:ARG:HD3	1.80	0.64
1:A:35:ARG:HH11	1:A:48:ARG:NH1	1.95	0.64
1:A:35:ARG:NH1	1:A:48:ARG:HH12	1.96	0.63
1:D:200:THR:HG22	1:D:248:VAL:HG22	1.83	0.61
1:D:250:PRO:HG2	1:D:253:GLN:HG3	1.83	0.60
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.37	0.59
2:B:72:PRO:HB2	2:B:97:ARG:HH12	1.71	0.56
1:A:262:GLN:HG2	1:A:269:PRO:HB3	1.89	0.55
2:B:19:LYS:O	2:B:72:PRO:HD2	2.09	0.51
1:A:204:TRP:HZ2	2:B:99:MET:HB2	1.76	0.50
1:D:2:SER:HB2	1:D:103:VAL:O	2.13	0.48
2:E:24:ASN:HB3	2:E:65:LEU:HD11	1.96	0.46
2:E:25:CYS:HB2	2:E:39:LEU:HD21	1.97	0.46
1:D:257:TYR:CD2	1:D:257:TYR:N	2.83	0.45
2:E:16:GLU:HG3	2:E:19:LYS:NZ	2.31	0.45
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.52	0.45
1:A:127:LYS:HD2	1:A:132:SER:HB2	1.98	0.44
1:A:19:GLU:HG3	1:A:20:PRO:HD2	1.99	0.43
1:A:51:TRP:CZ2	1:A:179:LEU:HD11	2.54	0.42
1:D:63:GLU:OE2	3:F:2:LEU:HB2	2.19	0.41
1:A:35:ARG:HG2	1:A:36:PHE:N	2.34	0.41
1:A:7:TYR:HB3	1:A:9:PHE:CZ	2.56	0.41
1:D:214:THR:HB	1:D:262:GLN:HB3	2.03	0.41
1:A:5:MET:HB2	1:A:168:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	268 (98%)	6 (2%)	0	100	100
1	D	273/276 (99%)	262 (96%)	11 (4%)	0	100	100
2	B	99/100 (99%)	96 (97%)	2 (2%)	1 (1%)	15	15
2	E	97/100 (97%)	93 (96%)	4 (4%)	0	100	100
3	C	8/10 (80%)	8 (100%)	0	0	100	100
3	F	8/10 (80%)	8 (100%)	0	0	100	100
All	All	759/772 (98%)	735 (97%)	23 (3%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	97	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	223 (96%)	9 (4%)	32	40
1	D	231/232 (100%)	221 (96%)	10 (4%)	29	35
2	B	96/95 (101%)	91 (95%)	5 (5%)	23	27
2	E	94/95 (99%)	89 (95%)	5 (5%)	22	26
3	C	10/10 (100%)	10 (100%)	0	100	100
3	F	10/10 (100%)	8 (80%)	2 (20%)	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	673/674 (100%)	642 (95%)	31 (5%)	27 32

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ARG
1	A	19	GLU
1	A	35	ARG
1	A	58	GLU
1	A	110	LEU
1	A	111	ARG
1	A	121	LYS
1	A	195	SER
1	A	232	GLU
2	B	0	MET
2	B	4	THR
2	B	48	LYS
2	B	70	PHE
2	B	89	GLN
1	D	35	ARG
1	D	75	ARG
1	D	111	ARG
1	D	128	GLU
1	D	131	ARG
1	D	132	SER
1	D	225	THR
1	D	226	GLN
1	D	249	VAL
1	D	257	TYR
2	E	20	SER
2	E	50	GLU
2	E	70	PHE
2	E	89	GLN
2	E	99	MET
3	F	2	LEU
3	F	8	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	32	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	141	GLN
1	D	174	ASN
1	D	180	GLN
1	D	218	GLN
1	D	260	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	301	-	5,5,5	0.07	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	276/276 (100%)	-0.18	1 (0%)	92	96	25, 38, 55, 65	0
1	D	274/276 (99%)	0.14	23 (8%)	11	16	25, 42, 70, 88	0
2	B	100/100 (100%)	-0.14	2 (2%)	65	75	24, 43, 62, 78	0
2	E	99/100 (99%)	-0.06	2 (2%)	65	75	29, 45, 73, 81	0
3	C	10/10 (100%)	-0.31	0	100	100	29, 37, 38, 39	0
3	F	10/10 (100%)	-0.28	0	100	100	32, 41, 43, 44	0
All	All	769/772 (99%)	-0.05	28 (3%)	42	55	24, 41, 66, 88	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	199	ALA	6.6
1	D	227	ASP	6.5
1	D	194	VAL	4.5
1	D	192	HIS	4.5
1	D	17	ARG	4.3
1	D	193	ALA	4.0
2	B	98	ASP	3.9
1	D	196	ASP	3.6
1	D	253	GLN	3.4
1	D	195	SER	3.3
1	D	274	TRP	3.3
1	D	16	GLY	3.1
2	E	98	ASP	3.0
1	D	18	GLY	2.7
1	D	248	VAL	2.7
1	A	184	ALA	2.7
1	D	1	GLY	2.6
1	D	247	VAL	2.5
1	D	201	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	97	ARG	2.4
1	D	197	HIS	2.4
1	D	254	GLU	2.4
1	D	15	PRO	2.3
1	D	200	THR	2.3
1	D	250	PRO	2.2
1	D	198	GLU	2.2
1	D	226	GLN	2.1
2	E	16	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	A	301	6/6	0.80	0.24	65,65,66,66	0

6.5 Other polymers (i)

There are no such residues in this entry.