



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 12:22 AM EDT

PDB ID : 7MB6
Title : SARS-CoV-2 Main Protease (Mpro) C145A in Complex with Cleavage Site Nsp6/7 (P6-P1)
Authors : Lockbaum, G.J.; Schiffer, C.A.
Deposited on : 2021-03-31
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow;"></div> </div> <p style="text-align: center;">7% 88% 11%</p>
1	B	306	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow;"></div> </div> <p style="text-align: center;">9% 86% 14%</p>
2	C	6	<div style="display: flex; align-items: center;"> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow;"></div> </div> <p style="text-align: center;">83% 17%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8991 atoms, of which 4307 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	305	4337	1427	2087	377	427	19	0	0	0
1	B	306	4451	1453	2167	384	427	20	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	ALA	CYS	engineered mutation	UNP P0DTD1
B	145	ALA	CYS	engineered mutation	UNP P0DTD1

- Molecule 2 is a protein called LYS-VAL-ALA-THR-VAL-GLN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	6	98	28	53	8	9	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	50	Total O 50 50	0	0
3	B	55	Total O 55 55	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3C-like proteinase



- Molecule 1: 3C-like proteinase



- Molecule 2: LYS-VAL-ALA-THR-VAL-GLN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	124.85Å 80.75Å 62.69Å 90.00° 91.33° 90.00°	Depositor
Resolution (Å)	28.58 – 2.21 28.58 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.4 (28.58-2.21) 95.1 (28.58-2.21)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.22Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.223 , 0.272 0.222 , 0.270	Depositor DCC
R_{free} test set	1562 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtrriage
Anisotropy	0.177	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.015 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8991	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2302	0.51	0/3146
1	B	0.27	0/2336	0.50	0/3186
2	C	0.33	0/44	0.58	0/57
All	All	0.27	0/4682	0.51	0/6389

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	-2	VAL	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2250	2087	2088	20	0
1	B	2284	2167	2168	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	45	53	53	0	0
3	A	50	0	0	0	1
3	B	55	0	0	0	1
All	All	4684	4307	4309	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:THR:HG21	1:A:44:CYS:O	1.97	0.64
1:B:111:THR:HG22	1:B:129:ALA:HB2	1.84	0.59
1:B:213:ILE:HG21	1:B:300:CYS:HB3	1.85	0.58
1:B:225:THR:O	1:B:262:LEU:HD22	2.04	0.57
1:A:17:MET:HG3	1:A:117:CYS:SG	2.45	0.57
1:B:267:SER:O	1:B:271:LEU:HD13	2.06	0.56
1:B:27:LEU:HD21	1:B:42:VAL:HB	1.88	0.55
1:B:234:ALA:HB1	1:B:239:TYR:CB	2.37	0.54
1:B:227:LEU:HD11	1:B:242:LEU:HD23	1.88	0.54
1:A:220:LEU:H	1:A:220:LEU:HD12	1.72	0.53
1:A:230:PHE:CZ	1:A:268:LEU:HD23	2.44	0.53
1:B:244:GLN:O	1:B:247:VAL:HG22	2.09	0.52
1:A:210:ALA:HB2	1:A:296:VAL:HG13	1.92	0.51
1:B:233:VAL:HG11	1:B:269:LYS:HD3	1.93	0.51
1:A:8:PHE:HB3	1:A:152:ILE:HD12	1.94	0.50
1:B:234:ALA:HB1	1:B:239:TYR:HB2	1.93	0.50
1:B:239:TYR:CE1	1:B:272:LEU:HD21	2.47	0.49
1:B:86:VAL:HG23	1:B:179:GLY:HA2	1.94	0.48
1:A:49:MET:HA	1:A:52:PRO:HG3	1.94	0.48
1:A:66:PHE:CE1	1:A:87:LEU:HD21	2.48	0.48
1:A:247:VAL:HG13	1:A:261:VAL:HG21	1.97	0.46
1:B:5:LYS:HD3	1:B:291:PHE:CZ	2.51	0.46
1:B:5:LYS:NZ	1:B:288:GLU:OE2	2.50	0.45
1:B:78:ILE:HG13	1:B:90:LYS:HG2	1.99	0.45
1:A:218:TRP:CH2	1:A:271:LEU:HD21	2.52	0.44
1:B:167:LEU:HB3	1:B:168:PRO:HD2	1.99	0.44
1:A:70:ALA:CB	1:A:75:LEU:HD11	2.48	0.44
1:B:207:TRP:CH2	1:B:281:ILE:HB	2.53	0.43
1:B:136:ILE:HD11	1:B:140:PHE:HE2	1.83	0.43
1:B:234:ALA:HB1	1:B:239:TYR:HB3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ASP:OD1	1:A:187:ASP:N	2.48	0.42
1:B:225:THR:OG1	1:B:226:THR:N	2.53	0.42
1:A:113:SER:O	1:A:149:GLY:HA2	2.20	0.42
1:A:19:GLN:OE1	1:A:26:THR:HG21	2.20	0.41
1:B:204:VAL:O	1:B:208:LEU:HG	2.21	0.41
1:A:230:PHE:HZ	1:A:268:LEU:HD23	1.84	0.41
1:B:56:ASP:O	1:B:59:ILE:HG22	2.20	0.41
1:B:148:VAL:HA	1:B:161:TYR:O	2.21	0.41
1:B:169:THR:HG23	1:B:171:VAL:HG22	2.03	0.40
1:A:49:MET:HB3	1:A:189:GLN:HG3	2.03	0.40
1:A:81:SER:O	1:A:87:LEU:HD12	2.21	0.40
1:A:230:PHE:CE1	1:A:268:LEU:HD23	2.57	0.40
1:A:30:LEU:HD21	1:A:32:LEU:HD11	2.03	0.40
1:A:101:TYR:HA	1:A:157:VAL:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:401:HOH:O	3:B:425:HOH:O[4_555]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/306 (99%)	291 (96%)	12 (4%)	0	100	100
1	B	304/306 (99%)	291 (96%)	13 (4%)	0	100	100
2	C	4/6 (67%)	4 (100%)	0	0	100	100
All	All	611/618 (99%)	586 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/262 (88%)	229 (99%)	3 (1%)	69	80
1	B	240/262 (92%)	237 (99%)	3 (1%)	69	80
2	C	5/5 (100%)	5 (100%)	0	100	100
All	All	477/529 (90%)	471 (99%)	6 (1%)	69	80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ARG
1	A	5	LYS
1	A	245	ASP
1	B	27	LEU
1	B	107	GLN
1	B	284	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/306 (99%)	0.36	20 (6%) 18 17	29, 53, 81, 96	0
1	B	306/306 (100%)	0.51	29 (9%) 8 6	32, 51, 79, 94	0
2	C	6/6 (100%)	0.71	0 100 100	45, 56, 63, 76	0
All	All	617/618 (99%)	0.44	49 (7%) 12 11	29, 52, 81, 96	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	PHE	4.6
1	B	272	LEU	4.4
1	B	220	LEU	4.2
1	B	29	GLY	4.2
1	B	255	ALA	4.0
1	A	260	ALA	3.7
1	A	226	THR	3.6
1	B	198	THR	3.5
1	B	30	LEU	3.5
1	B	276	MET	3.4
1	B	218	TRP	3.3
1	B	38	CYS	3.0
1	B	219	PHE	3.0
1	B	148	VAL	3.0
1	B	279	ARG	2.9
1	B	146	GLY	2.9
1	B	235	MET	2.9
1	B	50	LEU	2.8
1	A	233	VAL	2.7
1	A	198	THR	2.7
1	A	277	ASN	2.6
1	A	242	LEU	2.6
1	B	162	MET	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	117	CYS	2.5
1	B	18	VAL	2.5
1	B	216	ASP	2.5
1	B	213	ILE	2.4
1	B	28	ASN	2.4
1	B	17	MET	2.4
1	B	281	ILE	2.4
1	A	278	GLY	2.4
1	A	165	MET	2.4
1	A	29	GLY	2.3
1	B	149	GLY	2.3
1	A	234	ALA	2.3
1	A	56	ASP	2.3
1	A	229	ASP	2.2
1	B	37	TYR	2.2
1	A	146	GLY	2.2
1	B	36	VAL	2.1
1	A	144	SER	2.1
1	B	277	ASN	2.1
1	A	72	ASN	2.1
1	A	283	GLY	2.1
1	A	128	CYS	2.1
1	B	273	GLN	2.0
1	A	169	THR	2.0
1	B	39	PRO	2.0
1	A	206	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.