

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 09:30 AM EDT

PDB ID	:	6MDF
Title	:	Mevalonate kinase from Methanosarcina mazei with 5-phosphomevalonate
		bound
Authors	:	Miller, B.R.; Kung, Y.
Deposited on	:	2018-09-04
Resolution	:	2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.46 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



6MDF

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Mevalonate kinase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	303	Total	С	Ν	0	\mathbf{S}	0	1	0
	11	000	2202	1387	377	428	10	0	1	0
1	Р	301	Total	С	Ν	0	\mathbf{S}	0	0	0
	D	301	2143	1350	368	416	9	0	Δ	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-1	GLY	-	expression tag	UNP Q8PW39
А	0	HIS	-	expression tag	UNP Q8PW39
В	-1	GLY	-	expression tag	UNP Q8PW39
В	0	HIS	-	expression tag	UNP Q8PW39


• Molecule 2 is (3R)-3-HYDROXY-3-METHYL-5-(PHOSPHONOOXY)PENTANOIC ACID (three-letter code: PMV) (formula: C₆H₁₃O₇P) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C O P 14 6 7 1	0	0
2	В	1	Total C O P 14 6 7 1	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	133	Total O 133 133	0	0
4	В	75	Total O 75 75	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	53.97Å 58.58 Å 190.03 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.92 - 2.46	Depositor
% Data completeness	98.0 (51.92-2.46)	Depositor
(in resolution range)		-
R _{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.46 (at 2.45 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.200 , 0.267	Depositor
Wilson B-factor $(Å^2)$	39.6	Xtriage
Anisotropy	0.381	Xtriage
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4659	wwPDB-VP
Average B, all atoms $(Å^2)$	46.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	T a	Chain	Dec	Link	Bo	ond leng	ths	В	ond ang	les
Mol	Type	Chain	Res	LINK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	PMV	А	401	-	12,13,13	1.03	0	13,19,19	0.84	0
3	GOL	А	403	-	$5,\!5,\!5$	0.90	0	$5,\!5,\!5$	1.02	0
3	GOL	А	405	-	$5,\!5,\!5$	0.90	0	$5,\!5,\!5$	0.99	0
3	GOL	А	406	-	$5,\!5,\!5$	0.91	0	$5,\!5,\!5$	0.98	0
3	GOL	А	407	-	$5,\!5,\!5$	0.94	0	$5,\!5,\!5$	0.94	0
3	GOL	А	408	-	$5,\!5,\!5$	0.94	0	$5,\!5,\!5$	0.98	0
3	GOL	В	504	-	$5,\!5,\!5$	0.90	0	$5,\!5,\!5$	1.03	0
3	GOL	В	507	-	$5,\!5,\!5$	0.91	0	$5,\!5,\!5$	1.02	0
3	GOL	А	404	-	$5,\!5,\!5$	0.91	0	$5,\!5,\!5$	0.99	0
3	GOL	В	505	-	$5,\!5,\!5$	0.93	0	$5,\!5,\!5$	0.94	0
3	GOL	В	506	-	$5,\!5,\!5$	0.91	0	$5,\!5,\!5$	0.93	0
3	GOL	В	502	-	$5,\!5,\!5$	0.96	0	$5,\!5,\!5$	0.82	0
3	GOL	В	503	-	$5,\!5,\!5$	0.94	0	$5,\!5,\!5$	0.98	0
3	GOL	А	402	-	$5,\!5,\!5$	0.90	0	$5,\!5,\!5$	0.92	0
2	PMV	В	501	-	12,13,13	1.08	0	$13,\!19,\!19$	0.86	0

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PMV	А	401	-	-	8/13/13/13	-
3	GOL	А	403	-	-	0/4/4/4	-
3	GOL	А	405	-	-	2/4/4/4	-
3	GOL	А	406	-	-	2/4/4/4	-
3	GOL	А	407	-	-	0/4/4/4	-
3	GOL	А	408	-	-	0/4/4/4	-
3	GOL	В	504	-	-	2/4/4/4	-
3	GOL	В	507	-	-	0/4/4/4	-
3	GOL	А	404	-	-	1/4/4/4	-
3	GOL	В	505	-	-	2/4/4/4	-
3	GOL	В	506	-	-	2/4/4/4	-
3	GOL	В	502	-	-	4/4/4/4	-
3	GOL	В	503	-	-	0/4/4/4	-
3	GOL	А	402	-	-	2/4/4/4	-
2	PMV	В	501	_	_	3/13/13/13	_



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	401	PMV	C5-O5-P-OP1
2	А	401	PMV	O3A-C3-C4-C5
2	А	401	PMV	C1-C2-C3-C4
2	А	401	PMV	C1-C2-C3-O3A
2	А	401	PMV	C1-C2-C3-C3A
2	В	501	PMV	C5-O5-P-OP2
2	В	501	PMV	C5-O5-P-OP3
3	А	406	GOL	O1-C1-C2-O2
3	А	406	GOL	O1-C1-C2-C3
3	В	502	GOL	O1-C1-C2-C3
3	В	502	GOL	C1-C2-C3-O3
3	В	506	GOL	O1-C1-C2-C3
3	В	502	GOL	O1-C1-C2-O2
3	В	502	GOL	O2-C2-C3-O3
3	В	506	GOL	O1-C1-C2-O2
3	А	402	GOL	C1-C2-C3-O3
3	А	405	GOL	O1-C1-C2-C3
3	В	504	GOL	O1-C1-C2-C3
3	В	505	GOL	O1-C1-C2-C3
3	В	504	GOL	O1-C1-C2-O2
2	А	401	PMV	C2-C3-C4-C5
2	В	501	PMV	C5-O5-P-OP1
3	А	405	GOL	O1-C1-C2-O2
2	А	401	PMV	C3A-C3-C4-C5
3	А	402	GOL	O2-C2-C3-O3
3	В	505	GOL	O1-C1-C2-O2
2	А	401	PMV	C5-O5-P-OP3
3	А	404	GOL	O1-C1-C2-C3


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

