

wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2020 – 03:54 AM BST

PDB ID : 4MOM

Title: Pyranose 2-oxidase H450G mutant with 3-fluorinated galactose Authors: Tan, T.C.; Spadiut, O.; Gandini, R.; Haltrich, D.; Divne, C.

Deposited on : 2013-09-12

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	633	80%	11%		9%
1	В	633	78%	12%		9%
1	С	633	74%	16%	·	9%
1	D	633	77%	12%		9%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 19522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	576	Total	С	N	О	S	0	0	0
1	A	370	4536	2864	775	872	25	U	0	
1	В	576	Total	С	N	О	S	0	0	0
1	Б	370	4536	2864	775	872	25	U	U	0
1	С	574	Total	С	N	О	S	0	0	0
1		374	4520	2855	773	868	24	0	0	
1	D	574	Total	С	N	О	S	0	0	0
1		374	4520	2855	773	868	24	U	U	U

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	cloning artifact	UNP Q7ZA32
A	450	GLY	HIS	engineered mutation	UNP Q7ZA32
A	623	ALA	-	expression tag	UNP Q7ZA32
A	624	ALA	-	expression tag	UNP Q7ZA32
A	625	ALA	-	expression tag	UNP Q7ZA32
A	626	LEU	_	expression tag	UNP Q7ZA32
A	627	GLU	_	expression tag	UNP Q7ZA32
A	628	HIS	-	expression tag	UNP Q7ZA32
A	629	HIS	_	expression tag	UNP Q7ZA32
A	630	HIS	-	expression tag	UNP Q7ZA32
A	631	HIS	_	expression tag	UNP Q7ZA32
A	632	HIS	_	expression tag	UNP Q7ZA32
A	633	HIS	_	expression tag	UNP Q7ZA32
В	2	ALA	SER	cloning artifact	UNP Q7ZA32
В	450	GLY	HIS	engineered mutation	UNP Q7ZA32
В	623	ALA	_	expression tag	UNP Q7ZA32
В	624	ALA	=	expression tag	UNP Q7ZA32
В	625	ALA	-	expression tag	UNP Q7ZA32
В	626	LEU	=	expression tag	UNP Q7ZA32
В	627	GLU	-	expression tag	UNP Q7ZA32
В	628	HIS	-	expression tag	UNP Q7ZA32

 $\overline{\textit{Continued}}$ on next page...




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	629	HIS	_	expression tag	UNP Q7ZA32
В	630	HIS	_	expression tag	UNP Q7ZA32
В	631	HIS	-	expression tag	UNP Q7ZA32
В	632	HIS	_	expression tag	UNP Q7ZA32
В	633	HIS	-	expression tag	UNP Q7ZA32
С	2	ALA	SER	cloning artifact	UNP Q7ZA32
С	450	GLY	HIS	engineered mutation	UNP Q7ZA32
С	623	ALA	-	expression tag	UNP Q7ZA32
С	624	ALA	-	expression tag	UNP Q7ZA32
С	625	ALA	_	expression tag	UNP Q7ZA32
С	626	LEU	_	expression tag	UNP Q7ZA32
С	627	GLU	_	expression tag	UNP Q7ZA32
С	628	HIS	_	expression tag	UNP Q7ZA32
С	629	HIS	-	expression tag	UNP Q7ZA32
С	630	HIS	_	expression tag	UNP Q7ZA32
С	631	HIS	_	expression tag	UNP Q7ZA32
С	632	HIS	_	expression tag	UNP Q7ZA32
С	633	HIS	_	expression tag	UNP Q7ZA32
D	2	ALA	SER	cloning artifact	UNP Q7ZA32
D	450	GLY	HIS	engineered mutation	UNP Q7ZA32
D	623	ALA	_	expression tag	UNP Q7ZA32
D	624	ALA	_	expression tag	UNP Q7ZA32
D	625	ALA	-	expression tag	UNP Q7ZA32
D	626	LEU	-	expression tag	UNP Q7ZA32
D	627	GLU	_	expression tag	UNP Q7ZA32
D	628	HIS	_	expression tag	UNP Q7ZA32
D	629	HIS	-	expression tag	UNP Q7ZA32
D	630	HIS	-	expression tag	UNP Q7ZA32
D	631	HIS	-	expression tag	UNP Q7ZA32
D	632	HIS	-	expression tag	UNP Q7ZA32
D	633	HIS	-	expression tag	UNP Q7ZA32


• Molecule 2 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (three-letter code: FDA) (formula: $C_{27}H_{35}N_9O_{15}P_2$).





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0
2	A	1	53	27	9	15	2	U	0
9	В	1	Total	С	N	О	Р	0	0
2	Б	1	53	27	9	15	2	U	0
9	С	1	Total	С	N	О	Р	0	0
2		1	53	27	9	15	2	U	0
9	D	1	Total	С	N	О	Р	0	0
	ש	1	53	27	9	15	2	U	U


• Molecule 3 is 3-deoxy-3-fluoro-beta-D-galactopyranose (three-letter code: 2H5) (formula: $C_6H_{11}FO_5$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C F O 12 6 1 5	0	0
3	В	1	Total C F O	0	0
3	C	1	12 6 1 5 Total C F O	0	0
		1	12 6 1 5 Total C F O	0	0
3	D	1	12 6 1 5	0	0


• Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
1	Δ	1	Total C O	0	0
4	11	1	12 8 4	U	U
1	В	1	Total C O	0	0
4	D	1	16 10 6	U	U
4	C	1	Total C O	0	0
4	C	1	16 10 6	0	U
1	D	1	Total C O	0	0
4	ש	1	16 10 6	U	U

• Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
5	D	1	Total	С	N	О	S	0	0	
9	Б	1	12	6	1	4	1	0	U	
5	D	1	Total	С	N	О	S	0	0	
3	ש	1	11	6	1	3	1	0	U	

• Molecule 6 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
6	A	268	Total O 268 268	0	0
6	В	348	Total O 348 348	0	0
6	С	183	Total O 183 183	0	0
6	D	268	Total O 268 268	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyranose 2-oxidase



• Molecule 1: Pyranose 2-oxidase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	99.90Å 102.42Å 137.37Å	Danagitan
a, b, c, α , β , γ	90.00° 91.19° 90.00°	Depositor
Resolution (Å)	50.00 - 1.90	Depositor
rtesoration (A)	49.16 - 1.90	EDS
% Data completeness	99.2 (50.00-1.90)	Depositor
(in resolution range)	99.2 (49.16-1.90)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.85 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.238 , 0.280	Depositor
it, it free	0.247 , 0.289	DCC
R_{free} test set	3865 reflections (1.79%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.35\;,42.8$	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
	0.012 for -k,-h,-l	
Estimated twinning fraction	0.001 for k,h,-l	Xtriage
	0.020 for h,-k,-l	
F_o, F_c correlation	0.92	EDS
Total number of atoms	19522	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FDA, 2H5, MES, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	ond lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z >5	
1	A	1.03	3/4651~(0.1%)	0.99	7/6323~(0.1%)	
1	В	1.05	5/4651~(0.1%)	1.01	$10/6323 \; (0.2\%)$	
1	С	0.83	2/4635~(0.0%)	0.91	$11/6302 \; (0.2\%)$	
1	D	0.92	3/4635~(0.1%)	0.93	7/6302~(0.1%)	
All	All	0.96	$13/18572 \ (0.1\%)$	0.96	$35/25250 \; (0.1\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	В	564	CYS	CB-SG	-5.96	1.72	1.81
1	A	445	TRP	CD2-CE2	5.53	1.48	1.41
1	В	493	ASP	CB-CG	5.47	1.63	1.51
1	D	296	GLU	CD-OE1	5.42	1.31	1.25
1	В	407	TRP	CD2-CE2	5.41	1.47	1.41

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	139	ARG	NE-CZ-NH1	9.84	125.22	120.30
1	В	139	ARG	NE-CZ-NH2	-9.62	115.49	120.30
1	A	139	ARG	NE-CZ-NH1	9.31	124.96	120.30
1	D	470	ASP	CB-CG-OD1	8.68	126.11	118.30

Continued on next page...



 $Continued\ from\ previous\ page...$

\mathbf{Mol}	Chain	Res	Type	${f Atoms}$	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	D	139	ARG	NE-CZ-NH2	-8.34	116.13	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	436	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	4536	0	4384	34	0
1	В	4536	0	4384	47	0
1	С	4520	0	4371	62	0
1	D	4520	0	4371	43	0
2	A	53	0	28	3	0
2	В	53	0	29	1	0
2	С	53	0	29	1	0
2	D	53	0	29	2	0
3	A	12	0	11	4	0
3	В	12	0	11	0	0
3	С	12	0	11	2	0
3	D	12	0	11	1	0
4	A	12	0	14	0	0
4	В	16	0	22	1	0
4	С	16	0	22	0	0
4	D	16	0	22	3	0
5	В	12	0	13	1	0
5	D	11	0	9	1	0
6	A	268	0	0	2	0
6	В	348	0	0	12	0
6	С	183	0	0	4	0
6	D	268	0	0	3	0
All	All	19522	0	17771	191	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
5:D:804:MES:C2	5:D:804:MES:C6	2.11	1.26
1:C:361:LEU:HD22	1:C:546:VAL:HG22	1.38	1.02
1:B:110:GLN:HE21	1:B:167:HIS:HD1	1.07	0.96
1:C:110:GLN:HE21	1:C:167:HIS:HD1	1.14	0.93
1:A:460:GLN:HE22	1:A:463:ILE:H	0.93	0.92

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	574/633 (91%)	554 (96%)	19 (3%)	1 (0%)	47 38
1	В	574/633 (91%)	555 (97%)	19 (3%)	0	100 100
1	С	572/633 (90%)	547 (96%)	24 (4%)	1 (0%)	47 38
1	D	572/633~(90%)	547 (96%)	25 (4%)	0	100 100
All	All	$2292/2532 \ (90\%)$	2203 (96%)	87 (4%)	2 (0%)	51 43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Α	82	SER
1	С	617	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	503/547 (92%)	482 (96%)	21 (4%)	30 20
1	В	503/547~(92%)	489 (97%)	14 (3%)	43 36
1	С	501/547 (92%)	480 (96%)	21 (4%)	30 20
1	D	501/547~(92%)	480 (96%)	21 (4%)	30 20
All	All	$2008/2188 \; (92\%)$	1931 (96%)	77 (4%)	33 24

5 of 77 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	593	ASN
1	С	210	THR
1	D	421	GLU
1	С	45	ILE
1	С	168	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	207	GLN
1	С	263	GLN
1	D	324	HIS
1	С	110	GLN
1	D	341	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trans	Chain	Dag	Link	В	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FDA	A	801	1	51,58,58	1.60	8 (15%)	60,89,89	3.35	21 (35%)
3	2H5	A	802	-	12,12,12	1.19	1 (8%)	17,17,17	1.66	3 (17%)
4	1PE	В	704	-	15,15,15	0.63	0	14,14,14	0.28	0
2	FDA	D	801	1	51,58,58	1.57	8 (15%)	60,89,89	3.16	22 (36%)
5	MES	В	701	-	12,12,12	2.42	1 (8%)	14,16,16	1.69	3 (21%)
5	MES	D	804	-	10,10,12	2.00	1 (10%)	11,13,16	1.71	3 (27%)
4	1PE	D	803	-	15,15,15	0.61	0	14,14,14	0.22	0
4	1PE	С	803	-	15,15,15	0.51	0	14,14,14	0.23	0
2	FDA	В	702	1	51,58,58	1.70	13 (25%)	60,89,89	3.99	19 (31%)
3	2H5	D	802	-	12,12,12	0.66	0	17,17,17	1.66	3 (17%)
4	1PE	A	803	-	11,11,15	0.68	0	10,10,14	0.27	0
3	2H5	В	703	-	12,12,12	1.16	2 (16%)	17,17,17	2.27	8 (47%)
3	2H5	С	802	-	12,12,12	0.86	0	17,17,17	2.69	7 (41%)
2	FDA	С	801	1	51,58,58	1.73	11 (21%)	60,89,89	3.73	24 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDA	A	801	1	-	2/30/50/50	0/6/6/6
3	2H5	A	802	_	-	0/2/22/22	0/1/1/1
4	1PE	В	704	-	-	7/13/13/13	-
2	FDA	D	801	1	-	2/30/50/50	0/6/6/6
5	MES	В	701	_	-	0/6/14/14	0/1/1/1
5	MES	D	804	_	_	4/10/10/14	-
4	1PE	D	803	-	-	7/13/13/13	-

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	${f Torsions}$	Rings
4	1PE	С	803	_	-	6/13/13/13	-
2	FDA	В	702	1	-	5/30/50/50	0/6/6/6
3	2H5	D	802	-	-	2/2/22/22	0/1/1/1
4	1PE	A	803	-	-	4/9/9/13	-
3	2H5	В	703	_	-	0/2/22/22	0/1/1/1
3	2H5	С	802	_	-	2/2/22/22	0/1/1/1
2	FDA	С	801	1	-	3/30/50/50	0/6/6/6

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(\operatorname{\AA})$
5	В	701	MES	C8-S	-7.99	1.66	1.77
5	D	804	MES	C8-S	-6.04	1.68	1.77
2	С	801	FDA	C4-C4X	5.21	1.50	1.41
2	В	702	FDA	C4-C4X	4.73	1.49	1.41
2	D	801	FDA	C4-C4X	4.55	1.49	1.41

The worst 5 of 113 bond angle outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	С	801	FDA	C4-C4X-C10	-21.73	105.57	119.95
2	В	702	FDA	C4-N3-C2	16.57	129.13	115.14
2	A	801	FDA	C4-N3-C2	15.35	128.10	115.14
2	В	702	FDA	C1'-N10-C9A	14.25	129.51	118.29
2	D	801	FDA	C4-C4X-C10	-13.27	111.17	119.95

There are no chirality outliers.

5 of 44 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	801	FDA	PA-O3P-P-O5'
3	С	802	2H5	O5-C5-C6-O6
4	D	803	1PE	ОН7-С16-С26-ОН6
4	A	803	1PE	ОН6-С15-С25-ОН5
4	D	803	1PE	OH2-C12-C22-OH3

There are no ring outliers.


11 monomers are involved in 17 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	FDA	3	0
3	A	802	2H5	4	0
4	В	704	1PE	1	0
2	D	801	FDA	2	0
5	В	701	MES	1	0
5	D	804	MES	1	0
4	D	803	1PE	3	0
2	В	702	FDA	1	0
3	D	802	2H5	1	0
3	С	802	2H5	2	0
2	С	801	FDA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	576/633~(90%)	0.27	16 (2%) 53 56	11, 19, 37, 62	0
1	В	576/633 (90%)	0.32	13 (2%) 60 63	10, 18, 35, 61	0
1	С	574/633 (90%)	0.58	35 (6%) 21 24	14, 29, 51, 88	0
1	D	574/633 (90%)	0.41	20 (3%) 44 47	12, 22, 43, 64	0
All	All	$2300/2532 \ (90\%)$	0.39	84 (3%) 41 44	10, 22, 44, 88	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	618	PHE	6.4
1	D	343	ALA	5.7
1	D	382	ILE	5.7
1	С	343	ALA	5.5
1	С	618	PHE	5.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	1PE	С	803	16/16	0.75	0.25	43,50,64,69	0
4	1PE	A	803	12/16	0.83	0.20	27,31,36,36	0
4	1PE	В	704	16/16	0.84	0.19	28,41,53,54	0
4	1PE	D	803	16/16	0.84	0.17	31,37,48,51	0
3	2H5	С	802	12/12	0.86	0.15	39,42,47,50	0
3	2H5	A	802	12/12	0.87	0.14	29,31,34,34	0
5	MES	D	804	11/12	0.89	0.13	31,38,44,49	0
3	2H5	D	802	12/12	0.90	0.12	26,30,33,34	0
3	2H5	В	703	12/12	0.91	0.12	24,26,28,29	0
5	MES	В	701	12/12	0.93	0.22	28,38,41,44	0
2	FDA	С	801	53/53	0.95	0.11	18,20,28,32	0
2	FDA	В	702	53/53	0.97	0.10	6,9,15,20	0
2	FDA	A	801	53/53	0.97	0.10	7,9,12,16	0
2	FDA	D	801	53/53	0.97	0.09	11,13,19,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around FDA A 801: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

