



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 6, 2021 – 02:05 PM EDT

PDB ID : 7MOX
Title : Structure of HDAC2 in complex with an inhibitor (compound 14)
Authors : Klein, D.J.; Yu, W.
Deposited on : 2021-05-03
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.22
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.22


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	 2% 91% 7%
1	B	376	 0% 90% 7%
1	C	376	 8% 93%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone deacetylase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	368	3019	1937	507	548	27	0	17	0
1	B	366	3023	1938	503	553	29	0	19	0
1	C	368	2990	1909	507	548	26	0	6	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca 2 2	0	0
4	B	2	Total Ca 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ca	0	0
			2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is (1S)-N-[(1S)-7,7-dihydroxy-1-{4-[(1R,4S)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-6-yl]-1H-imidazol-2-yl}nonyl]-6-methyl-6-azaspiro[2.5]octane-1-carboxamide (three-letter code: ZLS) (formula: C₃₂H₄₆N₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	39	32	4	3	0	0
6	B	1	39	32	4	3	0	0
6	C	1	39	32	4	3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	352	352	352	0	0
7	B	411	415	415	0	4
7	C	302	305	305	0	3


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone deacetylase 2



- Molecule 1: Histone deacetylase 2



- Molecule 1: Histone deacetylase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.56Å 99.19Å 139.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.42 – 1.69 24.42 – 1.69	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.42-1.69) 100.0 (24.42-1.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.69Å)	Xtrriage
Refinement program	BUSTER 2.11.8 (10-DEC-2020)	Depositor
R, R_{free}	0.176 , 0.202 0.173 , 0.200	Depositor DCC
R_{free} test set	7204 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtrriage
Anisotropy	0.459	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10329	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, SO4, ZLS, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3126	0.61	1/4220 (0.0%)
1	B	0.45	0/3142	0.62	0/4241
1	C	0.38	0/3075	0.56	0/4152
All	All	0.43	0/9343	0.60	1/12613 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	THR	N-CA-C	-5.01	97.48	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3019	0	2932	11	0
1	B	3023	0	2943	15	0
1	C	2990	0	2888	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
3	C	20	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
5	A	21	0	30	0	0
5	B	14	0	20	1	0
5	C	14	0	20	1	0
6	A	39	0	0	0	0
6	B	39	0	0	0	0
6	C	39	0	0	0	0
7	A	352	0	0	0	0
7	B	415	0	0	0	0
7	C	305	0	0	0	0
All	All	10329	0	8833	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:282:HIS:HD2	1:B:316:GLU:OE2	1.60	0.84
1:C:29:HIS:HD2	1:C:31:MET:H	1.40	0.70
1:B:241:ILE:HG22	1:B:364:LYS:HE2	1.73	0.68
1:A:201:LYS:HD2	1:A:274[A]:CYS:SG	2.34	0.68
1:C:328:GLU:HG3	1:C:343:LYS:NZ	2.09	0.67
1:A:362:LYS:HE3	1:A:366:ARG:HE	1.60	0.65
1:B:44:LEU:HB2	1:B:49:TYR:CZ	2.36	0.61
1:B:43[A]:LEU:HD12	1:B:49:TYR:HE1	1.65	0.61
1:C:90:LYS:NZ	1:C:90:LYS:HA	2.17	0.59
1:B:43[A]:LEU:HD12	1:B:49:TYR:CE1	2.39	0.58
1:B:52[A]:MET:HB3	1:B:321:LEU:HD21	1.89	0.54
1:C:328:GLU:HG3	1:C:343:LYS:HZ3	1.72	0.53
1:A:241:ILE:HG12	1:A:360:MET:CE	2.41	0.51
1:C:52:MET:HG2	1:C:321:LEU:HD21	1.93	0.50
1:C:328:GLU:HG3	1:C:343:LYS:HZ2	1.77	0.49
1:A:52:MET:HG2	1:A:321:LEU:HD21	1.95	0.49
1:A:241:ILE:HG12	1:A:360:MET:HE3	1.94	0.49
5:B:406:PEG:O1	5:C:408:PEG:O4	2.27	0.48
1:B:43[B]:LEU:HD23	1:B:48:LEU:HD12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:90:LYS:HA	1:C:90:LYS:HZ2	1.81	0.46
1:A:51:LYS:NZ	1:A:323[A]:CYS:SG	2.89	0.46
1:B:139:GLY:HA2	1:B:157:ILE:HD11	1.97	0.46
1:B:263:GLY:O	1:B:282:HIS:HE1	1.99	0.46
1:C:139:GLY:HA2	1:C:157:ILE:HD11	1.98	0.45
1:B:201[B]:LYS:HD3	1:B:274[B]:CYS:HG	1.81	0.44
1:A:267:LEU:HD11	1:A:312:CYS:HB2	2.00	0.43
1:C:29:HIS:CD2	1:C:31:MET:H	2.27	0.43
1:C:241:ILE:HG21	1:C:360:MET:HE3	2.01	0.42
1:A:147:GLU:HG2	1:A:148:ALA:O	2.19	0.42
1:B:241:ILE:HG12	1:B:360:MET:CE	2.50	0.42
1:B:267:LEU:HD11	1:B:312:CYS:HB2	2.01	0.42
1:C:350:ASN:H	1:C:350:ASN:HD22	1.68	0.42
1:A:10:LYS:HB3	1:A:131:ASP:HB2	2.01	0.41
1:C:90:LYS:HA	1:C:90:LYS:HZ3	1.85	0.41
1:A:44:LEU:HD21	1:A:49[B]:TYR:CD1	2.56	0.41
1:B:201[B]:LYS:HG2	1:B:205:TYR:CD2	2.56	0.41
1:B:231:ASP:OD1	1:B:274[B]:CYS:SG	2.78	0.40
1:B:72:GLU:HG2	1:B:95:PHE:HZ	1.86	0.40
1:A:76:PHE:HE2	1:A:97:VAL:HG21	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	377/376 (100%)	372 (99%)	5 (1%)	0	100 100
1	B	380/376 (101%)	375 (99%)	5 (1%)	0	100 100
1	C	370/376 (98%)	365 (99%)	5 (1%)	0	100 100
All	All	1127/1128 (100%)	1112 (99%)	15 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/322 (101%)	318 (98%)	8 (2%)	47	29
1	B	331/322 (103%)	325 (98%)	6 (2%)	59	43
1	C	321/322 (100%)	313 (98%)	8 (2%)	47	29
All	All	978/966 (101%)	956 (98%)	22 (2%)	53	34

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LYS
1	A	140	LEU
1	A	141	HIS
1	A	212[A]	LEU
1	A	212[B]	LEU
1	A	227	PHE
1	A	240[A]	GLN
1	A	350	ASN
1	B	11	LYS
1	B	51	LYS
1	B	52[A]	MET
1	B	52[B]	MET
1	B	141	HIS
1	B	254	GLN
1	C	27	GLN
1	C	44	LEU
1	C	90	LYS
1	C	140	LEU
1	C	141	HIS
1	C	337	TYR
1	C	347	SER
1	C	350	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	350	ASN
1	A	365	GLN
1	B	240	GLN
1	B	282	HIS
1	B	354	GLN
1	C	29	HIS
1	C	169[A]	GLN
1	C	240	GLN
1	C	350	ASN
1	C	354	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 9 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	403	-	4,4,4	0.28	0	6,6,6	0.65	0
3	SO4	C	404	-	4,4,4	0.13	0	6,6,6	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PEG	C	409	-	6,6,6	0.11	0	5,5,5	0.05	0
3	SO4	B	402	-	4,4,4	0.13	0	6,6,6	0.15	0
3	SO4	A	404	-	4,4,4	0.14	0	6,6,6	0.06	0
5	PEG	A	409	-	6,6,6	0.11	0	5,5,5	0.09	0
6	ZLS	B	408	2	40,44,44	0.58	0	46,66,66	0.67	0
5	PEG	A	410	-	6,6,6	0.09	0	5,5,5	0.03	0
6	ZLS	C	410	2	40,44,44	0.57	0	46,66,66	0.65	0
5	PEG	B	407	-	6,6,6	0.07	0	5,5,5	0.10	0
3	SO4	C	402	-	4,4,4	0.15	0	6,6,6	0.06	0
6	ZLS	A	411	2	40,44,44	0.59	0	46,66,66	0.67	0
5	PEG	B	406	-	6,6,6	0.08	0	5,5,5	0.10	0
3	SO4	C	403	-	4,4,4	0.27	0	6,6,6	0.16	0
3	SO4	A	402	-	4,4,4	0.13	0	6,6,6	0.14	0
3	SO4	C	405	-	4,4,4	0.14	0	6,6,6	0.08	0
3	SO4	A	405	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	B	403	-	4,4,4	0.16	0	6,6,6	0.07	0
5	PEG	C	408	-	6,6,6	0.06	0	5,5,5	0.05	0
5	PEG	A	408	-	6,6,6	0.13	0	5,5,5	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	A	409	-	-	2/4/4/4	-
6	ZLS	B	408	2	-	0/24/65/65	0/7/6/6
5	PEG	C	409	-	-	0/4/4/4	-
6	ZLS	A	411	2	-	0/24/65/65	0/7/6/6
5	PEG	B	406	-	-	0/4/4/4	-
5	PEG	A	410	-	-	1/4/4/4	-
6	ZLS	C	410	2	-	0/24/65/65	0/7/6/6
5	PEG	B	407	-	-	1/4/4/4	-
5	PEG	C	408	-	-	1/4/4/4	-
5	PEG	A	408	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	410	PEG	C4-C3-O2-C2
5	B	407	PEG	O1-C1-C2-O2
5	A	409	PEG	C4-C3-O2-C2
5	A	409	PEG	C1-C2-O2-C3
5	C	408	PEG	C1-C2-O2-C3


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	406	PEG	1	0
5	C	408	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/376 (97%)	-0.01	9 (2%) 59 63	15, 22, 37, 45	5 (1%)
1	B	366/376 (97%)	-0.02	5 (1%) 75 79	13, 20, 36, 50	3 (0%)
1	C	368/376 (97%)	0.35	30 (8%) 11 13	19, 28, 44, 66	2 (0%)
All	All	1102/1128 (97%)	0.11	44 (3%) 38 42	13, 23, 40, 66	10 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	376	HIS	10.7
1	C	49	TYR	10.7
1	B	10	LYS	7.3
1	C	9	LYS	5.7
1	C	337	TYR	5.5
1	C	350	ASN	4.7
1	C	50	ARG	4.4
1	C	375	PRO	4.1
1	C	346	ILE	3.9
1	C	324	GLU	3.5
1	B	49	TYR	3.4
1	C	44	LEU	3.4
1	C	27	GLN	3.4
1	C	340	PRO	3.3
1	A	49[A]	TYR	3.3
1	A	350	ASN	3.3
1	C	28	GLY	3.1
1	A	27	GLN	3.0
1	C	58	HIS	2.8
1	C	368	PHE	2.7
1	C	204	GLU	2.7
1	A	375	PRO	2.6
1	C	86	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	341	ASP	2.6
1	A	72	GLU	2.6
1	A	167	TYR	2.5
1	C	87	GLU	2.4
1	C	240	GLN	2.4
1	C	374	LEU	2.4
1	C	342	PHE	2.3
1	C	348	PRO	2.3
1	C	83	ASP	2.3
1	C	88	TYR	2.3
1	A	86	SER	2.3
1	B	87	GLU	2.2
1	C	90	LYS	2.2
1	C	128	GLN	2.1
1	C	271	ARG	2.1
1	A	50[A]	ARG	2.0
1	B	50	ARG	2.0
1	B	197	VAL	2.0
1	C	347	SER	2.0
1	A	236	GLU	2.0
1	C	26	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	C	404	5/5	0.43	0.35	107,107,107,107	0
3	SO4	A	405	5/5	0.61	0.26	125,125,125,125	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PEG	A	408	7/7	0.62	0.34	60,61,61,62	0
5	PEG	A	410	7/7	0.70	0.18	66,66,66,66	0
3	SO4	A	404	5/5	0.72	0.36	116,116,116,116	0
5	PEG	C	409	7/7	0.72	0.37	64,65,65,65	0
5	PEG	A	409	7/7	0.73	0.23	57,58,58,58	0
5	PEG	B	407	7/7	0.75	0.15	54,55,55,55	0
3	SO4	A	403	5/5	0.83	0.24	33,34,36,37	0
5	PEG	B	406	7/7	0.87	0.16	71,71,71,71	0
5	PEG	C	408	7/7	0.88	0.12	51,51,52,52	0
6	ZLS	C	410	39/39	0.88	0.13	22,32,34,34	0
6	ZLS	A	411	39/39	0.90	0.11	18,26,31,31	0
3	SO4	A	402	5/5	0.91	0.15	70,70,70,70	0
3	SO4	C	403	5/5	0.91	0.26	51,51,52,52	0
3	SO4	C	402	5/5	0.92	0.14	114,114,114,114	0
6	ZLS	B	408	39/39	0.92	0.11	16,26,29,30	0
3	SO4	B	403	5/5	0.92	0.33	90,90,90,90	0
3	SO4	B	402	5/5	0.93	0.18	60,60,60,61	0
3	SO4	C	405	5/5	0.93	0.30	76,77,77,77	0
4	CA	B	405	1/1	0.98	0.11	42,42,42,42	0
2	ZN	C	401	1/1	0.99	0.03	21,21,21,21	0
4	CA	C	406	1/1	0.99	0.23	45,45,45,45	0
4	CA	A	407	1/1	0.99	0.11	45,45,45,45	0
4	CA	B	404	1/1	0.99	0.11	36,36,36,36	0
2	ZN	B	401	1/1	1.00	0.07	15,15,15,15	0
4	CA	C	407	1/1	1.00	0.11	44,44,44,44	0
2	ZN	A	401	1/1	1.00	0.06	16,16,16,16	0
4	CA	A	406	1/1	1.00	0.17	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZLS C 410:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZLS A 411:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.