



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 30, 2023 – 11:58 AM EDT

PDB ID : 3MPL
Title : Crystal Structure of Bordetella pertussis BvgS VFT2 domain (Double Mutant F375E/Q461E)
Authors : Herrou, J.; Bompard, C.; Wintjens, R.; Dupre, E.; Willery, E.; Villeret, V.; Locht, C.; Antoine, R.; Jacob-Dubuisson, F.
Deposited on : 2010-04-27
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Virulence sensor protein bvgS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	233	1799	1137	317	339	6	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	276	MET	-	expression tag	UNP P16575
A	277	ARG	-	expression tag	UNP P16575
A	278	GLY	-	expression tag	UNP P16575
A	279	SER	-	expression tag	UNP P16575
A	280	HIS	-	expression tag	UNP P16575
A	281	HIS	-	expression tag	UNP P16575
A	282	HIS	-	expression tag	UNP P16575
A	283	HIS	-	expression tag	UNP P16575
A	284	HIS	-	expression tag	UNP P16575
A	285	HIS	-	expression tag	UNP P16575
A	286	GLY	-	expression tag	UNP P16575
A	375	GLU	PHE	engineered mutation	UNP P16575
A	461	GLU	GLN	engineered mutation	UNP P16575

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	82	Total O 82 82	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	48.80Å 101.48Å 53.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 5.58 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.2 (20.00-2.10) 100.0 (5.58-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.46 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.188 , 0.245 0.188 , 0.246	Depositor DCC
R_{free} test set	776 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtrriage
Anisotropy	0.004	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.98 , 105.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1889	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	0/1830	0.92	2/2487 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	343	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	A	421	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1799	0	1809	29	0
2	A	8	0	12	1	0
3	A	82	0	0	5	0
All	All	1889	0	1821	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:444:PRO:HB2	1:A:469:TYR:CE2	2.19	0.78
1:A:393:ASN:HD21	1:A:464:ILE:H	1.39	0.67
1:A:398:VAL:HB	1:A:459:VAL:HG13	1.77	0.66
1:A:393:ASN:HD22	1:A:462:THR:HB	1.65	0.61
1:A:292:HIS:C	1:A:292:HIS:CD2	2.75	0.60
1:A:393:ASN:ND2	1:A:464:ILE:H	2.00	0.60
1:A:426:ILE:O	1:A:427:PRO:C	2.41	0.56
1:A:421:ARG:HG2	1:A:441:ALA:O	2.08	0.54
1:A:292:HIS:HE1	1:A:511:TYR:CD2	2.25	0.53
1:A:432:ARG:O	1:A:433:TYR:C	2.43	0.53
2:A:543:EDO:H22	3:A:82:HOH:O	2.09	0.51
1:A:292:HIS:CE1	1:A:511:TYR:HD2	2.29	0.50
1:A:408:ASP:HB2	1:A:411:HIS:CG	2.47	0.50
1:A:292:HIS:CD2	3:A:75:HOH:O	2.65	0.49
1:A:292:HIS:C	1:A:292:HIS:HD2	2.16	0.49
1:A:292:HIS:CE1	1:A:511:TYR:CD2	3.01	0.49
1:A:402:ASP:HB2	1:A:479:ARG:HH22	1.78	0.48
1:A:463:GLN:HE22	1:A:526:ARG:HH11	1.60	0.48
1:A:397:ILE:HG23	1:A:458:ALA:HB1	1.95	0.47
1:A:339:LEU:O	1:A:343:ARG:HG2	2.15	0.46
1:A:292:HIS:HD2	3:A:75:HOH:O	1.97	0.46
1:A:420:VAL:HG23	3:A:56:HOH:O	2.15	0.46
1:A:397:ILE:HD13	1:A:410:ASP:HA	1.99	0.44
1:A:463:GLN:HE22	1:A:526:ARG:NH1	2.14	0.44
1:A:446:GLU:O	1:A:450:MET:HG3	2.16	0.44
1:A:393:ASN:HD21	1:A:464:ILE:N	2.11	0.43
1:A:383:PHE:CD2	1:A:383:PHE:N	2.87	0.42
1:A:359:LEU:C	1:A:359:LEU:HD23	2.39	0.41
1:A:349:GLU:OE1	3:A:79:HOH:O	2.22	0.41
1:A:418:ALA:HA	1:A:439:VAL:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/267 (86%)	223 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/218 (87%)	183 (96%)	7 (4%)	34	35

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	383	PHE
1	A	421	ARG
1	A	423	SER
1	A	449	LEU
1	A	461	GLU
1	A	479	ARG
1	A	485	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	292	HIS
1	A	393	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	1	-	3,3,3	0.40	0	2,2,2	0.23	0
2	EDO	A	543	-	3,3,3	0.44	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1	-	-	0/1/1/1	-
2	EDO	A	543	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	543	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/267 (87%)	-0.02	18 (7%) 13 17	9, 21, 48, 62	3 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	404	ASP	3.9
1	A	432	ARG	3.4
1	A	431	ARG	3.0
1	A	403	PRO	2.8
1	A	435	GLN	2.8
1	A	292	HIS	2.7
1	A	485	ASP	2.7
1	A	382	SER	2.6
1	A	434	PRO	2.6
1	A	402	ASP	2.4
1	A	379	ALA	2.3
1	A	327	GLU	2.2
1	A	473	TYR	2.2
1	A	436	ALA	2.2
1	A	433	TYR	2.1
1	A	414	GLY	2.1
1	A	445	SER	2.0
1	A	383	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	543	4/4	0.92	0.15	37,39,39,39	0
2	EDO	A	1	4/4	0.97	0.07	18,19,20,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.